



LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

546

Il6i

1962-63



L

Return this book on or before the
Latest Date stamped below.

University of Illinois Library

~~APR 3 1 1964~~

~~APR 3 1965~~

~~DEC 15 1965~~

~~DEC 21 1968~~

~~JUL 2 1974~~

L161—H41



Digitized by the Internet Archive
in 2012 with funding from
University of Illinois Urbana-Champaign

<http://archive.org/details/inorganicsemi196263univ>

546
I 262
1962/63

TABLE OF CONTENTS

1962-63

SUMMER SESSION - 1962		Page
MOLECULAR ADDITION COMPOUNDS OF SILVER WITH OLEFIN AND AROMATIC MOLECULES Brad Wayland		1
NUCLEOPHILIC DISPLACEMENTS ON PHOSPHORUS G. D. Shier		8
APPLICATIONS OF GROUP THEORY TO INORGANIC CHEMISTRY Keith F. Purcell		13
RECENT REACTIONS OF THE METAL CARBONYLS L. Jane Park		22
ELECTRONIC CONFIGURATION OF TETRAHEDRAL OXYANIONS OF TRANSITION METALS P. Van Der Voorn		27
THE DEFECT SOLID STATE - AN INTRODUCTORY SURVEY Violet I. Imhof		32
FALL AND SPRING SESSION - 1962-1963		Page
COMPLEX CHEMISTRY OF PHOSPHINES AND PHOSPHINE OXIDES William J. Randall		1
SOME SELECTED USES OF OPTICAL ROTATORY DISPERSION IN COORDINATION CHEMISTRY L. J. Boucher		6
FAR INFRA-RED ABSORPTION OF TRANSITION-METAL COMPLEXES Michael R. Rosenthal		15
A MASS-SPECTROMETRIC STUDY OF SULFUR VAPOR FROM 81°C to 1,737°C. J. R. Marquart		36
STUDIES OF THE HAEMOPROTEINS B. W. Baylis		44
ANTI-FERROMAGNETISM IN SOME FIRST-ROW TRANSITION-METAL FLUORIDES J. C. Schapiro		56
NUCLEAR QUADRUPOLE RESONANCE IN INORGANIC CHEMISTRY Ronald H. Carlson		69



CONFIDENTIAL

SECRET

1. The purpose of this document is to provide a comprehensive overview of the current state of the project and to outline the key objectives for the next phase of development.

2. The project has been initiated in response to the strategic goals set forth in the company's annual report, with a primary focus on enhancing operational efficiency and reducing costs.

3. The initial phase of the project involved a thorough analysis of the existing processes, identifying areas of inefficiency and potential for improvement.

4. Key findings from the analysis include the need for improved communication channels, the implementation of standardized procedures, and the adoption of new technologies to streamline workflows.

5. The proposed solution involves the implementation of a new software platform, the restructuring of the organizational chart, and the establishment of a cross-functional team to oversee the project's progress.

6. The implementation of these changes is expected to result in a 15% increase in productivity and a 10% reduction in operational costs over the next six months.

7. The project team is committed to maintaining open communication and providing regular updates to all stakeholders throughout the implementation process.

8. The next steps include the final approval of the project plan, the allocation of resources, and the commencement of the implementation phase.

9. It is anticipated that the project will be completed by the end of the fiscal year, with a final evaluation to assess the impact of the changes on the organization's overall performance.

10. The success of this project will be a significant milestone for the company, demonstrating our ability to adapt to changing market conditions and drive innovation in our operations.

CONFIDENTIAL

COMPLEXES OF <u>O</u> -PHENYLENEBIS(DIMETHYLARSINE) L. V. Interrante	84
THE ZERO-VALENT COMPLEXES OF PLATINUM L. Jane Park	96
CHELATE STABILITY AND SUBSTITUTION ON AROMATIC LIGANDS E. J. Olszewski	100
CHEMICAL EFFECTS OF NUCLEAR TRANSFORMATIONS IN THE GAS PHASE Agnes Chia Wang	116
RECENT CHEMISTRY OF THE ALANES D. P. Eyman	121
RATE OF A SUBSTITUTION REACTION IN CHLOROFORM George Shier	126
INFRARED STUDIES OF TRIMETHYLTIN COMPOUNDS R. C. Burrows	127
BRIDGE ADDUCTS-THE INTERACTION OF BORON TRIFLUORIDE WITH TRANSITION-METAL CYANIDE COMPLEXES Gerald Feistel	128
N.M.R. COUPLING CONSTANTS AND HYBRIDIZATION IN ORGANO-TIN COMPOUNDS J. C. Schapiro	129
MOLECULAR EXCITATION IN SHOCK TUBES Sue F. Eilers	134
BORON POLYMERS Gerald Tennenhouse	141
MACROCYCLIC COORDINATION COMPOUNDS E. J. Olszewski	145
E. S. R. OF TRANSITION METAL ION COMPLEXES Peter C. Van Der Voorn	151
N. M. R. IN THE PRESENCE OF PARAMAGNETIC IONS Brad Wayland	158
AQUEOUS INFRARED SPECTROSCOPY AND ITS APPLICATIONS TO INORGANIC CHEMISTRY Ronald H. Carlson	172
FLASH PHOTOLYSIS AND KINETIC SPECTROSCOPY Michael R. Rosenthal	181
THE MOSSBAUER EFFECT AND ITS APPLICATIONS TO INORGANIC CHEMISTRY K. F. Purcell	187
HYDROLYSIS OF METAL IONS J. E. House, Jr.	193
MOLECULAR ORBITAL TREATMENT OF POLYHALOGEN AND RELATED SPECIES-203 Carolyn M. McArthur	

MOLECULAR ADDITION COMPOUNDS OF SILVER
WITH OLEFIN AND AROMATIC MOLECULES

Brad Wayland

June 19, 1962

I. INTRODUCTION

The first studies of silver olefin complexes were concerned with the effects of alkyl substituents on the relative stabilities of the complexes. Alkyl substitution in the olefin molecule created opposing electronic and steric effects on the relative complexing ability of the olefin (1). The position of the alkyl group was found to be particularly important. Studies of a series of butene isomers demonstrates that the cis 2-butene complex is much more stable than that of the trans isomer (2). A similar study of pentene isomers showed that the complex of cis-2-pentene was more stable than that of the trans isomer (3).

Olefin	Agentation Constant K_1
1-butene	119.4
2-methyl propene	71.5
cis-2-butene	62.5
trans-2-butene	24.6

The effect of ring size and strain on complexation of Ag^+ has been studied for a series of unsaturated ring compounds (4)(5). The relative equilibrium constants are found to increase as the strain in the ring increases.

Olefin	Relative Equilibrium Constant K_1
cyclopentene	23.6
cycloheptene	4.2
cyclohexene	3.65



The effects of ring strain on an exocyclic double bond have also been studied (5). Formation constants of the methylenecycloalkanes show only a slight dependency on ring size.

Olefin	Relative Equilibrium Constant K_1
methylenecyclobutane	17.9
methylenecyclopentane	13.3
methylenecycloheptane	14.4
methylenecyclohexane	11.9

Silver aromatic complexes are also subject to both steric and electronic effects (6)(7). Thus the substitution of a methyl group

aromatic	argentation constant K_1
benzene	1.10
toluene	1.19
ethyl benzene	.98
t-butylbenzene	.90
o-xylene	1.43
m-xylene	1.35
p-xylene	1.14
m-diethylbenzene	1.06
m-diisopropylbenzene	.93
p-di t-butylbenzene	.62
1,2,3,4 tetramethylbenzene	1.69
1,2,4,5 tetramethylbenzene	.88

onto the benzene ring increases the basicity of the aromatic molecule and the argentation constant increases. However, as the steric requirement of the alkyl group increase from methyl- to t-butyl- the argentation constant decreases. The position of the alkyl groups in the aromatic molecule also affects the formation of the addition compound with silver. Thus 1,2,3,4 tetramethylbenzene has a much larger argentation constant than 1,2,3,5 tetramethylbenzene. The energy of the highest occupied orbital in the aromatic is found by Fukui to be a good measure for complex formation. This indicates that the use of ionization potentials by Nakajima (9) to correlate the relative ease of complex formation is valid.

aromatic	K_1	relative delocal. energy	Energy of the highest occupied orbital
	2.41	1.333 β	-1.000
CH ₃ 	2.95	1.418 β	-.823
CH ₃  CH ₃	3.03	1.466 β	-.763
	3.08	1.600 β	-.618
	3.67	1.798 β	-.605
	3.94	1.370 β	-.705
	6.3	2.139 β	-.504

It is also recognized that polarizability is another factor which varies slightly for each aromatic molecule. Ion induced dipole interactions are not uniform from one aromatic molecule to the next. No correlation between polarizability and complex stability has yet been published, but a paper by Fukui is now being prepared (8).

II. STRUCTURE

The structures of silver olefin complexes are not well defined. It is generally assumed that the silver ion is symmetrically located above the plane of the olefin in the π cloud (10)(11). The suggested structure for the norbornadiene silver complexes places the silver ion in an exo configuration (12). This conforms with the model suggested by Winstein and Shatavsky for the electrophillic attack Br_2 on an olefin (13).



1:1 complex



1:2 complex

The structure of the silver aromatic complexes are better understood. An x-ray analysis of the silver benzene complex by Rundle showed that silver ions lie above and below the benzene ring over bonds one and four of each ring (14)(15). Thus silver benzene is an infinite polymer. A recent Raman spectra study has given similar results (16).




silver benzene complex

o carbon
 O Ag^+
 NO_3^- is excluded

For the general case of any aromatic, Mulliken suggests that the silver ion will coordinate preferentially above and between two adjacent unsubstituted ring carbons when these positions are available (17). This model for the silver aromatic structure explains why there are significant stability differences between the complexes formed by ortho, meta and para xylene (6)(7).


The structure of silver cyclooctatetraene has been extensively studied by x-ray analysis (18-21). Silver ion is found to interact with two nonadjacent π bonds of the C_8H_8 molecule at $\text{Ag}-\text{C}$ distances

of 2.46, 2.51, 2.78 and 2.84 Å. Somewhat longer Ag-C distances join the monomer unit into infinite chains along the c axis.




III. BONDING

The bonding in silver olefins was at first considered to be essentially the sharing of a pair of π electrons of the olefin with the Ag^+ (21)(23). This model was quickly discarded for free rotation would be allowed in the complexed olefin, and no rearrangement of cis and trans isomers was observed upon complexation (24). Winstein and Lucas suggested that the silver olefin bond involved three resonance forms (1).




Similar metal to carbon σ -bonding has been proposed by Babushkin, Gribov and Hellman (25)(26). Recent N.M.R. and I.R. studies by Powell and Sheppard indicate that the olefin is changed only slightly upon complexation (27). In terms of the Winstein and Lucas model, the no bond structure $\left(\begin{array}{c} \text{C} = \text{C} \\ \text{Ag} \\ + \end{array} \right)$ must predominate.

A more satisfying model for the silver olefin bond was suggested by Dewar (10).



The vacant 5s orbital on silver ion is capable of overlapping the bonding π orbital on the olefin to form a σ bond. The filled 4d orbitals on the silver ion have the proper symmetry to overlap the vacant antibonding molecular orbitals of the olefin. The combination of the two oppositely directed bonds leaves the olefin with a much smaller positive charge than would be the case in a normal π complex. If the silver ion is situated in the center of the π cloud in the ethylene silver complex, then the complex has C_{2v} symmetry. The molecular orbitals formed by the overlap of the silver 5s orbital with the π bonding orbital of ethylene both have A_1 symmetry. The electric dipole operator in the Z direction is also A_1 . Therefore, charge transfer from the $\sigma \rightarrow \sigma^*$ is allowed. This suggests that the most probable position for the silver ion in the complex is directly above the carbon carbon double bond.



Traynham and Sehert contend that little or no σ bond character can be ascribed to the silver olefin complex since the complex is readily dissociated and no isomerization or polymerization of the olefin is observed (4). Traynham and Sehert suggest that it is adequate to describe the silver olefin complex as a π complex in which only the deformation of the π orbitals on the olefin is significant.

The bonding in silver aromatic complexes can be described in much the same manner as that for the silver olefin complexes, however, the highest filled π molecular orbital of benzene is not of the proper symmetry to give charge transfer if the silver ion is placed on the benzene six fold axis (28)(17). The highest filled π molecular orbital of benzene belongs to the irreducible representation e_1 , while the lowest acceptor orbital of silver has A_1 symmetry. The silver ion moves to a position of lower symmetry where charge transfer becomes allowed. The empty e_2 π orbitals overlap the silver d orbitals which partially balances the charge transfer (29).

Rundle and Smith observed that the benzene molecule in the silver benzene complex is significantly distorted (14)(15). The carbon-carbon bond nearest to the silver ion becomes shorter. This distortion can only be explained as due to the polarization of the π electrons by silver increasing the electron density in the bonds nearest the positive silver ion. Thus polarization terms must also make a significant contribution to the total bonding.

Origin of Steric Effects

Cis olefins in general form much more stable complexes with silver than the trans isomer (2)(3). If the silver ion lies above the plane of the olefin in the π cloud, then steric effects should be small. Cis olefins, having two alkyl groups on one side of the molecule, are strained due to bond oppositions. The strain is partially relieved by a slight twisting about the double bond. The twisting about the double bond deforms the π orbital which, according to Gardner, facilitates polarization of the π electrons and thus stabilizes the complex (11). The effects of ring strain on cyclic olefins may be similarly explained. A. Cope reported that trans-cyclooctene preferentially reacts with silver and may be quantitatively separated from the cis isomer (30). This appears to be an anomalous result. However, a conformational analysis of cyclooctene shows that the trans isomer is more strained than the cis isomer. Thus the π electron cloud of the trans is more easily polarized and a stronger bond results

Applications of Silver Olefin Complexation

The stability differences between cis and trans olefin complexes with silver have been used by the organic chemist to separate geometrical isomers (30)(34). Silver ion has also been used as a means of trapping and stabilizing the short lived cyclobutadiene molecule(35). Complex formation may possibly be used as a means of trapping other short lived olefin intermediates in organic reactions.

BIBLIOGRAPHY

1. S. Winstein and H. Lucas, J. Am. Chem. Soc., 60, 836 (1938).
2. F. Hepner, K. Trueblood and H. Lucas, *ibid.*, 74, 1333 (1952).
3. H. Lucas, R. Moore, and D. Pressman, *ibid.*, 65, 227 (1943).
4. J. Traynham and M. Sehert, *ibid.*, 78, 4024 (1956).
5. J. Traynham and J. Olechowski, *ibid.*, 81, 571 (1959).
6. N. Ogimachi, J. Andrews, and R. Keefer, *ibid.*, 78, 2210 (1956).
7. J. Andrews and R. Keefer, *ibid.*, 71, 3644 (1949).
8. K. Fukui, A. Imamura, F. Yonezawa, C. Nagata, Bull. Chem. Japan,
9. F. Nakajima, The Science Reports of the Fohohu University, Ser. I., 41, 171 (1957).
10. M. Dewar, Bull. soc. chim., 18, C79 (1951).
11. P. Gardner, R. Brandon, and N. Nix, Chem. and Ind., 1958, 1363.
12. J. Traynham, J. Org. Chem., 26, 4694 (1961).
13. S. Winstein and M. Shatavsky, Chem. and Ind. 1956, 56.
14. R. Rundle and J. Goring, J. Am. Chem. Soc., 72, 5337 (1950).
15. H. Smith and R. Rundle, *ibid.*, 80, 5075 (1958).
16. Sh. Roskin, Doklady Acad. Nauk. S. S. S. R., 141, No. 4, 900 (1961).
17. R. Mulliken, J. Am. Chem. Soc., 74, 821 (1952).
18. F. Mathews and W. Lipscomb, J. Phys. Chem. 63, 845 (1959).
19. F. Mathews, Dissertation Abstr., 20, 3971 (1960).
20. S. Nyburg and J. Hilton, Acta. Cryst., 12, 116 (1959).
21. F. Mathews and W. Lipscomb, J. Am. Chem. Soc., 80, 4745 (1958).
22. G. Bennett and G. Willis, J. Chem. Soc., 1929 259.

23. H. Drew, F. Pinkard, W. Wardlaw and E. Cox, *ibid.*, 1932, 988.
24. J. Anderson, *ibid.*, 1936, 1042.
25. A. Babashkin, L. Gribov and A. Hellman, *Doklady Acad. Sci. U. R. R. S.*, 12346 (1958).
26. A. Hellman, L. Gribov, and A. Babashkin, *Chem. Soc. Spec. Publ.*, No. 13, 116 (1959).
27. D. Powell and N. Sheppard, *J. Chem. Soc.*, 1960, 2519.
28. R. Rundle and J. Corbett, *J. Am. Chem. Soc.*, 79, 757 (1957).
29. D. Brown, *J. Inorg. Nucl. Chem.*, 10, 39 (1959).
30. A. Cope, R. A. Pike, and C. Spencer, *J. Am. Chem. Soc.*, 75, 3212 (1953).
31. Nichols, *ibid.*, 74, 1091 (1952).
32. W. Jones, *J. Chem. Soc.*, 1954, 1808.
33. A. Romyns and J. Lucas, *J. Am. Chem. Soc.*, 79, 4339 (1957).
34. L. Zakharkin and V. Korneva, *Doklady Akad. Nauk S. S. S. R.*, 132, 1078 (1960).
35. W. Avram, E. Marica and C. Nenitzescu, *Chem. Ber.*, 92, 1088 (1959).

NUCLEOPHILIC DISPLACEMENTS ON PHOSPHORUS

G. D. Shier

June 26, 1962

I. INTRODUCTION

The reactions of phosphorus compounds are interesting because of their analogy to carbon chemistry, their importance in living systems, and the availability of "d" orbitals for bonding. The existence of five coordinate phosphorus in PCl_5 and pentaphenyl phosphorus raises the possibility of metastable five coordinate intermediates in displacement reactions, and this possibility has been explored. This seminar will deal with some of the less complicated and more intensively studied reactions of phosphorus compounds and will try to interpret them in terms of kinetic and mechanistic parameters.

II. THE REACTIONS OF ACID FLUORIDES

The most thoroughly investigated class of compounds is the R_2POF series where R can be an alkyl, alkoxy, or amide group. Aksnes(1) has studied the effect of alkoxy versus alkyl substitution in the bimolecular alkaline hydrolysis reaction of R_2POF . His data are given in Table I. It is apparent that the activation energy increases and the rate decrease with the replacement of alkyl groups by alkoxy groups.

The inductive effect of oxygen in a sigma bond to phosphorus is more strongly electron withdrawing than for a bond between carbon and phosphorus, and this decreases the shielding of the phosphorus nuclear charge in the case of alkoxy substituents. However, as the phosphorus nucleus becomes less shielded, conjugation between the electron pairs on oxygen and the phosphorus atom becomes more favorable. On this basis a decrease in shielding of phosphorus will facilitate bond making and retard bond breaking. An increase in the conjugation effect will retard bond making, reducing the residual positive charge on phosphorus seen by an incoming nucleophile. The effect of conjugation on bond breaking will depend on the ability of the leaving group to strengthen its bond to phosphorus by conjugation. Implicit in this argument is the assumption of a five coordinate transition state.

Halmann(2) has investigated the hydrolysis of R_2POF where R is a methyl or isopropoxy group. He found no ^{18}O exchange for the phosphoryl group during the reaction, and general acid-base catalysis. The lack of ^{18}O oxygen exchange is interpreted as eliminating fast reversible addition of water in the transition state, but reversibility implies equivalence of the oxygen of the attacking group and of the phosphoryl group, which is questionable. Base catalysis may be due to greater rate of attack of hydroxide as compared to water, and acid catalysis may be due to protonation of the leaving group. Acid catalysis has not been observed where chloride is the leaving group, and hydrogen bonding is much less effective for chloride than for fluoride.

The effect of the calculated entropy of activation in Table I is to oppose the effect of the activation energy. Aksnes(3) has used the shift of the infrared stretching frequency of the phosphoryl group upon hydrogen bonding with phenol as a measure of the polarizability of the phosphorus compound. This work indicates that alkoxy groups enhance polarizability. On this basis the entropy effect can be explained by greater delocalization of charge in transition state and consequently greater desolvation of the transition complex. Desolvation causes an increase in entropy giving a less negative overall entropy of activation. Steric effects often appear in entropy terms, and so it must be assumed that the steric effects of the ethyl and methoxy groups are similar.

Heath(4) has worked on the kinetics of alkaline hydrolysis of R_2POF compounds where R is a primary or secondary alkyl amido group. His data is neither good enough nor complete enough to separate the various steric, inductive, and conjugative factors, but it does indicate great steric hindrance for compounds where both R groups are secondary amido groups. Models are said to show these compounds to be very rigid.

The reactions of R_2POF compounds have been studied by other workers with varying results due to the complexity of the effects. The rates of reaction with oximes(5) and hydroxamic acids(6) have been found to correlate well with the basicities of the nucleophiles. Larsson has studied the alkaline hydrolysis of alkyl methylphosphonofluoridates(7), and the reaction of these compounds with hydrogen peroxide(8). Epstein et al. (9) have found that catechol reacts faster than phenol with alkyl methylphosphonofluoridates due to hydrogen bonding in the transition state.

III. THE REACTIONS OF ACID CHLORIDES

Dostrovsky and Halmann(10) studied the solvolysis, aminolysis, and reactions of anions with phosphorochloridates. For the reactions with anions they found the order $F^- > EtO^- > C_6H_5O^- > EtOH, C_6H_5S^-, acetate$. The reactivity of fluoride ion was about six times slower in 60% aqueous ethyl alcohol than in dry alcohol. This is possibly due to stronger hydrogen bonding between fluoride ion and water as compared to ethyl alcohol.

Aminolysis of diisopropyl phosphonochloridate with di-normalbutyl amine in several solvents showed a rather interesting behavior. The rate of the reaction increased smoothly with the increase in dielectric constant of the solvent for nonhydroxylic solvents, but the rate in ethanol with dielectric constant 26 was between benzene and dioxane with dielectric constant 2.3. This is especially interesting in view of the fact that almost all other work has been done in hydroxylic solvents.

Miller(12) has reacted O,O-diphenyl phosphorochloridothioate with a series of oxygen and sulfur nucleophiles and found that the plot of pK_a of the nucleophile versus the negative logarithm of the k_a rate constant for ten nucleophiles gives a curved line with little scatter. The anions of benzophenone and acetone oximes are much more reactive than their basicities would indicate, and a specific mechanism has been proposed for them.

A thorough discussion of the hydrolysis of phosphonochloridates(13) has been given, the conclusions being similar to those for the phosphonofluoridates.

IV. OTHER DISPLACEMENT REACTIONS

Optically active o-ethyl ethylphosphonothioic acid has been isolated (14) and used to study the stereospecificity of displacements on phosphorus. Displacement reactions carried out using hydrogen sulfide, n-propyl mercaptide, and ethoxide ions in ethanolic solution indicated that inversion of configuration had taken place. A preliminary communication on other similar work has appeared (15).

The kinetics of alkaline hydrolysis of phosphonate esters has been studied(16) and found to proceed by SN_2 attack of hydroxide on phosphorus. The rates of hydrolysis decrease smoothly with increasing size of the alkyl group, but the activation energy and preexponential factors show no regular pattern.

The reaction of optically active methylethylphenylbenzylphosphonium iodide with hydroxide and alkoxide ions has been studied (17, 18). The reaction with hydroxide gives optically pure methylethylphenylphosphine oxide. The authors say that the hydroxide ion can add reversibly to any face of the tetrahedron, and the conjugate base of any of the trigonal bipyramidal intermediates can be formed by reversible reaction with another hydroxide ion. Since the benzyl group forms the most stable anion, the only conjugate base that decomposes is the one where the oxygen atom and the benzyl group are colinear.

V. SUMMARY AND CONCLUSIONS

It has not been found necessary to postulate the existence of a stable five-coordinate intermediate in any of the displacement reactions on phosphorus that have been studied. The evidence favors a straight forward SN_2 reaction mechanism where bond making and bond breaking influence the rate to varying degrees. Some authors(12) cite the strong rate dependence on the basicity of the nucleophile as evidence for bond making as the rate determining step. This hypothesis has not been tested with a variety of leaving groups or in non hydroxylic solvents. The strong phosphorus fluorine bond probably accounts partially for the slower reactions

and higher activation energies of phosphorofluoridates as compared to phosphorochloridates. The existence of a rapid reversible equilibrium step preceding displacement has not been demonstrated in any case, and it seems better to assume a straight forward nucleophilic attack. Steric effects on rate constants are analogous to those found in carbon chemistry.

Much work remains to be done in the area of organo-phosphorous mechanisms. In particular, it would seem desirable to know more about conjugative versus inductive effects, solvation effects, and the effect of variation of the leaving group.

References

1. G. Aksnes: Acta. Chem. Scand., 14, 1515 (1960).
2. M. Halmann: J. Chem. Soc., 305 (1959).
3. G. Aksnes and T. Gramstad: Acta. Chem. Scand., 14 1485(1960).
4. D. F. Heath: J. Chem. Soc., 3796, 3804 (1956).
5. A. L. Green and B. Saville: J. Chem. Soc., 3887 (1956).
6. A. L. Green, G. L. Sainsburg, B. Saville, and M. Stansfield: J. Chem. Soc., 1583 (1958).
7. L. Larsson: Acta. Chem. Scand. 11, 1131 (1957).
8. L. Larsson: Acta. Chem. Scand. 12, 723 (1958).
9. J. Epstein, D. H. Rosenblatt, M. M. Demek: J. Am. Chem. Soc., 78, 341 (1956).
10. I. Dostrovsky and M. Halmann: J. Chem. Soc., 502, 508, 511, 516 (1953).
11. I. Dostrovsky and M. Halmann: J. Chem. Soc., 1004 (1956).
12. B. Miller: J. Am. Chem. Soc., 84, 403 (1962).
13. R. F. Hudson and L. Keay: J. Chem. Soc., 1859, 1865 (1960).
14. H. S. Aaron, R. T. Uyeda, H. F. Frack, and J. I. Miller: J. Am. Chem. Soc., 84, 617 (1962).
15. M. Green and R. F. Hudson: Proc. Chem. Soc., 227 (1959).
16. R. F. Hudson and L. Keay: J. Chem. Soc., 2463 (1956).
17. A. Blade-Font, C. A. Vander Werf, and W. E. McEwen: J. Am. Chem. Soc., 82, 2396 (1960).
18. C. B. Parisek, W. E. McEwen, and C. A. Vander Werf: *ibid.*, 82, 5503 (1960).

Table I

<u>Compound</u>	<u>K(hydroxide)</u>	\bar{E}_A	\bar{S}^I	<u>I.R. Stretch(cm^{-1})</u> P=0	<u>P-F</u>	<u>Phenol Association</u> at 20° in CCl_4
$(\text{MeO})_2\text{POF}$	18.2	11.6	-16	1308	864	26.3
$(\text{MeO})\text{EtPOF}$	49.1	10.1	-19	1295	858	82.5
Et_2POF	726	8.3	-20	1278	845	129.9

APPLICATIONS OF GROUP THEORY TO INORGANIC CHEMISTRY

Keith F. Purcell

July 12, 1962

INTRODUCTION


Group theory can be applied quite easily to many problems in inorganic chemistry and through an understanding of a few basic principles, one may attain a great deal of insight into otherwise incomprehensible phenomena. It is the purpose of this seminar to demonstrate the success of group theory and the ease with which it may be applied to inorganic problems. The mathematical foundations of the theory will be taken for granted but they can be found in several texts (1,2,3,4,5).

STRUCTURE DETERMINATIONS

Many molecular structures have been ascertained or at least reduced to a few possibilities by application of group theory to the infrared and Raman spectra of the compounds. To illustrate the procedure of making a structure assignment, the investigations of Lord, et. al., (6) into the structure of IF_5 will be reviewed.

The first step in this process is to write down all the possible spatial arrangements of the atoms in the molecule. For each structure we determine the point group to which it belongs on the basis of whatever symmetry elements the configuration possesses. Finally, the number of infrared active and Raman active vibrational modes for each point group are determined. Through a comparison of the theoretical with the observed number of frequencies we can limit the number of possible structures or in some cases conclude the actual structure.

For IF_5 there are seven possible symmetries but only one will be discussed in detail. If the five fluorine atoms are arranged in a square based pyramid with the iodine atom at the center of the square, the point group which describes this structure is C_{4v} .



Utilizing the displacement coordinate approach, the motions of each atom are defined in terms of three vectors in the x, y, and z directions (displacement coordinates). The reducible representation generated by the displacement coordinates is found by performing each of the symmetry operations of the group on the molecule and noting which vectors are taken into themselves or their negatives. Each vector taken into itself contributes +1 to the character, into its negative, -1, and all others, 0. The sum (character) is then placed in the column of the symmetry operation.

The reducible representation is decomposed into the irreducible representations which it contains by means of the following equation,

$$n^{(\lambda)} = \frac{1}{h} \sum_j h_j \chi_j^{(\lambda)} \chi_j$$

h = number of operations in the group

h_j = number of operations in the j th set of operations

$\chi_j^{(\lambda)}$ = the character of the λ th irreducible representation for the j th operation

χ_j = the character of the reducible representation for the j th operation

$n^{(\lambda)}$ = the number of times the λ th irreducible representation is contained in the reducible representation.

The normal modes of motion of the molecule are represented analytically by normal coordinates, one coordinate for each mode. It can be shown that each normal coordinate generates an irreducible representation of the point group. The symmetries of the normal modes of motion are finally arrived at by subtracting out the symmetries of the translational and rotational degrees of freedom. (See last page). As a final check, the final number of symmetry species should equal $3N-6$ (for non-linear molecules).

$$\begin{aligned} \Gamma(\text{motion}) &= 4A_1 + A_2 + 2B_1 + B_2 + 5E \\ \Gamma(\text{vib.}) &= 3A_1 + 2B_1 + B_2 + 3E \end{aligned}$$

Now that the symmetries of the normal modes are known, the optically active modes are determined from the fact that only those transitions are allowed for which the transition integral

$$\int \psi_x |R| \psi_y d\tau \neq 0$$

This is equivalent to saying that the direct product of the irreducible representations corresponding to each function in the integral must contain the totally symmetric irreducible representation of the point group.

Since the ground state vibrational symmetry of any molecule is the totally symmetric symmetry, only those modes whose symmetries are the same as any of the "T"s are infrared active. Those modes whose symmetries correspond to any of the "α"s are Raman active. In addition, any totally symmetric modes are Raman active and polarized. With these rules in mind we can compare the number of Raman lines, the number of infrared lines, polarized lines and coincident lines (total and polarized) found by experiment with the number predicted for each point group and eliminate unlikely symmetries. This analysis led Lord, et. al., to the conclusion that IF_5 possesses C_{4v} symmetry.

HYBRID ORBITAL FORMATION

Again we will use IF_5 as an example. To determine the iodine atomic orbitals which will combine to give hybrid orbitals of the proper symmetry, five labelled lines (representing sigma bonds) are drawn from the iodine atom to the fluorine atoms. These bonds are the basis for a reducible representation as before. (See last page.)

$$\Gamma(\text{hybrid}) = 2A_1 + B_1 + E$$



Possible combinations with these symmetries are

sd^4	sdp^3
sd^2p^2	pd^4
spd^3	d^2p^3


Since the valence shell configuration for iodine is $5s^2 5p^5 5d^0$, we can predict that the combination of atomic functions with the lowest "hybridization energy" will be the correct one and, in this case, d^2p^3 is the most likely candidate. It is interesting to note that Kimball (7) predicted the symmetry of IF_5 to be C_{4v} on the basis of this method.

It is also interesting to note that, as one would anticipate, the d_{z^2} orbital is involved in the axial bond while the $d_{x^2-y^2}$ is involved in formation of the equatorial bonds.

MOLECULAR ORBITAL SCHEMES OF COMPLEXES

In constructing a molecular orbital scheme due care must be taken to "match" ligand and metal orbitals of the same symmetry. The $NiCl_4^{2-}$ ion has been chosen to demonstrate the method. This ion has been shown to possess nearly, if not exactly, T_d symmetry (8). As in the previous section, we first draw the ion with "labelled bonds" and generate a reducible


representation with these, this time for the ligand orbitals. From the four Cl^- sigma bonds we can expect four molecular or symmetry orbitals. (See last page)



$$(\text{4 orb.}) = A_1 + T_2$$

(symmetries of ligand molecular orbitals)

From the character table it is seen that the Ni(II) d orbitals possess E (d_{z^2} , $d_{x^2-y^2}$) and T_2 (d_{xy} , d_{xz} , d_{yz}) symmetries. The s orbital is always totally symmetric. Hence, ligand and metal orbitals of the same symmetry are joined in the diagram and the electrons added (9). Note that this procedure, which is identical to that of the last section, yields the well known sp^3 hybridization for tetrahedral transition ion complexes.



CRYSTAL FIELD THEORY

Since the maximum M_L for Ni(II) (d^8) is three and there are two unpaired electrons, the ground state term symbol for the free ion is $3F$. Due to electronic repulsions there is a $3P$ level just above the $3F$ level. Here, the splitting of these states by a crystal field will be considered and then further splitting by spin-orbit coupling will be discussed.

The angular portions of the free ion wave functions are spherical harmonics, and, as such, generate the irreducible representations of the full rotation group. The behavior of a member of this group under various symmetry operations of the finite groups is summarized in the following table (10).

$$\chi^L(E) = (2L + 1)$$

L	C_2	$\chi(C_2)$	L	C_3	$\chi(C_3)$	L	C_4	$\chi(C_4)$
0		1	0		1	0		1
1		-1	1		0	1		1
2		1	2		-1	2		-1
3		-1	3		1	3		-1


The Ni(II) ion in our example is subjected to a T_d field and hence the free ion terms are split. A reducible representation is generated under the T_d character table using the above tables. When decomposed, this reducible representation yields the symmetries of the various energy levels which have arisen as a result of the tetrahedral field. (See last page)

$$\begin{aligned} \Gamma(3F) &= A_2 + T_1 + T_2 \\ \Gamma(3P) &= T_1 \end{aligned}$$

To determine the effect of spin-orbit coupling on these levels, the symmetry of the spin function must be determined. A theorem, valid for integral spins, states that $\Gamma(S = n) = \Gamma(L = n)$. Therefore, the spin symmetry is found in the same manner as the orbital symmetry. For Ni(II), $S = 1$. The spin-orbit splitting is obtained by forming the direct product and decomposing the reducible representation into the (spin * orbit) irreducible representations. (See last page)

$$\begin{aligned} \Gamma(S = 1) \times \Gamma A_2 &= T_2 \\ " \times \Gamma T_1 &= A_1 + E + T_1 + T_2 \\ " \times \Gamma T_2 &= A_1 + E + T_1 + T_2 \end{aligned} \quad \begin{aligned} & \\ & \text{(As a check, the number} \\ & \text{of new levels must} \\ & \text{equal } (2L + 1)(2S + 1) \end{aligned}$$

It is not possible, at this point, to tell what the order of energy levels is. These must be calculated.



As was discussed in the infrared section, only transitions are allowed which correspond to the transition integral containing the totally symmetric representation. From these considerations, where $R = T_x + T_y + T_z$, three transitions are allowed. The effect of spin-orbit coupling is to split these three bands.

Cotton, et. al., (11) have studied the spectrum of $NiCl_4^{2-}$ and found the ν_2 and ν_3 bands in the visible region of the spectrum. The ν_3 band is quite obviously split (by s-o coupling) into at least four peaks. Liehr and Ballhausen (12) have constructed the E vs. Dq diagram for $T_d Ni(II)$ and Cotton's work agrees very well with theory.

C_{4v}	E	$2C_4$	C_2	2	v	2	d
A_1	1	1	1	1	1	1	$T_z, \alpha_{zz}, \alpha_{xx}, \alpha_{yy}$
A_2	1	1	1	-1	-1	-1	R_z
B_1	1	-1	1	1	-1	-1	$\alpha_{xx} - \alpha_{yy}$
B_2	1	-1	1	-1	1	1	α_{xy}
E	2	0	-2	0	0	0	$(\alpha_{xz}, \alpha_{yz})(T_x, T_y)$ (R_x, R_y)

For Normal Modes

\square (motion)	18	2	z-2	4	2
--------------------	----	---	-----	---	---

For Hybrids

\square (hyb.)	5	1	1	3	1
------------------	---	---	---	---	---

T_d	E	$8C_3$	$3C_2$	(ixC_2) $6C_2$	(ixC_4) $6C_4$	
A_1	1	1	1	1	1	$\alpha_{xx} + \alpha_{yy} + \alpha_{zz}$
A_2	1	1	1	-1	-1	
E	2	-1	2	0	0	$\alpha_{xx} - \alpha_{yy}, \alpha_{xx} + \alpha_{yy} - 2\alpha_{zz}$
T_1	3	0	-1	-1	1	(T_x, T_y, T_z)
T_2	3	0	-1	1	-1	(R_x, R_y, R_z) $(\alpha_{xy}, \alpha_{xz}, \alpha_{yz})$

For Ligand Molecular Orbitals

Γ (orb.)	4	1	0	2	0
--------------------	---	---	---	---	---

For T_d Field Splitting

Γ (3F)	7	1	-1	-1	-1
Γ (3P)	3	0	-1	-1	1

For Spin-Orbit Coupling

Γ (S=1)	3	0	-1	-1	1
" $\times A_2$	3	0	-1	1	-1
" $\times T_1$	9	0	1	1	1
" $\times T_2$	9	0	1	-1	-1

1. Eyring, H., Walter, J., and Kimble, G. E., Quantum Chemistry, John Wiley and Sons, New York (1944), Chap. X and Appendix.
2. Wilson, E. B., Jr., Decius, J. C., and Cross P. C., Molecular Vibrations, McGraw-Hill Book Company, Inc., New York (1955), Chaps. 5-7, 10.
3. Streitwieser, A., Jr., Molecular Orbital Theory for Organic Chemists, John Wiley and Sons, Inc., New York (1961), Chap. 3.
4. Landau, L. D., and Lifshitz, E.M., Quantum Mechanics, Pergamon Press, London (1958), pp. 314-352.
5. Margenau, H., and Murphy, G.M., The Mathematics of Physics and Chemistry, D. Van Nostrand Company, Inc., New York (1943) , pp. 526-563.
6. Lord, R.C., Lynch, M.A., Jr., Schumb, W. C., and Slowinski, E. J., Jr., J. Am. Chem. Soc., 72, 522 (1950).
7. Kimball, G. E., J. Chem. Phys., 8, 188 (1940).
8. Gill, N. S., and Nyholm, R.S., J. Chem. Soc., 3997 (1959).
9. Ballhausen, C. J., and Liehr, A.D., J. Mol. Spec., 2, 342 (1958).
10. Wigner, E. P., (Trans. by J. J. Griffin), Group Theory, Academic Press, New York (1959), Eqn. 15.7, p. 155.
11. Goodgame, D. M. L., Goodgame, M., and Cotton, F. A., J. Am. Chem. Soc., 83, 4161 (1961).
12. Liehr, A. D., and Ballhausen, C. J., Ann. Phys. (New York), 6, 134 (1959).

MOLECULAR SPECTROSCOPY

Eyring, H., Walter, J., and Kimball, G. E., op. cit., Chap. XIV.

Wilson, E. B., Jr., Decius, J. C., and Cross, P. C., op. cit.

Orgel, L. E., Inorg. Chem., 1, 25 (1962).

MOLECULAR ORBITAL THEORY

Streitwieser, A., Jr., op. cit.

Daudel, R., Lefebvre, R., Moser, C., Quantum Chemistry, Interscience Publishers, Inc., New York (1959) pp. 331-354.

Ballhausen, C. J., and Gray, H. B., Inorg. Chem. 1, 111 (1962)

Ballhausen, C. J., and Liehr, A. D., J. Mol. Spec., 2, 342(1958).

CRYSTAL FIELD THEORY

Manch, W., and Fernilius, W. C., *J. Chem. Educ.*, 38, 192 (1961).

Orgel, L. E., *An Introduction to Transition-Metal Chemistry*,
Methuen and Co. Ltd., London (1961), Chap. 6.

Ballhausen, C. J., and Gray, H. B., *op. cit.*

Meek, D. W., Drago, R. S., and Piper, T. S., *Inorg. Chem.*,
1, 285 (1962).

RECENT REACTIONS OF THE METAL CARBONYLS

L. Jane Park

July 17, 1962

In the past two years, developments in the area of metal carbonyls have been quite numerous. The carbonyls are unique for two reasons: first, they are restricted to the transition metals and second, they show a pronounced preference for the low valence states. The object of this seminar is to present a brief review of some of the recent work on carbonyls.

Bonding in the carbonyls involves the overlap of the d, as well as S and P, orbitals of the metal with the π -orbitals of ligands. However, the stability of the carbonyls depends upon back donation from the metals d shell to the ligand. Thus if the carbon monoxide molecule may be written as $C^- \equiv O^+$, complexation with a metal produces $M^- - C \equiv O^+$, and back donation yields $M=C=O$. The mode of bonding thus indicates the reason for restriction of the carbonyls to the transition metals.

The carbonyls are so varied that it is difficult to classify them. Classification of reactions in this seminar is on the basis of the entering ligand in a given reaction.

I. Hydrocarbons

The added stability of the "mixed" compounds and the preparative techniques involving the hydrocarbons substituted carbonyls are important reasons for the rapid accumulation of data in this area. Both unsaturated ring systems and simple olefins are quite common as ligands in substituted metal carbonyls.

A. Ring Systems

Tropylium tricarbonyl chromium complexes have been studied by Munro and Pauson. In addition to so-called "normal" reactions of this complex (1), they have observed a ring contraction (2) and an intermolecular coupling of the ligand of the complex ion (3). Other tropylium complexes include tricarbonyl tropylium vanadium, in which vanadium formally has an oxidation state of (-1) (4).

Substitution of the sulfur atom by an $Fe(CO)_3$ group occurs when $Fe_3(CO)_{12}$ is reacted with thiophene (5). The process is believed to be related to the mechanism of the metal desulfurization process in organic reactions. A structure is proposed for the product.

Various resonance structures may be written for the biphenylene molecule, one of which includes a cyclobutadiene derivative. However, it has been found that only the six membered rings are utilized in π -bonded complexes (6). When two metal-carbonyl groups are bonded to biphenylene, the resulting structure is believed to be trans, from x-ray studies.

Iron pentacarbonyl has the unusual property that when reacted with 1,5 cyclooctadiene, no organo-metallic compound is formed but rather conjugation of the diene is observed (7). Other non-conjugated dienes have been found to form complexes in which the dienes are conjugated.

B. Simple Olefins

Manganese, cobalt, nickel and palladium carbonyls containing an allyl group have been studied (8). Since the atoms in a π -connected allyl group are found to be coplanar, there are believed to be two isomers possible with terminally substituted allyl groups. The allyls seem to be more thermally stable than alkyl and aryl derivatives of the metals.

Other workers have studied the protonation of allyl-type complexes, including those of iron, cobalt and manganese with HCl (9). In the case of π -cyclopentadienyl iron dicarbonyl sigma propene (10), addition of the proton causes a unique rearrangement to a cation containing a π -bonded propene group. The reaction is irreversible.

A compound analogous to the above mentioned ones is formed when $\text{Mo}(\text{CO})_6$ is reacted with acrylonitrile (11). Analysis data are consistent with the formula $(\text{CH}_2\text{CHCN})_2\text{Mo}(\text{CO})_2$. A structure has been formulated but is in question because it does not account for the observed insolubility of the product.

Reaction of $\text{Co}_2(\text{CO})_8$ with acetylenes followed by acid treatment produces a compound whose stoichiometry is $[\text{Co}_3(\text{CO})_9(\text{C}_2\text{H}_2\text{R})]$. A structure is proposed which is consistent with infra-red data. The same compound results when $\text{Co}_2(\text{CO})_8$ is reacted with tri halogeno methyl derivatives (12).

II. Carbon Monoxide

Much interest has been aroused by the carbonylation reaction because the synthesis of aldehydes, acetals (13) and alcohols (14) are believed to be catalyzed by a cobalt complex formed by insertion of a carbon monoxide group into the coordination sphere. For example, reaction between carbon monoxide, hydrogen and ethyl ortho formate produces 1,1 diethoxy-propane (13). The inserted carbon monoxide apparently becomes part of an aldehyde which then reacts with a molecule of the ortho ester to form an acetal.

Reaction between cobalt carbonyl hydride and olefins produces an alcohol, one of the intermediate steps being carbonylation of an alkyl cobalt tetra carbonyl complex (14). Addition of triphenyl phosphine at 0°C allows isolation of the acyl intermediate.

Reduction of the silver amine complexes with carbon monoxide is believed to proceed by a carbonylation-type mechanism, in which a carbon monoxide group is inserted into the coordination sphere. The rate law is found to depend upon the concentration of carbon monoxide (15).

Bonding is believed to exist between two different transition metals in a number of carbonyl derivatives (28,29). The first one of these to be observed has the formula $C_5H_5MoC_5H_5W(CO)_6$. Structures are proposed for these substances.

Another type of metal-metal bonding is believed to occur when $Fe(CO)_5$ is reacted with organo-tin compounds (30). The proposed structure contains no tin-tin or iron-iron bonds.

The expected product of the reaction between a platinum metal fluoride and carbon monoxide is a lower fluoride, the carbon monoxide acting as a reducing agent. There have been observed, however, compounds with stoichiometries $Pt(CO)_2F_8$ and $Rh(CO)_2F_3$ (31). The compounds exhibit abnormally high valence states and coordination numbers.

Alkyl manganese pentacarbonyl will react with primary and secondary amines to produce the compound with formula $RCOMn(CO)_4NHR'R''$ (32). Since no carbon monoxide is given off, the mechanism is apparently not a simple displacement. Evidence has been advanced for an amine displacement of the alkyl group followed by attack of the alkyl group on one of the metal-carbonyl ligands. The mechanism is said to resemble a carbonylation-type reaction of alkyl manganese pentacarbonyl.

BIBLIOGRAPHY

1. J. D. Munro and P. L. Pausen, J. Chem. Soc., 1961, 3475.
2. Ibid, 3479.
3. Ibid, 3484.
4. R. P. M. Werner and S. A. Manastyrskyj, J. Am. Chem. Soc., 83, 2023 (1961).
5. W. D. Kaez, R. B. King, T. A. Manuel, L. D. Nichols and F. G. A. Stone, J. Am. Chem. Soc., 83, 4749 (1960).
6. J. Chatt, R. G. Guy and U. R. Watson, J. Chem. Soc., 1961, 2332.
7. J. E. Arnet and R. Pettit, J. Am. Chem. Soc., 83, 2954.
8. W. R. McClellan, H. H. Hoehn, H. N. Cripps, E. L. Muetterties and B. W. Hawk, J. Am. Chem. Soc., 83, 1601 (1961).
9. F. J. Impastato and K. G. Ihrman, J. Am. Chem. Soc., 83, 3726 (1961).
10. M. L. H. Green and P. Nagy, Proc. Chem. Soc., 1961, 378.
11. A. G. Massey and L. E. Orgel, Chem. and Ind., 1961, 436.
12. W. T. Dent, L. A. Duncanson, R. G. Guy, W. B. Reed and B. L. Shaw, Proc. Chem. Soc., 1961, 169.
13. F. Piacenti, C. Cioni and P. Pino, Chem. and Ind., 1960, 1240.
14. R. F. Heck and D. S. Breslow, J. Am. Chem. Soc. 83, 4203.
15. S. Nakamura and J. Halpern, J. Am. Chem. Soc., 83, 4102.
16. G. Booth and J. Chatt, Proc. Chem. Soc., 1961, 67.
17. R. B. King and F. G. A. Stone, J. Am. Chem. Soc., 82, 3833 (1960).
18. F. Basolo and A. Wojcicki, J. Am. Chem. Soc., 83, 520.
19. Ibid, 525.
20. A. Wojcicki and F. Basolo, J. Inorg. Nucl. Chem. 17, 77.
21. H. B. Fray and A. Wojcicki, Proc. Chem. Soc., 1960, 358.
22. T. A. Manuel, S. L. Stafford and F. G. A. Stone, J. Am. Chem. Soc., 83, 249 (1961).

23. T. D. Coyle, R. B. King, E. Pitcher, S. L. Stafford, P. Treichel and F. G. A. Stone, *J. Inorg. Nucl. Chem.*, 20, 172 (1961).
24. R. B. King, P. Treichel and F. G. A. Stone, *Proc. Chem. Soc.*, 1961, 69.
25. W. R. McClellan, *J. Am. Chem. Soc.*, 83, 1598.
26. C. C. Addison, M. Kilner and A. Wojcicki, *Proc. Chem. Soc.*, 1961, 202.
27. C. C. Addison, M. Kilner and A. Wojcicki, *J. Chem. Soc.*, 1961, 4839.
28. J. F. Tilney-Bassett, *Proc. Chem. Soc.*, 1960, 419.
29. R. B. King, P. Treichel and F. G. A. Stone, *Chem. and Ind.*, 1961, 747.
30. R. B. King and F. G. A. Stone, *J. Am. Chem. Soc.*, 82, 3833 (1960)
31. D. W. A. Sharp, *Proc. Chem. Soc.*, 1960, 317.
32. K. A. Keblys and A. H. Filbey, *J. Am. Chem. Soc.*, 82, 4204 (1960).
33. F. Calderazzo and F. A. Cotton, *Inorg. Chem.*, 1, 30 (1962).

ELECTRONIC CONFIGURATION OF TETRAHEDRAL
OXYANIONS OF TRANSITION METALS

P. Van Der Voorn

July 24, 1962

A number of models have been proposed for the electronic configurations of transition metal complexes in general, and of tetrahedral complexes in specific. It appears that ligand field theory can account for the experimental evidence in a more quantitative manner than any other model.

The first model which could, at least qualitatively, account for the observed spectra of tetrahedral complexes was the Bethe-Van Vleck ionic model. However when an attempt is made to make quantitative predictions, this model soon proves inadequate.

The intensity of a transition is commonly expressed in terms of the oscillator strength which is defined as

$$f = 4.32 \times 10^{-9} \int \epsilon_{\nu} d\nu, \text{ i.e. constant } \times \text{ area}$$

In the case of a Gaussian curve this can be approximated as

$$f = 4.60 \times 10^{-9} \epsilon_{\max} \nu_{1/2}$$

where $\nu_{1/2}$ is the width of the band where $\epsilon_{\nu} = 1/2 \epsilon_{\max}$
In theory

$$f = 1.085 \times 10^{-9} \left[\text{av}_I \sum_{II} \left| \int \psi_I^* \vec{r} \psi_{II} d\tau \right|^2 \right]$$

where ψ_I and ψ_{II} are the wavefunctions of the initial and final states, ν_{II} is the frequency of the band and

$$\vec{r} = \underline{i} x + \underline{j} y + \underline{k} z$$

$\text{av}_I \sum_{II}$ means summed over the excited states and this sum averaged over the initial states. e.g. if there is a three fold degenerate ground state and a two fold degenerate excited state, there are a total of 6 combinations which are averaged over three, because of the triply degenerate ground state (1).

If the e and t_2 wave functions in the Bethe-Van Vleck ionic model, are composed solely of $(3d)^n$ electronic configurations, no visible electronic dipole transitions are allowed. Vibronic perturbations lead to bands with intensities of only $f = 10^{-4}$ to 10^{-5} , compared to observed values of $f = 10^{-3}$.


Two alternate mechanisms have been suggested to account for the observed spectra:

- 1) mixing of $3d$ and $4p$ wave functions due to the tetrahedral crystal field, or
- 2) mixing of $3d$ and ligand wave functions due to the tetrahedral ligand field.

Assuming mechanism I above, Ballhausen and Liehr calculated

oscillator strengths for two tetrahedral complexes which have been well characterized i.e. CuCl_4^- and CoCl_4^- . There was very poor agreement between calculated and experimental values, hence mixing of 3d and ligand wavefunctions must occur and a molecular orbital model must be used (2).

One such model involves σ -bonding only. The linear combinations which, through group theory, can be formed from the σ bonds of the four ligands are found to be of a_1 and t_2 symmetry and are combined with the 3d (e and t_2) and 4s (a_1) metal atomic orbitals to form molecular orbitals




Oscillator strengths calculated for CoCl_4^- and CuCl_4^- , agree quite closely with experimental values but agreement for MnO_4^- and CrO_4^- is very poor (2).

The next step is to include π -bonding which must play an important role with oxygen as the ligand. The first such model for CrO_4^- and MnO_4^- was proposed by Wolfsberg and Helmholz(3). Linear combinations of the ligand orbitals lead to combinations of A_1 , E , T_2 and T_1 symmetry which are combined with the proper metal atomic orbitals of E , T_2 and A_1 symmetry; T_1 forms a non bonding orbital. Wolfsberg and Helmholz made extensive quantum mechanical calculations in order to find the relative energies of the molecular orbitals and were able to obtain values for the energies of the absorption bands of CrO_4^- and MnO_4^- , which agree fairly closely with observed values, as shown below:

	obs	MnO_4^-	calc	obs	CrO_4^-	calc
1st transition	2.29 ev		1.68ev	3.25ev		2.42ev
2nd transition	3.96ev		2.78ev	4.59ev		3.15ev

The (necessary) oversimplifications led to some surprising results as pointed out by Ballhausen and Liehr (2): Large stabilization of the metal non π -bonding orbital of e symmetry and inversion of bonding energy levels in going from MnO_4^- to CrO_4^- . They proposed an alternate energy level diagram but established the order of the molecular orbitals from chemical intuition and

they did not calculate the relative energies.



The diagram above is that for MnO_4^- or CrO_4^{2-} which have a "closed shell" structure. Oscillator strengths were also calculated for these two ions, with the results below

			f(calc)(2)	f(obs)	
CrO_4^{2-}	ν_1	27,000 cm^{-1}	6.7×10^{-2}	8.9×10^{-2}	(3)
	ν_2	37,000 cm^{-1}	0.092	0.11	(3)
MnO_4^-	ν_1	18,500 cm^{-1}	3.9×10^{-2}	3.2×10^{-2}	(4)
	ν_2	32,000 cm^{-1}	0.073	0.07	(4)

In order to arrive at these values, effective nuclear charges had to be postulated, but it was found that the results are rather insensitive to variations in the assumed charges.

The diagram above can be extended even further by including the metal 4p orbitals of t_2 symmetry to give an extra antibonding t_2^* molecular orbital, and the oxygen lone pairs of a_1 and t_2 symmetry to give two extra non-bonding molecular orbitals (5).

In 1939 Teltow(6) suggested that a correlation might exist between the long wavelength absorption edge and the ionic radii for ions with an electronic structure similar to CrO_4^{2-} and MnO_4^- . Carington and coworkers(7) plotted these two quantities for a number of ions and found a straight line relationship for the 3d and 4d transition metals; the 5d transition metals seemed to fall on a separate straight line, possibly because of increasing importance of 5f orbitals in bonding.

A considerable amount of electron spin resonance work has been done on these ions. The first work (8) reported has turned out to be partly in error because the spectra were obtained from samples dissolved in alkali glasses, rather than from single crystals. For the ions MnO_4^- , CrO_4^{2-} and ReO_4^{2-} , all containing one unpaired electron, one line was observed at 20°K. The

line width was highly dependent on temperature which is evidence for spin-orbit coupling. Spectra were also obtained for MnO_4^{3-} and FeO_4^{2-} as well as RuO_4^{2-} and ReO_4^{3-} , but later work on the first pair of ions has disproved the earlier results which appeared to indicate that the two unpaired electrons were present in a triply degenerate orbital, in agreement with the model proposed by Wolfsberg and Helmholz.

FeO_4^{2-} in a K_2CrO_4 crystal gave two lines with little temperature broadening and a g-value (2.000 ± 0.004) equal to the free spin value. This indicates that there are two unpaired electrons in a doubly degenerate molecular orbital. Two electrons in a triply degenerate orbital would give rise to spin-orbit coupling and a g-value considerably different from 2.000. Furthermore, the e^* molecular orbital must be well separated from the lower lying completely filled orbitals and the higher lying unoccupied levels (9). The two lines are a result of zero-field splitting(15).

Similar results were obtained for MnO_4^{3-} in Na_3VO_4 which gave six pairs of lines because the ^{55}Mn nucleus has a spin $I = 5/2$ (9).

In early work, two peaks were obtained also for RuO_4^{2-} and ReO_4^{3-} both with temperature broadening(8) but these results, similarly to those for FeO_4^{2-} and MnO_4^{3-} , may also turn out to be incorrect when single crystal studies are made.

In the study of MnO_4^{2-} in K_2CrO_4 crystal, six hyperfine lines and some quadrupole lines were observed (9). Quantum mechanical calculations by Schonland (10) show that the experimental results can be explained only on the assumption that the unpaired electron occupies a doubly degenerate orbital. The e^* orbital is largely concentrated in a 3d orbital centered on the Mn atom while the t_2^* orbital is spread out over the whole molecule. Attempts to fit the data to the diagram of Wolfsberg and Helmholz ended in failure.

The spectra of a number of anions had been interpreted by Carrington and coworkers (7) on the basis of the model proposed by Wolfsberg and Helmholz(3) but the above evidence (9,10) in favor of the model of Balhausen and Liehr (2) has prompted them to reconsider their assignments (11). The transitions $t_1 \rightarrow e^*$ and $t_1 \rightarrow t_2^*$ in MnO_4^{2-} are split because of Coulomb effects caused by the presence of one electron in the e^* level. Calculations indicate that the $e^* \rightarrow t_2^*$ transition should be comparable in intensity to the d-d transitions in transition metals, while the $t_1 \rightarrow e^*$ and $t_1 \rightarrow t_2^*$ should be very intense. They further came to the following conclusions:

1. There is strong bonding through the a_1 and t_2 levels which may be compared to the d^3s hybrid orbitals.
2. The unpaired electrons in a doubly degenerate level, are concentrated mainly on the metal ion and hence the electrons in the bonding e level are concentrated mainly in the oxygen atoms. Wolfsberg and Helmholz seem to have overestimated the importance of π -bonding.
3. The intense bands of the visible and UV regions are due to charge transfer transitions from oxygen to metal atoms.

All the evidence presented so far seems to favor the Ballhausen and Liehr model greatly over that of Wolfsberg and Helmholtz. However FeO_4^{3-} which has 3 electrons beyond the closed shell structure of MnO_4^- and should therefore have only one unpaired electron according to the Ballhausen and Liehr model, forms an exception. It has been found to have a magnetic moment of 3.6-3.7 B.M. compared to a theoretical value of 3.87 B.M. for three unpaired electrons (12). Lott and Symons (13) have attempted to prepare MnO_4^{4-} containing tervalent Mn which would also have three extra electrons. Although Mn(IV) may have been prepared there was no evidence for a tetrahedral species. For two other ions: FeO_4^{4-} and CoO_4^{4-} , which contain 4 and 5 electrons resp. beyond the "closed shell" structure of MnO_4^- , no magnetic moments have been reported (14).

References

1. C. J. Ballhausen, Progress In Inorganic Chemistry, Vol. II, Edited by F. A. Cotton, Interscience Publishers Inc., New York (1960), p. 251.
2. C. J. Ballhausen and A. D. Liehr, J. Mol. Spectroscopy, 2, 342 (1958).
3. M. Wolfsberg and L. Helmholtz, J. Chem. Phys. 20, 837, (1952).
4. Landolt-Börnstein, "Atom und Molekular Physik" Vol. I, Parts 3 and 4, Springer, Berlin (1951, 1955).
5. A. D. Liehr, J. Chem. Ed., 39, 135 (1962).
6. Y. Teltow, Z. phys. Chem., B43, 198 (1939).
7. A. Carrington, D. Schonland, M.C.R. Symons, J. Chem. Soc., 659 (1957).
8. A. Carrington, D. J. E. Ingram, D. Schonland, M.C.R. Symons, J. Chem. Soc., 4710 (1956).
9. A. Carrington, D. J. E. Ingram, K. A. K. Lott, D. S. Schonland, and M. C. R. Symons, Proc. Roy. Soc., A254, 101 (1960).
10. D. S. Schonland, Proc. Roy. Soc., A254, 101 (1960).
11. A. Carrington and M. C. R. Symons, J. Chem. Soc., 889 (1960).
12. W. Klemm, Angew. Chem., 66, 468 (1954).
13. K. A. K. Lott and M. C. R. Symons, J. Chem. Soc., 829 (1959).
14. R. Scholder, Angew. Chem., 66, 461 (1954).
15. A. Carrington and H. C. Longuet-Higgins Quart. Revs., 14, 427 (1960).

THE DEFECT SOLID STATE
AN INTRODUCTORY SURVEY

Violet I. Imhof

July 26, 1962

INTRODUCTION

Since the classic x-ray experiments by Laue in 1912 it has been realized that not only are there defects and disorder in the lattices of real crystals but also these imperfections are closely connected with the physical and chemical properties of solids. In this seminar we shall attempt to provide a general survey of this field with emphasis on an understanding of the nature of some of the fundamental defect types, so that the current literature may be read with comprehension. There have been several recent introductory reviews (1,2). More detailed material may be found in references 3-9 and in the series "Solid State Physics". (10)

PHONONS

The motion of atoms in a crystalline lattice is restricted to oscillation about an equilibrium point by the force field of neighboring atoms. The frequency of this mechanical vibration is designated ν_0 . For a one dimensional vibration where Hooke's Law holds, the allowed vibrational levels are given by $E_n = (n + 1/2) h \nu_0$, where n is a position integer or zero. Thus the energy difference between two states where $n = n'$ and $n = n''$ is $E = E_{n''} - E_{n'} = (n'' - n') h \nu_0$. The term phonon is given to the minimum ΔE , $h \nu_0$ (11).

No interaction between phonons in different vibrational planes would be observed if forces between atoms strictly followed Hooke's Law. The forces are not strictly Hooke's Law forces, however, and interaction does occur. Large amounts of phonon interaction can result in movement of atoms or ions into the lattice to produce vaporization. A smaller degree of interaction can cause the production of free electrons and positive holes or may elevate free electrons to higher energy states in the case of metals or alloys (9). The phonon concept has proven useful for example in treating thermal and electrical conduction (2) and semiconductivity (9).

ELECTRONS AND POSITIVE HOLES

We have just seen that phonon interaction in a crystal in thermal equilibrium can produce free electrons and electron absences (positive holes) in what would normally be an insulator.

These imperfections can also be generated by irradiation, high field conditions, or impurities, among other things. It is these electrons and holes that are the current carriers in semiconductors. They will be discussed more fully in connection with semiconductivity in a later section.

EXCITONS

It would be expected that an electron and positive hole would annihilate each other. It is possible however that prior to this, attraction would lead to the formation of a bound pair. The interaction should be much the same as in the hydrogen atom. The term exciton was first used for this pair by Frenkel (12). Wannier has identified excitons with wave functions and energy levels (13). Using these results Hiller and Marcus (14) showed that in general anisotropy exists in the motion of excitons and that the effective mass of an exciton corresponding to a dipole transition is of the order:

$$m_{\text{eff}} \approx (m_e / f_{n_0}) (R_0 / a)$$

where m_e = electronic mass, f_{n_0} = oscillator strength of the corresponding isolated atomic transition, a = equivalent Bohr radius of an internal exciton orbit, $R_0 = (3/4\pi n_0)^{1/3}$ and n_0 = atomic density. Thus if f is about one and a about 3\AA , the exciton mass is approximately the electronic mass. Thus an exciton may either be viewed as a neutral particle produced by partial recombination of an electron and a positive hole or as an excited state of an atom or ion capable of migration from one lattice cell to another.

Experimentally the exciton is inferred from peaks on the low energy side of the first absorption band of the crystal. Irradiation with light of the frequency of the first absorption band produces photoconductivity indicating ionization. Irradiation at the frequency of the subsidiary peaks however produces no photoconductivity indicating that the electrons are raised to an excited state below the ionization limit (15,16,17,18).

LATTICE DEFECTS

Of more chemical interest are those types of defects which concern the misplacement, displacement or nonplacement of atoms in the crystal. That is, there are three general classes of lattice defect. a) Atoms or ions may occupy metastable positions off the lattice sites, called interstitial positions. b) Lattice sites may be occupied by foreign atoms. If the impurity atoms are small enough they may also occupy interstitial positions. It is also possible that in a crystal containing A and B, B atoms might be found on A sites and vice versa. This does occur in alloys but is improbable in ionic lattices due to unfavorable electrostatic interactions. c) Lattice sites may be vacant. These positions are termed vacancies rather than holes to distinguish this ionic defect from the electronic defect, positive holes.

Interstitial atoms and vacancies may be discussed in terms of two possible kinds of disorder postulated by Frenkel(19) and Schottky(20,21). In the case of a Frenkel defect, ions of one kind may migrate to interstitial positions leaving behind an equal number of vacancies. In general it is the smaller cations which migrate but although it is much less likely anions may also take up interstitial positions. This is termed anti-Frenkel disorder. It is also conceivable that both ions may be found in these positions, this is unlikely and has never been observed. Schottky defects on the other hand do not necessitate interstitial ions. Here the ion migrates to the surface of a crystal, leaving a vacancy behind. To preserve electrical neutrality the number of cation and anion vacancies must be the same.

Although for a given set of conditions one type or another of disorder is apt to predominate, the various types are capable both of separate existence and coexistence. In fact for any real crystal at temperatures greater than 0°K some disorder must exist, for although energy is expended in defect formation, entropy is increased at the same time (22) Accurate knowledge of the energy necessary to form disorders would allow predictions of the type of disorder to be expected in a given case. These calculations have been attempted, but involve many assumptions, particularly about the effect of polarization occurring as a defect is produced (23,24,25,26) Stone gives a comprehensive review in the book by Garner(5). In general Schottky disorder is favored for crystals where ionic sizes are similar and neither the Van der Waals energy nor the dielectric constant are particularly high. The alkali metal halides meet these conditions fairly well, and in general show Schottky defects. Silver halides on the other hand better meet the conditions favoring Frenkel defects, i.e. a large difference in ionic sizes, high Van der Waals energy and dielectric constant.

There are various systems of notation in use. In that due to Rees (27) the nature of the atom occupying a lattice site and the nature of the site are given. Thus $A \mid \square_A$ means an atom of A on an A site. The symbol $A_{1-x} \mid \square_A$ means that the fraction 1-x of the A sites are occupied by A atoms. Similarly, an atom of A in an interstitial position is $A \mid \Delta$. In other notations (1) 0 represents an interstitial species, \square a vacancy and $A \cdot (B)$ the replacement of B by A. The subscript i may also be used to represent interstitial species.

IMPURITY ATOMS


The addition of foreign atoms to a metal lattice changes the number of free electrons. In nonmetals either new electronic levels are produced or existing levels are modified. Extensive reviews on the dependence of crystalline disorder on impurity content are available (6,1).

Here we will only discuss one aspect, semiconductivity.

The classic example is the case of silicon which has the tetrahedral diamond lattice. An arsenic atom added as an impurity can only fit itself into the tetrahedral arrangement by forming an As^+ ion and a free or nearly free electron. The bonding energy of this electron will be decreased by the square of the dielectric constant of the medium. For arsenic in silicon this is 0.09 eV. Hence at normal temperatures the electron is essentially ionized and the material is an N type semiconductor. If boron is used as the impurity atom it must accept an electron from the lattice in order to assume a tetrahedral configuration. This electron borrowing leaves a positive hole behind. The hole is not localized at any particular site and at real temperatures is ionized. Since conduction is by positive holes, this is a P type semiconductor.

The simplest model for the energy levels in insulating materials is shown diagrammatically below. (9) The filled band consists of permitted energy levels separated by a broad forbidden region from the conduction band where the electron is free to move about the crystal. In N type semiconductors there are donor levels a short distance below the conduction band. Electrons from these levels can easily be ionized to the conduction band. In P type semiconductors there are acceptor levels just above the highest filled band. Ionization of an electron from the filled band to an acceptor level leaves a conducting positive hole behind.

In more sophisticated model consideration is given to the existence in the forbidden zone of discrete levels of both donor and acceptor character (28,29). The model due to Morin also considers the effect of d orbitals (30).



COMPLEX DEFECTS - COLOR CENTERS

If a crystal of an alkali halide is heated with alkali metal vapor it undergoes discoloration due to a new absorption band in the visible region (31). This same band is also produced if the crystal is irradiated with x-rays. In this case another absorption also appears in the ultraviolet region (32). These are the F and V bands respectively.

The first model for the defect responsible for the F band, the F center, was proposed by de Boer(33). It pictures a defect consisting of an anion vacancy and a trapped electron. It bears a formal resemblance to the hydrogen atom. If modification is made for the effect of the centers crystalline surroundings and the electronic mass is replaced by an effective mass, calculations using Bohr energy levels give good order of magnitude results for the frequency of the F band. This model is somewhat oversimplified as shown by photoconductivity experiments (34). More detailed analyses have been made of the energetics involved (35, 3a) but the basic model holds. Another defect, the F' center, is believed to be an anion with two trapped electrons. It is produced by irradiation of discolored alkali metal halides with F center light (36).

The V band may also be produced by the alkali halide crystal in halogen vapor. The V centers responsible are formally the enantiomorphs of F centers. Seitz (3c) distinguishes four types of V centers on the basis of recent experimental evidence: V_1 , a hole bound to a single cation vacancy; V_3 , one hole bound to two cation vacancies; V_2 , two holes bound to two cation vacancies; V_4 , one hole bound to a unit of an anion vacancy and two cation vacancies.

CONCLUSION

An understanding of the nature of the defect solid state is basic to a grasp of such phenomena, as semiconductivity, luminescence, the magnetic properties of solids and the nature of solid reactions. While it has not been possible to discuss all possible crystal imperfections it is hoped that some insight into the nature of the problem has been provided.

BIBLIOGRAPHY

1. J. M. Honig, J. Chem. Educ., 34, 224 (1957).
2. Idem., Ibid., 34, 343 (1957).
3. F. Seitz, Revs. Mod. Phys., 26, 7(1954), a) pp 50-53, b) pp. 46-50, c) p. 57.
4. N. Mott, R. Gurney, "Electronic Processes in Ionic Crystals," 2nd ed., Clarendon, Oxford, 1950.
5. W. E. Garner, "Chemistry of the Solid State," Academic Press, New York, 1955.
6. K. Hauffe, "Reaktionen in und an festen Stoffen," Springer, Berlin, 1955.
7. F. Seitz, "Imperfections in Nearly Perfect Crystals", W. Shockley, ed., Wiley, New York, 1952.
8. W. Jost, "Diffusion in Solids, Liquids, Gases", Academic Press, New York, 1952.
9. T. J. Gray, "The Defect Solid State", Interscience, New York, 1957.
10. F. Seitz, D. Turnbull, "Solid State Physics", Academic Press, New York, 1954-1962.
11. I. Tamm, Z. Physik, 76, 849 (1932).
12. J. Frenkel, Phys. Rev., 37, 17, 1276 (1931).
13. G. Wannier, ibid., 52, 191 (1937).
14. W. Hiller, A. Marcus, Ibid., 84, 809 (1951).

15. L. Apker, E. Taft, Ibid., 82, 814 (1951).
16. Idem., Ibid., 81, 698 (1951).
17. Idem., Ibid., 79, 964 (1950).
18. M. Helb, Ibid., 81, 702 (1951).
19. J. Frenkel, Z. Phys., 35, 652 (1926).
20. W. Schottky, Z. phys. Chem., 29B, 335 (1935).
21. C. Wagner, W. Schottky, Ibid., 11B, 163 (1930).
22. Fowler, Guggenheim, "Statistical Thermodynamics", Cambridge, the University, 1939.
23. W. Jost, Z. physik Chem., 169A, 129 (1934).
24. Idem., J. Chem. Phys., 1, 466 (1933).
25. N. Mott, M. Littleton, Trans. Faraday Soc., 34, 485 (1938).
26. R. Hutner, E. Ritter, F. DuPre, J. Chem. Phys., 17, 198, 204 (1949), 18, 379 (1950).
27. A. Rees, "The Chemistry of the Defect Solid State", Methuen, London, 1955.
28. A. Rose, Phys. Rev., 97, 322 (1955).
29. W. Shockley, Bell System Tech. J., 28, 435 (1949).
30. F. Morin, Phys. Rev., 93, 1195, 1199 (1954).
31. F. Seitz, J. Chem. Phys., 6, 130 (1938).
32. R. Pohl, Phys. Z., 39, 36 (1938).
33. J. de Boer, Rec. trav. chim., Pays-Bas, 56, 301 (1937).
34. R. Pohl, G. Glaser, Ann. Phys., 29, 239 (1937).
35. J. Markham, Phys. Rev., 80, 500 (1952).
36. H. Pick, Ann. Phys., 31, 365 (1938).

For articles on current research see also the journal, "Physics and Chemistry of the Solid State".

COMPLEX CHEMISTRY OF PHOSPHINES AND PHOSPHINE OXIDES

William J. Randall

October 2, 1962


INTRODUCTION

The complex chemistry of ammonia and amines has been known for almost a century. Although the analogous phosphines were available, little characterization of metal complexes of these was done until relatively recently. In the ammonia series, stability decreases with an increase in the number of alkyl or aryl substituents on the nitrogen atom; however, the exact reverse is true in the phosphine series. Although PH_3 will coordinate, it is a strong acid under those conditions and will lose protons readily to form the metal phosphide. An explanation is that the phosphorus atom is large enough to accommodate organic groups and coordinate as well, and the organic groups in turn increase the basicity of the phosphorus by donation of electron density to the phosphorus. This is illustrated in the increasing dipole moment of the series PH_3 , PH_2R , PHR_2 , and PR_3 . The stability of these complexes is illustrated in the ability of the ligand to stabilize unusual oxidation states of certain metals.

PHOSPHINES


Most of the investigations have dealt with tertiary phosphine complexes. In general, these are made by allowing an anhydrous metal salt to react with the phosphine either in a solvent or in the fused state, followed by filtration of the insoluble complex or extraction and then precipitation, respectively. Such procedures were used in forming complexes of Ru(II) and Os(II) (1). These form dinuclear complexes of the formulae $[\text{Ru}_2(\text{PR}_3)_6\text{Cl}_3]\text{Cl}$ and $[\text{Os}_2(\text{PR}_3)_6\text{Cl}_3]\text{Cl}$. Chromium forms $[\text{Cr}(\text{PR}_3)_2\text{X}_3]_2$ ($\text{X}=\text{Cl}, \text{Br}$) from a benzene or toluene solution. (2,3) The halides CrX_2 ($\text{X}=\text{Cl}, \text{Br}$) give $[\text{Cr}(\text{PR}_3)_2\text{X}_2]$, which compounds are fairly unstable and perhaps polymeric. Tertiary phosphines also form compounds with silicon halides of the formula $[\text{Si}(\text{PR}_3)_2\text{X}_4]$ (4).

Nickel bromide and triethylphosphine form $[\text{Ni}(\text{PEt}_3)_2\text{Br}_2]_2$ which collects as dark-violet-red crystals. Treatment of this compound with bromine gives a compound which is violet-black in crystalline form and dark green in solution. Dipole moment studies indicate that it has a tetragonal pyramidal structure (in solution)



It is definitely an example of nickel (III) (5).

Using HReO_4 and excess PR_3 in ethanol in the presence of hydrohalic acids, Freni and Valenti obtained $\text{Re}(\text{PR}_3)_2\text{Cl}_3$, which upon treatment with hydrazine gives $[\text{Re}(\text{PR}_3)_2\text{X}_2](6)$. The $\text{Re}(\text{II})$ compound was thought to be polymeric:



The paramagnetism of the compound corresponds to one unpaired electron, which indicates that the species in solution is square planar with cis chlorides. The $[\text{Re}(\text{PR}_3)_2\text{X}_3]$ compounds are non-electrolytes, very poorly soluble, diamagnetic, and non-reactive toward elemental halogens. Hence the structure is thought to be dimeric with the rhenium atom coordinately saturated. Chatt and Rowe (7) found that the compounds reported to be $\text{Re}(\text{PPh}_3)_2\text{Cl}_3$, $\text{Re}(\text{PPh}_3)_2\text{Br}_2$, and $\text{Re}(\text{PPh}_3)_2\text{I}_2$ were really $\text{ReOCl}_3 \cdot (\text{PPh}_3)_2$, $\text{ReO}(\text{OEt})(\text{PPh}_3)_2\text{Br}_2$, and $\text{ReO}(\text{OEt})\text{I}_2(\text{PPh}_3)_2\text{I}_2$, respectively. These workers concluded from their investigations that the formation of "oxo" and "alkoxy" complexes of rhenium is a general phenomenon when alcohol is used as the preparatory medium. Colton and Wilkinson also report a $\text{Re}(\text{PPh}_3)_2\text{I}_4$ (8). Other complexes of tertiary phosphines are reviewed in references 9 and 10.

Thermodynamic studies of the adducts of trialkyl phosphines with diacetyl-bisbenzoylhydrazino-nickel(II), showed them to be 4-11 times as stable as those with trialkyl amines. Triphenyl phosphine would not form an adduct with this metal species owing to the steric interaction of the three phenyl groups with the metal complex. However, the ligands PEt_3 , PPr_3 , and PBu_3 give ΔF values of 8.43, 7.85, and 7.85 (kcal/mole), respectively (11).

Triphenyl phosphine with -onium iodomercurate salts-- M_2HgI_4 , MHgI_3 , MHg_2I_5 , MHg_3I_8 (M = quaternary cation)-- in acetone, or ethanol, forms iodotriphenylphosphine mercury(II) complexes and either an -onium iodomercurate with a higher (iodine/mercury) ratio than the reactant or a quaternary iodide (12). Further reaction of these complexes with alkyl halides gives the corresponding phosphonium halomercurates (13).


Compounds of the form $[\text{Ag}(\text{P}(\text{C}_6\text{H}_{11})_3)_2\text{I}]$, $[\text{Ag}(\text{P}(\text{C}_6\text{H}_{11})_3)\text{I}]$, and $[\text{Ag}(\text{P}(\text{C}_6\text{H}_{11})_3)_2\text{SCN}]$ have been reported. The latter forms monoclinic crystals of the space group C or C_2/c with cell dimensions $a = 16.80$, $b = 9.30$, and $c = 24.64$ (Å), $\beta = 108^\circ 40'$, and density = 1.33g./cm^3 . Cryoscopic measurements in benzene indicate that the compound is a monomer in solution (14).

The compound $\text{P}(\text{PhCH}_2)\text{Et}_2$ and excess NiBr_2 in toluene give $[\text{NiBr}_2 \cdot \text{P}(\text{PhCH}_2)\text{Et}_2]$ (15). The structure of some first-row transition metal phosphine complexes is discussed in reference 16.

The ditertiary phosphines, $R_2P(CH_2)_nPR_2$, are analogous to alkylene diamines and form similar complexes (10,17,18,19,20). Also reported are complexes of the general composition $[M(Ph_2P-PPh_2)_2-X_2]$, which are made from $NiBr_2$, $FdCl_2$, $CoBr_2$, CoI_2 , (21,22) and Cu_2Br_2 (21).

A departure from the ordinary synthetic route is seen in the mixing of $P(PhCH_2)_2H_2^+Br^-$ and K_2PdBr_4 in aqueous-alcoholic solution, which gives the product $[Pd(P(PhCH_2)_2H)_2Br_2]$ (pale yellow crystals) (23). The preparation of other secondary phosphine complexes and primary phosphine complexes is analogous to that described above.

The compounds $PHPh_2$ and $CoBr_2$ give three products which are thought to be $[Co(PHPh_2)_3Br_3]$, $[Co(PHPh_2)_3Br]Br$, and $[(PHPh_2)_4-Co(Br)_2Co(PHPh_2)_4]Br_2$. The corresponding nickel bromide gives $[Ni(PHPh_2)_2(PPh_2)_2]$, and with $PHEt_2$ gives $[Ni(PHEt_2)_4Br_2]$ (24). The reaction of $PHPh_2$ on palladium halides or halopalladates produces $[Pd(PHPh_2)(PPh_2)X]_2$, (a), and some $[Pd(PHPh_2)_2X]$, (b), ($X=Cl, Br, I$) and $[Pd(PHPh_2)_2(PPh_2)_2]$, (c). The bromide also forms $[Pd(PHPh_2)_3Br]Br$, (d). The tendency to form complexes containing the anionic ligand (PPh_2^-) decreases in the order $Cl > Br > I$. The presence of hydrohalic acids opposes the formation of these species and favors the dihalide complex--excess $PHPh_2$ favors the formation of (a) and (c) in alcohol and (d) in benzene. The use of alcohols as solvents favors the formation of (a) and (b). Compound (a) is a dimer which does not undergo ligand exchange with PPh_3 , but does with orthophenanthroline to form (26)



Primary phosphine complexes with CoX_2 ($X=Cl, Br, NO_2$) are thought to be tetrahedral, and other complexes have the following geometries: $[Co(PH_2Ph)_4I_2]$, octahedral; $[Co(PH_2Ph)_3I]$, trigonal bipyramidal; $[(PH_2Ph)_2ClPd-PPh-Pd(PH_2Ph)_2Cl]$, binuclear (square planar Pd). Those of chromium are like those containing the $PHPh_2$ ligand. Copper(I) chloride forms a tetramer, $[Cu(PH_2Ph)Cl]_4$ (27).

PHOSPHINE OXIDES

Pickard and Kenyon reported a phosphine oxide compound of iron in 1906 (28). However, insufficient understanding of complex chemistry was available for a reasonable explanation of the structure. Divalent and trivalent chromium form $[Cr(OPEt_3)_2Cl_2]$ and $[Cr(OPR_3)_3(SCN)_3]$, respectively. These are non-electrolytes, as proved by conductivity and molecular-weight measurements (29). Complexes of trivalent vanadium (30) and nickel(II) bromide (31) have also been characterized. Tri-n-octylphosphine oxide forms a complex with thorium, which allows thorium(IV) to be separated from sulfates (32). Some first-row transition metal complexes with phosphine oxides are discussed in reference 16.

UNUSUAL PRODUCTS

Phosphines have been used to stabilize some hydrido (33), organometallic (34-38) and carbonyl (39-54) compounds. The compounds $(Et_2P-AlCl_2)_3$ and $(Et_2P-AlCl_3:OPCl_3)$ (55) have also been described.

SUMMARY

Phosphines and phosphine oxides as ligands provide an area of complex chemistry that is characterized by highly colored compounds, some of which have quite unusual structures and oxidation numbers.

SELECTED BIBLIOGRAPHY

1. J. Chatt and R. G. Hayter, J. Chem. Soc., 896(1961).
2. K. Issleib and H. O. Fröhlich, Z. anorg. u. allgem. Chem., 298, 84 (1959).
3. K. Issleib and A. Tzschach, Z. anorg. u. allgem. Chem., 297, 121 (1958).
4. K. Issleib and H. Reinhold, Z. anorg. u. allgem. Chem., 314, 113 (1962).
5. K. A. Jensen and B. Nygaard, Acta. Chem. Scand., 3, 474 (1949).
6. M. Freni and V. Valenti, J. Inorg. Nucl. Chem., 16, 240 (1961).
7. J. Chatt and G. A. Rowe, Chem. and Ind., 92 (1962).
8. R. Colton and G. Wilkinson, Chem. and Ind., 1314 (1959).
9. J. C. Bailar, Jr., and D. H. Busch, The Chemistry of Coordination Compounds, J. C. Bailar, Jr., Ed., Reinhold Publishing Corp. New York, 1956, p. 83.
10. C. E. Wymore, Doctoral Dissertation, University of Illinois, 1956, pp. 1-29.
11. L. Sacconi, G. Lombardo, and P. Paolette, J. Am. Chem. Soc., 82, 4186 (1960).
12. G. B. Deacon and B. O. West, J. Chem. Soc., 5127 (1961).
13. G. B. Deacon and B. O. West, J. Inorg. Nucl. Chem., 24, 169 (1962).
14. A. Turco, C. Panattoni, and E. Frasson, Ricerca sci., 29, 544(1959).
15. K. Issleib and E. Pelebe, Chem. Ber., 92, 3152 (1959).
16. J. T. Donoghue, "Some Recent Developments in Tetrahedral Complexes," Inorganic Seminars, University of Illinois, May 15, 1962, pp. 200-212.
17. C. E. Wymore and J. C. Bailar, Jr., J. Inorg. Nucl. Chem., 14, 42 (1960).
18. K. Issleib and G. Hohlfield, Z. anorg. u. allgem. Chem., 312, 169 (1961).
19. J. Chatt and R. G. Hayter, J. Chem. Soc., 5507 (1961).
20. J. Chatt, F. A. Hart, and D. T. Rosevear, J. Chem. Soc., 5504 (1961).
21. K. Issleib and G. Schwager, Z. anorg. u. allgem. Chem., 310, 43 (1961).
22. K. Issleib and G. Schwager, Z. anorg. u. allgem. Chem., 311, 82 (1961).
23. R. G. Hinton and F. G. Mann, J. Chem. Soc., 2835 (1959).

24. K. Issleib and E. Wenschuh, Z. anorg. u. allgem. Chem. 305, 15 (1960).
25. K. Issleib and G. Düll, Z. anorg. u. allgem. Chem., 305, 1 (1960).
26. R. G. Hayter, J. Am. Chem. Soc., 84, 3046 (1962).
27. K. Issleib and G. Walde, Z. anorg. u. allgem. Chem., 312, 287 (1961).
28. R. H. Pickard and J. Kenyon, J. Chem. Soc., (Transactions), 89, 262 (1906).
29. K. Issleib, A. Tzschach, and H. O. Fröhlich, Z. anorg. u. allgem. Chem., 298, 164 (1959).
30. K. Issleib and G. Bohn, Z. anorg. u. allgem. Chem., 301, 188 (1959).
31. K. Issleib and W. Seidel, Chem. Ber., 92, 2681 (1959).
32. W. J. Ross and J. C. White, Anal. Chem., 31, 1847 (1959).
33. J. Chatt, T. A. Duncanson, and B. L. Shaw, Chem. and Ind., 859 (1958).
34. G. Booth and J. Chatt, Proc. Chem. Soc., 67 (1961).
35. J. Chatt and B. L. Shaw, J. Chem. Soc., 285 (1961).
36. J. Chatt and B. L. Shaw, Chem. and Ind., 675 (1959).
37. L. S. Meriwether, C. E. Colthup, M. L. Fiene, and F. A. Cotton, J. Inorg. Nucl. Chem., 11, 181 (1959).
38. R. A. Plowman and F.G.A. Stone, Inorg. Chem., 1, 518 (1962).
39. D. W. McBride, S. L. Stafford, and F.G.A. Stone, Inorg. Chem., 1, 386 (1962).
40. R. F. Heck and D. S. Breslow, J. Am. Chem. Soc., 82, 4438 (1960).
41. J. Chatt, P. L. Fauson, and L. M. Venanzi, Organometallic Chemistry, H. Zeiss, Ed., Reinhold Publishing Corp., New York, 1960, pp. 468, 477 and 478.
42. R. D. Gorsick, J. Am. Chem. Soc., 84, 2487 (1962).
43. W. Hieber, G. Faulhaber, and T. Theübert, Z. anorg. u. allgem. Chem., 314, 125 (1962).
44. A. Sacco and M. Freni, J. Inorg. Nucl. Chem., 8, 566 (1958).
45. G. Booth and J. Chatt, J. Chem. Soc., 2099 (1962).
46. W. Hieber and E. Lindner, Chem. Ber., 95, 273 (1962).
47. G. N. Schräuzer, J. Am. Chem. Soc., 81, 5310 (1959).
48. W. Hieber and J. Peterhaus, Z. Naturforsch., 14b, 462 (1959).
49. W. Hieber and W. Freyer, Chem. Ber., 92, 1765 (1959).
50. W. Hieber and A. Hipp, Chem. Ber., 92, 2085 (1959).
51. T. S. Meriwether and M. L. Fiene, J. Am. Chem. Soc., 81, 4200 (1959).
52. W. Hieber and W. Schropp, Z. Naturforsch., 14b, 460 (1959).
53. W. Hieber, K. Englert, and K. Rieger, Z. anorg. u. allgem. Chem., 300, 1295 (1959).
54. E. W. Abel, M. A. Bennett, and G. Wilkinson, J. Chem. Soc., 2323 (1959).
55. G. Fritz and G. Trenczek, Z. anorg. u. allgem. Chem., 313, 236 (1961).

SOME SELECTED USES OF OPTICAL ROTATORY
DISPERSION IN COORDINATION CHEMISTRY

L. J. Boucher

October 9, 1962

INTRODUCTION

The octahedral configuration of certain hexacoordinate complexes implies the existence of mirror-image isomers, and hence optical activity (the ability of single molecules to rotate plane polarized light). This possibility was experimentally realized in 1911 by Werner when he resolved the $[\text{Co}(\text{en})_2\text{NH}_3\text{Cl}]^{2+}$ complex. Thus at the very heart of Werner's theory of coordination compounds lies the phenomenon of optical activity and with it the experimental measurement of the angle of rotation, α . It has since become customary to measure the change of rotation with wavelength of incident light. This is known as Optical Rotatory Dispersion (O.R.D.).

The measurement of O.R.D. has steadily grown in importance and today its importance as a research tool is evidenced by the keen interest in it shown by coordination chemists both here and abroad. Although O.R.D. does not have as wide an application to inorganic chemistry as other physical methods such as N.M.R., I.R., and electronic absorption spectroscopy, it does have enough important applications to warrant at least a cursory knowledge of its fundamentals by inorganic chemists and in particular by those concerned with coordination.

The theoretical aspects of the optical rotatory power of transition-metal complexes will not be dealt with here other than to say that such theories do exist (quantum mechanical as well as classical) which purport to relate the optical rotatory power to fundamental mechanical properties of the molecule under consideration. The present state of these theories is somewhat unfinished, but for those who are interested in further pursuit, the references noted provide a starting point: Moffit(2), Sugano(3), Liehr(4), and Piper(5).

A molecule may possess optical rotatory power if and only if it is non-superimposable upon its mirror image. This is equivalent to saying that an active molecule can neither have a center nor a plane of symmetry. Lowery(6) has used the term "dissymmetric" for figures possessing axes of symmetry but no planes center of symmetry and no alternating axis of symmetry. Dissymmetric figures are not necessarily assymmetric. The simple Drude equation(7) expresses the variation of rotatory power with the wavelength of plane polarized light:

$$[\alpha]_D = \sum_i \frac{k_i}{\lambda^2 - \lambda_i^2} \quad (1)$$

where k_i is a constant and λ_i is the wavelength of the maximum absorption of the optically active absorption band. The observed

rotation α is experimentally determined. It is related to the specific rotation $[\alpha]$ by the following expression

$$[\alpha] = \frac{100}{dc} \cdot \alpha \quad (2)$$

where d is the cell length in decimeters and c is the concentration of optically active substances in gr./100 ml. of solution. In the literature, O.R.D. curves are usually plotted in terms of the molecular rotation $[M]$ which is related to $[\alpha]$ by the following expression:

$$[M] = \frac{[\alpha] (\text{Mol. wt.})}{100} \quad (3)$$

Equation (1) is satisfactory when λ is far removed from the wavelength of the optically active absorption band. If it is within the absorption band, a more complicated expression must be used.

According to Djerassi(8), there are three types of O.R.D. curves: (1) plain curves with no maximum or minimum corresponding to the Drude equation; (2) single Cotton-effect curves with only one maximum and one minimum; and (3) multiple Cotton effects with two or more peaks and troughs. The rapid increase and reversal of sign of optical rotatory power with changing wavelength in an absorption band is known as an anomalous dispersion or as the Cotton effect when associated with a particular electronic transition.




Figure 1 - O.R.D. curve showing Cotton effect.

Figure 1 represents an ideal case in which the observed rotation is due to a single electronic transition and there are no contributions to the rotatory dispersion from other bands. The center of inversion, λ_0 , usually coincides with the wavelength of an absorption maximum. The absorption band is commonly referred to as an optically active absorption band. Both a chromophore and an optically active center must be present within a molecule to produce the Cotton effect. The chromophore may be at the center of activity, or the chromophore and active center may exist in different parts of the molecule, provided there is an interaction between them. The latter condition is called the vicinal effect and will be discussed below.

ASSIGNMENT OF CONFIGURATION

A knowledge of the relative configurations of optical antipodes is of great importance in determining stereochange in some substitution reactions at octahedral metal sites. An example of this is the reaction of l-cis[Co(en)₂Cl₂]⁺ with ethanol and hydrogen fluoride in a 1:1 ratio in the presence of silver carbonate to form d-cis[Co(en)₂F₂]⁺. It should be emphasized that the sign of rotation cannot be used to determine relative configuration. The d or l that is usually quoted in the literature tells only whether an optical antipode gives a positive or negative rotation of plane polarized light of a certain wavelength (usually the sodium D line). In the above example the O.R.D. curves of the reactant and product are very similar, leading to a postulated retention of configuration instead of an opponent inversion. Although O.R.D. measurements alone cannot determine relative configuration, it is possible to assign relative configurations, and even absolute configurations when they are used in conjunction with special x-ray crystallographic techniques.

Pioneer work by Mathieu on O.R.D. measurements of a number of cobalt complexes of the general formula d-cis[Co(en)₂AA'ⁿ⁺] with A and A' = NH₃, H₂O, Cl⁻, NCS⁻, Br⁻, NO₂⁻ or AA' = CO₃²⁻, OX²⁻; the complexes d-cis[Cr(en)₂Cl₂]Cl, d-[Co(en)₂OX]Cl and d-cis[Ir(OX)₂Cl₂](10); and 9Me(en)₃³⁺, [Me(OX)₃]³⁻, with Me = Cr³⁺, Rh³⁺ AND Ir³⁺ (11). He showed that complexes of similar composition which had O.R.D. curves with the same sign and some general shape of the Cotton effect for similar electronic transitions possess the same configuration, whereas if the slopes of the dispersion curves are of opposite sign, the configurations are opposite. Mathieu's postulate was made in conjunction with Werner's solubility rule, i.e., for analogous complex ions, the enantiomorphic forms which produce the less soluble salts with a common resolving agent are considered to have the same generic configuration.

The absolute configuration of an antipode can be determined by special x-ray crystallographic techniques alone. The experiments of Saito and coworkers (12) indicate the absolute configuration of 2-D[Co(en)₃]Cl₃.NaCl.6H₂O to be D(+).





Figure 2. - Absolute configuration of D(+)[Co(en)₃]³⁺.

This figure represents D(+) complex cation, which possesses D₃ symmetry (three fold axes out of plane of paper). If the enclosed areas in Figure 2 are considered as blades of propeller and the


darkened lines as the leading edges of the three propellers. Clockwise rotation of the figure about its three-fold axis shows that the blades will be driven away from the viewer. If the dashed lines and darkened lines are interchanged with those opposite them and the same procedure is carried out the blades now move toward the viewer. The former is called a left-handed system and the latter a right-handed system. Piper(13) has proposed that the left-handed system be designated Λ and the right handed system be designated Δ . The absolute configuration of other $[\text{Me}(\text{en})_3]^{3+}$ complexes may now be determined by simply recording O.R.D. curves and comparing the sign of the Cotton effect for similar electronic transitions. With more and more x-ray determinations being carried out and with the help of O.R.D. measurements, the assignment of absolute configurations to a host of complexes will be possible.

COMPLEXES WITH OPTICALLY ACTIVE LIGANDS

Pfeiffer (14) prepared square -planar complexes of Cu(II) and Ni(II) of the following formulations:



I



II

In these, the ligand was optically active. In each case the O.R.D. curves showed strong Cotton effects. Neither of the ligands itself showed any Cotton effect in the wavelength range investigated. Mellor (15) proposed that the electronic transition giving the Cotton effect was due to a metal-ligand bond chromophore (Me-N). The absorption band is optically active because of the so-called vicinal effect (16). Since the metal itself is not the site of the optical activity, the asymmetric carbon in the ligand must interact with the chromophore, making the absorption band optically active.

The essential nature of this interaction or coupling is still unknown. The effect is noted even when the asymmetric carbon is removed beyond the position adjacent to the chromophore. Pfeiffer and Christeliet (17) applied this effect to measure the O.R.D. of some Cu(II) complexes of naturally-occurring optically active Δ -amino acids. The O.R.D. curves for all the complexes (similar to Structure II) and the observed Cotton effects were similar in shape and were in the same region of the spectrum. It was thus shown that the configuration of the asymmetric centers in the different acids is the same and hence that all natural amino acids have the same configuration. Recently the same technique was applied in an attempt to determine the relative configurations of Δ -amino acids with more than one center of assymetry (18). The

data did not allow an unambiguous conclusion as to the configuration of the amino acids, but it was possible to correlate the relative configurations of the Δ asymmetry with the shape and the position of the Cotton effects.

In those cases where an optically active ligand (A-A) is in a complex of the type $[\text{Co}(\text{A-A})_2\text{B}_2]^{n+}$, both the cis and trans forms are optically active and show Cotton effects near the first absorption band. The optical activity in the trans isomer is due to the vicinal effect of the asymmetric center on the ligand. Inasmuch as the optical activity is due mainly to the arrangement of the groups about the central atom, O.R.D. curves of complexes of similar (cis) configurations should have the same characteristics whether the coordinating base is optically active or not, (e.g., l-propylenediamine versus ethylenediamine). The trans configuration, on the other hand, will show a greatly different O.R.D. curve, since the activity arises from the presence of the optically active base.

Bailar(19) and McReynolds(20) used O.R.D. measurements to assign a cis configuration to $[\text{Co}(\underline{l}\text{-pn})_2\text{Cl}_2]\text{Cl}$ and $[\text{Co}(\underline{l}\text{-pn})_2(\text{NO}_2)_2]\text{NO}_2$ and a trans configuration to $[\text{Co}(\underline{l}\text{-pn})_2(\text{NH}_3)_2]\text{Cl}_3$ on the basis of the striking similarity of the curves of the first two complexes to those of d-cis $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$ and d-cis $[\text{Co}(\text{en})_2(\text{NO}_2)_2]\text{NO}_2$. The latter complex was assigned a trans configuration because of the dissimilarity of O.R.D. curves to that of the corresponding ethylenediamine complex d-cis $[\text{Co}(\text{en})_2(\text{NH}_3)_2]\text{Cl}_3$. This sort of analysis requires a knowledge of the O.R.D. curves of analogous compounds which can exist in a active form only in the cis configuration. The same procedure was used to assign the trans configuration to $[\text{Co}(\underline{l}\text{-pn})_2\text{NH}_3\text{Cl}]\text{Cl}_2$ and $[\text{Co}(\underline{l}\text{-pn})_2(\text{NCS})_2]\text{Cl}$ (21). It was also found that the O.R.D. curve was inverted by replacing l-pn by d-pn in trans- $[\text{Co}(\underline{l}\text{-pn})_2(\text{NH}_3)_2]\text{Cl}_3$, therefore showing that the optical activity of the complex is intimately connected with the optical activity of the base.

In order to investigate the nature of the induced asymmetry that arises when propylenediamine (and other optically active amines) coordinates with a metal that in itself is not dissymmetric, O'Brien and Tool (22) measured O.R.D. curves of Zn^{2+} , Cd^{2+} , Ag^+ , and Pt^{2+} -l-propylenediamine complexes in the visible region of the spectrum. The curves showed no Cotton effects and could be described by a single one-term Drude equation (one optically active electronic transition in the far ultraviolet). The dispersive power was found to be independent of the metal and the nature of the optically active amine, but dependent on the formation function \bar{n} . It was concluded therefore that the activity of the complex arises only from the ligands and not from any induced asymmetry around the central atom.

ABSORPTION SPECTRA

Hidaka and coworkers claim that in order to understand fully the nature of absorption bands and especially their splitting and overlapping, O.R.D. curves must be carefully analyzed. While studying the complexes D-cis $[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$ and D-cis $[\text{Co}(\text{en})_2\text{ClNO}_2]\text{Cl}$, they found that the Cotton effect corresponding to the first absorption

band is actually a multiple Cotton effect made up of the superposition of two Cotton effects of the same or opposite sign and the tail of a Cotton effect in the far ultraviolet. The multiple Cotton effects are of equal magnitude and arise because of the splitting of the first absorption band into components \underline{I}_a and \underline{I}_b . The second absorption band gives rise to only one Cotton effect (of lesser magnitude than the first). Further studies (24) of the complexes \underline{D} -Na₂[Co(EDTA)NO₂], \underline{D} -K[Co(EDTA)], \underline{D} -K₂[Co(EDTA)Cl], and \underline{D} -cis[Co(en)₂(NO₂)₂]I lead to the same results, i.e., the first absorption-band splitting gives rise to a multiple Cotton effect. These workers also reported that \underline{D} -cis[Co(en)₂(NO₂)₂]I shows only one Cotton effect at the first absorption band, which may in fact be a superposition of two Cotton effects of the same sign that are too close together for resolution. The observed absorption spectrum of \underline{D} -K[Co(EDTA)] does not show any splitting in the first absorption band due to the diffuseness of the bands.

In a later paper, Hidaka and coworkers (25) studied the complexes \underline{D} -Na[Co(en)(OX)₂] and \underline{D} -K[Co(en)(Malo)₂]. The multiple Cotton effect was also observed for the split first absorption band and a single Cotton effect for the second absorption band. Shimura(26) observed the same result with an optically active ligand where the metal atom is not in the site of activity in the complex. The O.R.D. curve for [Co(NH₃)₄(l-leuc)](ClO₄)₂ exhibited multiple Cotton effects of opposite sign near the first absorption band. These can be attributed to the vicinal effect between the cobalt atom and the asymmetric carbon of the amino acid. It was also found that \underline{D} -[Co(en)₂l-leuc]²⁺ has a rotational power greater than \underline{D} -[Co(NH₃)₄l-leuc]²⁺ and shows only one Cotton effect at the first absorption band. It seems probable that the dissymmetry around the central cobalt atom contributes more profoundly to the rotatory power in the region of the first absorption band than the asymmetric carbon atom in the coordinated ligand. Neither complex shows any Cotton effect in the vicinity of its second absorption band. It also was noted that the center of inversion (wavelength where the Cotton effect crosses the zero rotation axis) was almost always in the longer wavelength region in comparison with the position of the absorption maximum. Some representative data for this section are listed in Table I.

MISCELLANEOUS

There are many other less broad applications of O.R.D. to coordination chemistry cited in the literature, some of which deserve mention. Albinas and coworkers (27) have found that the change in O.R.D. of a dissymmetric complex ion when the optically inactive counter ion associated with it is changed can be related to the degree of ion pairing. Larson (28) maintains that by studying the variation of O.R.D. with concentration of counter ion, the nature of the forces responsible for the association effects or upon the coordination sites preferred by the different ligands can be decided. Kling (29) and Woldbye (30), in a study of the effect of differences between similar chelates on O.R.D. of complexes, report that when ethylenediamine is replaced by trimethylenediamine in

Table I. Comparison of Center of Inversion and Absorption Maximum for Some Cobalt Complexes. (First Absorption Band)

<u>Compound</u>	<u>$\nu_{\text{abs.}}^*$</u>	<u>$\nu_{\text{rot.}}^*$</u>	<u>Cotton Effect</u>	<u>Ref.</u>
<u>D-cis</u> [Co(en) ₂ Cl ₂]Cl Ia	49.5	48	-	23
Ib	57.5	57	+	
<u>D-cis</u> [Co(en) ₂ ClNO ₂]Cl	60 (60)	55 (55)	- -	23
<u>D-Na</u> [Co(EDTA)NO ₂]	51.4 60.4	49 59	- +	24
<u>D-K</u> [Co(EDTA)]	55.9	51.5 61.0	- +	24
<u>D-CisK</u> ₂ [Co(EDTA)Cl]	52.0	51.8 57-63	-	24
<u>D-Cis</u> [Co(en) ₂ (NO ₂) ₂]I	65.5 (65.5)	69 (69)	+ +	24
<u>D-Na</u> [Co(en)(OX) ₂]	55.6	52 60	- -	25
<u>D-Na</u> [Co(en)(Malo) ₂]	55.6	50 56.0	- +	25
[Co(NH ₃) ₄ (<u>l-leuc</u>)](ClO ₄) ₂	60.8	54.6 64.6	+ -	26
<u>D</u> -[Co(en) ₂ (<u>l-leuc</u>)](ClO ₄) ₂	61.5	59.5	+	26

*
 ν in units of 10^{13} sec^{-1}

$[\text{Co}(\text{en})_3]^{3+}$ and $[\text{Cr}(\text{en})_3]^{3+}$, the molecular rotation contributed to the first absorption band is reduced by a factor of between 5 and 10 even though the absorption spectra are almost identical. In work being done in the new field of bio-inorganic chemistry, O.R.D. has been used to study the interaction of metal ions with sites in hemoproteins(31) and metalloproteins (32).

CONCLUSIONS

This seminar has attempted to show the importance of O.R.D. in solving structural problems in coordination chemistry. The theory and measurement of circular dichroism, a closely related topic has not been mentioned since this would require another seminar of equal length (for an introduction to this topic see Jaegen (33)).

References

1. Werner, A., Ber., 44, 1887 (1911).
2. Moffitt, J. W., J. Chem. Phys., 25, 1189 (1956).
3. Sugano, S., J. Chem. Phys., 33, 1883 (1960).
4. Liehr, A. D., "Advances in the Chemistry of the Coordination Compounds", S. Kirschner, Ed., The Macmillan Company, New York, 1961, pp. 422-429.
5. Piper, T. S., J. Chem. Phys., 36, 2224 (1962).
6. Lowry, T.M., "Optical Rotatory Power", Langmans Green and Company, New York, 1935, pp. 26-27.
7. Drude, P., "Lehrbuch der Optik", 2nd Ed., S. Hirzel Verlag, Leipzig, 1906.
8. Djerassi, C., "Optical Rotatory Dispersion", McGraw-Hill Book Company, Inc., New York, 1960, pp. 11-17.
9. Matoush, W. R., and Basolo, F., J. Am. Chem. Soc., 78, 3972(1956).
10. Mathieu, J. P., Bull. soc. chim. [5], 3, 497 (1936).
11. Mathieu, J. P., J. chim. phys., 33, 78 (1936).
12. Saito, Y., Nakatsu, K., Shiro, M., and Kuroya, H., Acta Cryst., 8, 729 (1955).
13. Piper, T. S., J. Am. Chem. Soc., 83, 3908 (1961).
14. Pfeiffer, P., Christeleit, W., Hesse, Th., Pjitzner, H., and Thielert, H. F. Prakt. Chem., 150, 261 (1938).
15. Mellor, D. P., J. Proc. Roy. Soc. N. S. Wales, 75, 157 (1942).
16. Kuhn, W., "Freundenbergs Stereochemie Lfg 2", F. Deuticke, Leipzig, 1933, p. 317.
17. Pfeiffer, P., and Christelert, W., Z. physiol. Chem., 245, 197 (1937).
18. Izumiya, J., Winitz, M., Bianbaum, S. M., and Greenstein, J.P., J. Am. Chem. Soc., 78, 1602 (1956).
19. Bailar, J. C., Jr., and McReynolds, J. P., J. Am. Chem. Soc., 61, 3199 (1939).
20. McReynolds, J. P., University of Illinois thesis, 1938.
21. O'Brien, T.D., McReynolds, J. P., and Bailar, J.C., Jr., J. Am. Chem. Soc., 70, 749 (1948).
22. O'Brien, T. D., and Tool, R., J. Am. Chem. Soc., 77, 1368(1955).
23. Hidaka, J., Shimura, Y., and Tsuchida, R., Bull. Chem. Soc., Japan, 31, 921 (1958).
24. Hidaka, J., Shimura, Y., and Tsuchida, R., Bull. Chem. Soc. Japan, 33, 847 (1960).
25. Hidaka, J., Shimura, Y., and Tsuchida, R., Bull. Chem. Soc. Japan, 35, 567 (1962).

26. Shimura, Y., Bull. Chem. Soc. Japan, 31, 315 (1958).
27. Albinak, M. J., Bhatnagar, P. C., Kirschner, S., and Sonnessa, A. J., Can. J. Chem., 39, 2360 (1961).
28. Larsson, R., Proc. 7th Int. Conf. on Coord. Chem., Paper 7B2, 326 (1962).
29. Kling, O., and Woldbye, F., Acta Chem. Scand., 15, 704 (1961).
30. Woldbye, F., Proc. 7th Int. Conf. on Coord. Chem., Paper 1E1, 41 (1962).
31. Eichorn, G. L. and Oshahr, A. J., "Advances in the Chemistry of the Coordination Compounds", S. Kirschner, Ed., The Macmillan Company, New York, 1961, pp. 216-223.
32. Ulmer, D. D., Li, T. K., and Valles, B. L., Proc. 7th Int. Conf. on Coord. Chem., Paper 3D2, 129 (1962).
33. Jaeger, F. M., "Optical Activity and High Temperature Measurements", McGraw-Hill Book Company, Inc., New York, 1930, pp. 215-234.

FAR INFRA-RED ABSORPTION OF TRANSITION-METAL COMPLEXES

Michael R. Rosenthal

October 16, 1962

INTRODUCTION

In the last 10 years, a great deal of investigation has been carried out in the far infra-red region. This region, extending from approximately 20-200 μ (50-500 cm^{-1}), had been relatively little investigated earlier owing to instrumental difficulties. These difficulties included low energy, atmospheric absorption, and stray radiation. Raman data were used along with infra-red data in the "fundamental" region for pertinent problems. Fortunately, techniques have now been developed which allow accurate spectra to be determined rapidly in this region. The information gathered has been particularly valuable in structure determinations, bonding studies, and in developing analytical techniques. The far infra-red is much more sensitive to changes in structure than the "fundamental" region, often allowing the difference of only a methylene group to be detected. Heavy-atom stretching frequencies and bending frequencies are usually found in this region. The purpose of this discussion is to review the uses and limitations of the far infra-red and to illustrate its applications to transition-metal chemistry.

APPARATUS

The development of the double-beam double-pass instrument has almost eliminated the problem of stray radiation. The drawback, however, is that this produces a greater energy loss than a single-beam instrument. Flushing with dry nitrogen has eliminated the problem of atmospheric water-vapor absorption. References (1) and (2) discuss apparatus in detail.

Grating spectrometers are useful for very low wavelength measurements (below 250 cm^{-1}). Duncanson, (4), discusses the construction and use of such an instrument. Orr(5) discusses recent advances in infra-red instrumentation.

PHASE

Carbonyl compounds and cyclopentadienyl complexes are generally observed in the gas phase. The procedure usually comprises sealing the compound in vacuo in gas cells and placing in a heating jacket(1).

More commonly encountered are liquid or solution spectra, the latter of which are best obtained with a double-beam instrument, using the pure solvent as the reference. However, if solvent absorption is high (75%), only small differences between large numbers result and much difficulty follows. Well-matched cells are an important criterion. The use of the wedge cell has helped alleviate this difficulty. Reference (2) contains detailed information on solvents.

Solid-state techniques enjoy wide popularity. Mulling the compound in mineral oil or hexachlorobutadiene is common. A drawback to the mull method is that energy tends to fall off at shorter wavelengths (Tyndall Effect). The sample may also be pressed into KBr, CsBr, or CsI discs. Meloche and Kalbus(3) discuss sources of error in this method as a result of anion exchange, pressure, and orientation effects. Occasionally either a powder or film on halide windows or single-crystal work is reported.

It should be kept in mind that shifts due to physical state are larger in this region than at higher wave numbers.

OPTICS (1,2)

To be useful, a prism must transmit the radiation used and give high dispersion. That is, the rate of change of refractive index with wavelength must be large. The following is a table of approximate values of prism ranges

KBr	15-27 μ	370-665 cm.^{-1}
CsBr	24-38 μ	265-415 cm.^{-1}
TlBr } KRS-5	24-40 μ	250-415 cm.^{-1}
TlI }		
CsI	24-50 μ	200 cm.^{-1} -415 cm.^{-1}
-Grating	50-200 μ	50-200 cm.^{-1}

VIBRATIONAL SPECTRA IN DIATOMIC MOLECULES (1)

Before proceeding to specific data, it seems worthwhile to discuss briefly the theoretical considerations of vibrational spectra. A diatomic molecule behaves as a harmonic oscillator in which

$$E = Hr \left(n + \frac{1}{2} \right)$$

where n is the vibrational quantum number. As the molecule vibrates, the dipole moment vector changes with the same frequency. If the molecule is bathed with radiation, the dipole moment vector will "lock in" with the electromagnetic vector of the same frequency, thus allowing for absorption and emission. The process may be expressed as follows:

$$\Delta E = hr \left(n + 1 + \frac{1}{2} \right) - hr \left(n + \frac{1}{2} \right) = hr$$

Molecular vibrations are found in the infra red from 1-200 μ . (50-10,000 cm.^{-1}) If as the molecule vibrates, polarizability changes with the frequency of the vibration, it will show the Raman effect.

The procedure is to bathe the molecule with monochromatic radiation of a frequency much higher than its vibrational frequency. Most of the radiation will be scattered, but now and then a

Solid-state techniques enjoy wide popularity. Mulling the compound in mineral oil or hexachlorobutadiene is common. A drawback to the mull method is that energy tends to fall off at shorter wavelengths (Tyndall Effect). The sample may also be pressed into KBr, CsBr, or CsI discs. Meloche and Kalbus(3) discuss sources of error in this method as a result of anion exchange, pressure, and orientation effects. Occasionally either a powder or film on halide windows or single-crystal work is reported.

It should be kept in mind that shifts due to physical state are larger in this region than at higher wave numbers.

OPTICS (1,2)

To be useful, a prism must transmit the radiation used and give high dispersion. That is, the rate of change of refractive index with wavelength must be large. The following is a table of approximate values of prism ranges

KBr	15-27 μ	370-665 cm.^{-1}
CsBr	24-38 μ	265-415 cm.^{-1}
TlBr } KRS-5	24-40 μ	250-415 cm.^{-1}
TlI }		
CsI	24-50 μ	200 cm.^{-1} -415 cm.^{-1}
-Grating	50-200 μ	50-200 cm.^{-1}

VIBRATIONAL SPECTRA IN DIATOMIC MOLECULES (1)

Before proceeding to specific data, it seems worthwhile to discuss briefly the theoretical considerations of vibrational spectra. A diatomic molecule behaves as a harmonic oscillator in which

$$E = Hr \left(n + \frac{1}{2} \right)$$

where n is the vibrational quantum number. As the molecule vibrates, the dipole moment vector changes with the same frequency. If the molecule is bathed with radiation, the dipole moment vector will "lock in" with the electromagnetic vector of the same frequency, thus allowing for absorption and emission. The process may be expressed as follows:

$$\Delta E = hr \left(n + 1 + \frac{1}{2} \right) - hr \left(n + \frac{1}{2} \right) = hr$$

Molecular vibrations are found in the infra red from 1-200 μ . (50-10,000 cm.^{-1}) If as the molecule vibrates, polarizability changes with the frequency of the vibration, it will show the Raman effect.

The procedure is to bathe the molecule with monochromatic radiation of a frequency much higher than its vibrational frequency. Most of the radiation will be scattered, but now and then a

quantum will be absorbed, and the molecule will go to a higher energy state. It will then re-emit the energy as a quantum of lower energy than the incident radiation. This technique is valuable for some modes of vibration.

Although emission spectra are useful, we usually are more interested in absorption spectra resulting because the Boltzmann distribution favors the lower state. Overtones appear if Δn is larger than 1, and we try to avoid these by proper sample preparation. For a diatomic molecule

$$r = \frac{1}{c} \left(\frac{K}{4\pi^2 \mu} \right)$$

where μ = reduced mass, K = bond force constant, c = velocity of light, and r = oscillator frequency.

VIBRATIONAL SPECTRA OF POLYATOMIC MOLECULES (1)

The problem becomes much more complicated when a polyatomic system is considered. If there are n atoms, there are $3n$ degrees of freedom, of which 3 are translational and 3 (non-linear) or 2 (linear) are rotational. Since these 6 (or 5) do not change the relative positions of the atoms there are $3n-5$ (linear) or $3n-6$ (non-linear) normal vibrations in a molecule of n atoms. A normal vibration is formally defined as one in which there is no change in the center of mass or angular momentum of the molecule. If an experimental frequency is known for a vibration along with molecular parameters, the bond force constant may be calculated. Herzberg (6) discusses this situation.

The mass effect is worthy of mention. If co-ordination occurs through the apex atom of a triatomic molecule, the motion of that atom will be restricted. Therefore, the frequency shift upon complexation will be greatest for that atom which moves more when uncomplexed. For this reason, H_2O exhibits a small shift when compared to NO_2 . This is due to the fact that N motion in NO_2 is greater than O motion in water, which in turn depends on the fact that H (in H_2O) is so much lighter than O (in NO_2).

Group frequencies are valuable in assigning vibrations (101). However, we must realize that often interactions will occur between vibrations of the same symmetry (coupling). These lead to unexpected values. In many cases, a "pure" stretching or bending frequency will not exist. If, however, a bond is very strong, coupling may not occur even if symmetry permits. Carbonyl and nitrile frequencies illustrate this point.

As a molecule becomes larger, analysis becomes more and more complicated. Purcell(7) and references therein apply group theoretical analysis to complicated molecules.

VIBRATIONAL SPECTRA OF COMPLEXES (1)

Several empirical rules may often be applied carefully to spectral analysis of complexes: (a) Ligand frequencies and force constants will not usually vary greatly from free to bound

ligand; (b) Vibrations of bound ligands will not normally couple strongly to vibrations of other ligands or to skeletal vibrations; (c) Slight coupling of ligand vibrations to other molecular vibrations will not usually destroy degeneracies inherent in the free ligand.

OTHER REFERENCES

Lawson(8) contains sparse information on this subject. Bentley and Wolfarth (9) list spectra-structure correlations for a large number of aliphatic and aromatic hydrocarbons. Miller, et al (10), list spectra-structure correlations for 208 inorganic salts containing polyatomic ions. Ferraro (49) discusses inorganic infra-red spectroscopy, though little of the material relates to the far infra-red.

Tabulated below is far infra-red data on transition-metal complexes. Because of space limitations, interpretation of this data is not included. This may be secured from the listed references.

Class and Reference	Compound	Vibrational (cm ⁻¹)	Assignment	Phase	Crystal	
Thiocyanate(11)	(NH ₄) ₃ V(NCS) ₆ ·4H ₂ O	478	NCS bend	mull,	KBr	
	K ₃ Cr(NCS) ₆ ·4H ₂ O	476		solution		
	NH ₄ Cr(NH ₃) ₂ (NCS) ₄	466				
	K ₂ Co(NCS) ₄ ·4H ₂ O	475				
	K ₄ Ni(NCS) ₆ ·4H ₂ O	469				
	K ₂ Zn(NCS) ₄ ·4H ₂ O	479, 470				
	K ₃ Rh(SCN) ₆	471, 452, 438				
	Tl ₃ Rh(SCN) ₆	455, 428				
	K ₂ Pd(SCN) ₄	473, 465, 439, 430				
	K ₂ Cd(SCN) ₄ ·2H ₂ O	456, 453, 453, 417				
	K ₂ Pt(SCN) ₆	454, 426, 415				
	K ₂ Hg(SCN) ₄	451, 457, 436, 431, 416				
	K ₃ Mo(CNS) ₆ ·4H ₂ O	476				
	(NH ₄) ₃ Mo(CNS) ₆ ·HCl·H ₂ O	474				
Tetra(p-chlorophenyl) porphine (12)	Uncomplexed	491	C-Cl stretch	KBr	KBr	
	V complex	498		disc		
	Cu "	502				
	Co "	503.5				
	Ni "	504.5				
	Zn "	497.5				
Oxalato(13)	K ₂ ZnOx ₂ ·2H ₂ O	(a) 519, 428, 419 (b) 420, 417 (c) 377, 364, 382, 370, 358, 370, 340, 336, 358, 364, 364, 347	(d) 291, 339, 350, 328, -	(a) ring def. + CO ₂ bend (b) MO str. + ring def. (c) CO str. + CO ₂ bend (d) ?	mull	CsBr
	K ₂ CuOx ₂ ·2H ₂ O	481, 420				
	K ₂ PdOx ₂ ·2H ₂ O	469, 417				
	K ₂ PtOx ₂ ·2H ₂ O	469, 405				
	K ₃ FeOx ₃ ·3H ₂ O	498, 366				
	K ₃ VOx ₃ ·3H ₂ O	497, 368				
	K ₃ CrOx ₃ ·3H ₂ O	485, 415				
	K ₃ CoOx ₃ ·3H ₂ O	472, 446				
	K ₃ AlOx ₃ ·3H ₂ O	436, 485				
	Cr(NH ₃) ₄ OxCl	486, 496, 366				
(14)	Pd(NH ₃) ₂ Ox	374, 363, 299	none	KBr	CsBr	
	K ₂ [PdOx ₂]·2H ₂ O	374, 357		disc		
	Na ₂ [CuOx ₂ (H ₂ O) ₂]	387, 346				
	Na[CroX ₂ (H ₂ O) ₂]·5H ₂ O	398, 315				
	K[CroX ₂ (H ₂ O) ₂]·3H ₂ O	399, 315				

Class and Reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Crystals
2-(0-hydroxyphenyl)-imidazoline (16)	Cu ⁺ ₂ complex	384	M-N str.	KBr disc	CsBr
	Zn ⁺ ₂ complex	374			
	Ni ⁺ ₂ complex	336 or 367			
2-(0-hydroxyphenyl)benzimidazole (16)	Cu ⁺ ₂ complex	355	M-N str.	KBr disc	CsBr
	Zn ⁺ ₂ complex	367			
	Co ⁺ ₂ complex	378			
Pyridine (17)	Py	403	ring vibration	mull	KBr
	MnpyCl ₂	417			
	MnpyBr ₂	424, 419			
	MnpyI ₂	423, 418			
	FePy ₄ (NCS) ₂	426, 416			
	CoPy ₂ Cl ₂	427			
	CoPy ₂ Br ₂	418			
	CoPy ₂ I ₂	423, 415			
	CoPy ₂ (NCS) ₂	422, 418			
	CoPy ₄ (NCS) ₂	426, 417			
	NiPy ₂ Cl ₂	437			
	NiPy ₄ (NCS) ₂	440			
	NiPy ₄ Br ₂	427			
	NiPy ₄ I ₂	430, 417			
	CuPy ₂ Cl ₂	440			
	CuPy ₂ Br ₂	439			
	CuPy ₂ (NCS) ₂	416			
	CuPy ₄ (BF ₄) ₂	433			
	AsPy ₂ BF ₄	421			
	ZnPy ₂ Cl ₂	417			
	ZnPy ₂ I ₂	420			
	ZnPy ₂ (NCS) ₂	422, 410			
	CdPy ₂ (NCS) ₂	416, 413			
	CdPy ₄ (BF ₄) ₂	412 ^φ			
	HgPy ₂ (BF ₄) ₂	425			
	MnPy ₂ Cl ₂	417			
	CoPy ₂ Cl ₂	427			
NiPy ₂ Cl ₂	437				
CuPy ₂ Cl ₂	440				
MnPy ₂ Cl ₂	422				
CuPy ₂ Br ₂	439				

Class and Reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Optics
Fyridinium Salts (17)	(pyH) ₂ MnCl ₄	413	none	mull	CsBr
	(pyH) ₂ CoCl ₄	414			
Unidentate Carbonate (18)	[Co(NH ₃) ₅ CO ₃]Br	362	Co-O str.	mull	CsBr
	[Co(ND ₃) ₅ CO ₃]Br	351			
	[Co(NH ₃) ₅ CO ₃]I	360			
	[Co(ND ₃) ₅ CO ₃]I	341			
	[Co(NH ₃) ₅ CO ₃]NO ₃ · 1/2 H ₂ O	351			
	[Co(ND ₃) ₅ CO ₃]NO ₃ · 1/2 H ₂ O	344			
	[Co(NH ₃) ₅ CO ₃]ClO ₄	350			
	[Co(NH ₃) ₅ CO ₃]ClO ₄	338			
	[Co(NH ₃) ₄ CO ₃]Cl	(a) 395 (b) 430	(a) CoO str. (b) Co-O bend	mull	CsBr
	deuterated [Co(NH ₃) ₄ CO ₃]ClO ₄	378 392			
	deuterated [Coen ₂ CO ₃]Br	374 353			
deuterated [Coen ₂ CO ₃]ClO ₄	372 362				
deuterated KCo(NH ₃) ₂ (CO ₃) ₂	348 366				
deuterated [Co(NH ₃) ₆][Co(CO ₃) ₃]	362 394, 391				
Ethylenediamine (19)	Pten ₂ + ₂	(a) 533 (b) ?	(a) M-N str. (b) log K av	mull	KBr
	Rhen ₂ + ₃	580			
	Coen ₃ + ₂	585	16.3		
	Pden ₂ + ₃	573	13.5		
	Cren ₃ + ₂	567	?		
	Hgen ₂ + ₂	?	11.7		
	Cuen ₂ + ₂	538	9.8		
	Nien ₃ + ₂	515	6.0		
	Coen ₃ + ₂	502	4.6		
	Znen ₃ + ₂	478	4.0		
	Cden ₃	478	4.0		
(20)	[Pten ₂]Cl ₂	(a) 478(455) (b) 589, 546 (509)	(a) ring bend(?) (b) MN str.	liq. para- ffin	KBr
	[Pten ₂]PtCl ₄	466(445) 583, 566 538			

Class and Reference	Compound	Vibration(cm^{-1})	Assignment	Phase	Other
	[Pt _{en} Cl ₂]	(536, 523, 497)	parentheses		
	[Pd _{en} Cl ₂]	466(445)	denote deuterated bands		
	[Pd _{en}] ₂ PtCl ₄	472(451)			
	[Pd _{en}] ₂ PtCl ₄	457			
	[Pd _{en}] ₂ Cl ₂	448			
	[Cu _{en}] ₂ PtCl ₄	-			
	[Ni _{en}] ₃ PtCl ₄	-			
Titanates	BaTiO ₃	550	TiO ₄ octa-hedra vibration	KBr disc	KBr
Ferrites (23)	CoFe ₂ O ₄	374			
	FeFe ₂ O ₄	370-80	r ₂ -a complicated vibration	KBr disc	KRS-5
	MgFe ₂ O ₄	406			
	MnFe ₂ O ₄	392			
	NiFe ₂ O ₄	396			
	ZnFe ₂ O ₄	393			
	Ni _{0.3} Zn _{0.7} Fe ₂ O ₄	409			
Manganites (24)	NiMn ₂ O ₄	465 broad			
	CuMn ₂ O ₄	465 sharp	r ₂	KBr disc	KBr
Ethylene (25)	[Pd(C ₂ H ₄)Cl ₂] ₂	415 weak	may be M-ethylene	mull	KBr
	[Pt(C ₂ H ₄)Cl ₂] ₂	not investigated			
	K[Pt(C ₂ H ₄)Cl ₃].H ₂ O	405 very weak			
	K[Pt(C ₂ H ₄)Cl ₃]	-			
	trans-[Pt(C ₂ H ₄)Cl ₂ (NH ₃)]	-			
1:5 cyclo octadiene (26)	CuCl ₂ .C ₈ H ₁₂	455	none	?	?
	PdCl ₂ .C ₈ H ₁₂	450			
RuO ₄ (27)	RuO ₄	330	Ru-O str.	liquid film	CsBr foreprism; grating

Class and reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Optical
Nitrito	ONO ⁻	(a) 1335 (b) 1250 (c) -	(a) asymm. N-O str. (b) symm. N-O str. (c) a-b	mull	KBr
	Nitro (28,29)	K ₂ Ca[Ni(NO ₂) ₆]	1355 1325		30
K ₃ [Co(NO ₂) ₆]		1396, 1381 1332		55	
K ₂ [Pt(NO ₂) ₄]		1436, 1410 1350 1336		61	
CH ₃ NO ₂		1582 1384		-	
Sandwich Compds. (30-33)	K ₃ [Co(NO ₂) ₆]	413			
	K ₂ [Pt(NO ₂) ₆]	450			
	[Co(NH ₃) ₆] ⁺³	(a) 502 (b) 1	(a) Co-NH ₃ str. (b) no predicted bands	mull	KBr
	[Co(NH ₃) ₅ NO ₂] ⁺²	513, 499, 488		3	
	1,6-[Co(NH ₃) ₄ (NO ₂) ₂] ⁺²	511, 504		1	
	1,2-[Co(NH ₃) ₄ (NO ₂) ₂] ⁺²	514, 483, 469		4	
	1,2,4-[Co(NH ₃) ₃ (NO ₂) ₃] ⁺²	481, 464		2	
	1,2,3-[Co(NH ₃) ₂ (NO ₂) ₄] ⁺²	507, 438		1	
	1,6-[Co(NH ₃) ₂ (NO ₂) ₄] ⁺²	-		-	
	[Co(NH ₃) ₅ NO ₂] ⁺²	499			
	[Co(NH ₃) ₅ ONO] ⁺²	?			
	[Cr(NH ₃) ₅ ONO] ⁺²	476, 456, 444			
	[(NH ₃) ₃ Co(OH)(NO ₂)-Co(NH ₃) ₃] ⁺³	500			
Sandwich Compds. (30-33)	[Cr(NH ₃) ₆] ⁺³	455			
	[Cr(NH ₃) ₅ Cl] ⁺²	470, 460, 431			
	[Cr(NH ₃) ₅ ONO] ⁺²	476, 466, 444			
	[Cr(NH ₃) ₅ N ₃] ⁺²	448			
	B[Cr(NH ₃) ₅ NO ₂] ⁺²	513, 499, 488			
Sandwich Compds. (30-33)	Cp ₂ Fe	(a) 478 (b) 492	(a) antisymm. ring metal str. (b) antisymm. ring tilt	mull	KBr
	Cp ₂ Ni	355	antisymm. ring metal str.	mull	KBr
	Cp ₂ Ru	446	antisymm. ring metal str.	mull	KBr

*reference
*ring-metal
*ring bend
*Calculated

Class and Reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Comments
Organo Pd and Pt (34)	(C ₆ H ₆) ₂ Cr	456 (a) 492 (b)	(a) skeletal str.	sublimed	Csbr
	deuterated	421	(b) skeletal tilt	film	
	(PEt ₃) ₂ Pd me Br	510	M-CH ₃	KI disc	Grubb Farsons GS2A
	(PEt ₃) ₂ Pd me (SCN)	526	str.		
	(PEt ₃) ₂ Pd me (CN)	502			
	(PEt ₃) ₂ Pd me ₂	491, 457			
	(AsEt ₃) ₂ Pdme ₂	498, 479			
	(Pbenz ₃) ₂ Pd me ₂	529, 482			
	bipy Pd me ₂	534, 522			
	cis-(PEt ₃) ₂ Ptme ₂	523, 506			
(35)	cis-PtCl ₂ (Pme ₃) ₂	689, 630		mull,	?
	cis-PtCl ₂ (Pet ₃) ₂	643, 637		solution	
	trans-PtCl ₂ (Pet ₃) ₂	632			
	cis-Ptme ₂ (Pme ₃) ₂	(a) 525, 508	(b) 678, 671	(a) Pt-C str in benzene	?
	cis-Ptet ₂ (Pme ₃) ₂	511, 500	678, 672	(b) Pt-C	
	cis-Ptpr ₂ (Pme ₃) ₂	609, 601	676, 669	str. in	
	cis-Ptme ₂ (Pet ₃) ₂	526, 506	641, 629	nujol mull	
	cis-Ptet ₂ (Pet ₃) ₂	516, 496	-		
	Ptme ₂ (Et ₂ PCH ₂ CH ₂ PEt ₂)	521, 512	625		
	Ptme ₂ (Et ₂ SCH ₂ CH ₂ SEt ₂)	555, 548	-		
	trans-Pt(NO ₃)me(PR ₃) ₂	(a) 567	(b) 566	(c) 675	(d) 630
	trans-PtClme(PR ₃) ₂	551	551	675	630
	trans-PtBrme(PR ₃) ₂	546	548	675	633
	trans-PtI ₂ me(PR ₃) ₂	538	541	675	633
	trans-Pt(NO ₂)me(PR ₃) ₂	545	544	676	633
	trans-Pt(SCN)me(PR ₃) ₂	558	556	675	626
	trans-Pt(CN)me(PR ₃) ₂	-	516	-	637

Class and reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Optics
	cis-PtCl ₂ me ₂ (PR ₃) ₂	(a) 527 (b) 686, 675 (c) 642, 630	Pt-C str. (a) R=et benzene	-	?
	trans-PtCl ₂ et(PR ₃) ₂	538	(b)R=me		
	trans-PtIet(PR ₃) ₂	522	nujol		
	trans-PtCl ₂ pr(PR ₃) ₂	- 674	(c)R=et nujol		
	PtI ₂ me ₂ (PET ₃) ₂	(a) 536 (b) 540	Pt-C str. (a) nujol		
	PtCl ₂ me ₂ (PET ₃) ₂	545, 456	(b) benzene		
	PtI ₂ me ₂ (Ppr ₃) ₂	525, 452			
(36)	cis-Pt(CH ₃) ₂ (Pme ₃) ₂	477, 466	Pt-C	benzene	Grubb Parsons GS2A
	cis-Pt(CD ₃) ₂ (Pme ₃) ₂	522, 510	str.		
	trans-PtCl(CH ₃)(Pme ₃) ₂	516			
	trans-PtCl(CD ₃)(Pme ₃) ₂	563			
(37)	trans(CH ₃ CO)Pt(PET ₃) ₂ X	(a) 541 (b) 583 545 583 539 587 537 584 532 580 554 581	(a) PtC str. (b) skeletal	CCl ₄ , mull	Grubb Parsons GS2A
	X = NO ₃				
	NCS				
	Cl				
	Br				
	I				
	NO ₂				
	trans-(CH ₃ CO)Pd(PET ₃) ₂ X	(a) 521 (b) 577 520 576 515 572 513 571 509 570 517 575	(a) Pd-C str. (b) skeletal	CCl ₄ , mull's	Grubb Parsons GS2A
	X = NO ₃				
	NCS				
	Cl				
	Br				
	I				
	NO ₂				
Propene		(a) 370 (b) 330 (c) 306	(a) antisymm. ClPtCl str. (b) PtCl str. (c) PtCl str.	mull	Grubb Parsons GS2A
(50)	K[PtCl ₃ (C ₃ H ₆)]·H ₂ O	370			
	K[PtCl ₃ (C ₃ D ₆)]·H ₂ O	393	trans to propene		

Class and Reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase
(38)	Coacac ₃	(a) 466	(a) MO str. (b) force constant	KBr disc KBr
	Cracac ₃	(b) 2.40		
	Feacac ₃	459		
	Cuacac ₃	434		
(39)		455	(a) MO str. (b) force constant (c) log K form.	KBr disc KBr
	Znacac ₂	(a) 422		
	Coacac ₂	(b) 1.50		
	Niacac ₂	422		
	Cuacac ₂	452		
(40)	Fdacac ₂	464	MO str.	KBr disc KBr
	Niacac ₂	452		
	Niacac ₂ .py ₂	439		
	VOacac ₂	480		
(41)	VOacac ₂ .py ₂	463	No assignments but metal sensitive	CsBr
	Alacac ₃	400		
	Cracac ₃	360		
	Mnacac ₃	339		
	Feacac ₃	301		
Ammines (42)	Co(NH ₃) ₆ Cl ₂	318	Co-N str.	mull, KBr disc CsBr
	Co(NH ₃) ₆ Br ₂	317		
	Co(NH ₃) ₆ I ₂	308		
	Co(NH ₃) ₆ Cl ₃	328		
	Co(NH ₃) ₆ Br ₃	322		
	Co(NH ₃) ₆ I ₃	320		
	Hg(NH ₃) ₂ Cl ₂	513		
(43)	Hg(NH ₃) ₂ Br ₂	499	(a) Co-N str. (b) NCoN def.	mull, KBr disc CsBr
	Co(NH ₃) ₆ Br ₃	(a) 464		
	Co(NH ₃) ₆ I ₃	(b) 325		
(44)	trans-[Co(NH ₃) ₄ Cl ₂]Cl	(a) 501	(a) antisym CoX str. (b) CoN str. (c) NCoX def. (d) NCoN def.	same
	trans-[Co(NH ₃) ₄ Br ₂]Br	(b) 463		
(45)		(c) 353	(a) antisym CoX str. (b) CoN str. (c) NCoX def. (d) NCoN def.	same
		(d) 290		
(46)		318	(a) antisym CoX str. (b) CoN str. (c) NCoX def. (d) NCoN def.	same
		294		

Class and Reference	Compound	Vibration (cm ⁻¹)			Assignment	Phase	Crystals
		(a)	(b)	(c)			
(44)	[Co(NH ₃) ₅ F]Br ₂	504, 493	449	310	(a) antiphase	same	same
	[Co(NH ₃) ₅ Cl]Br ₂	487, 479	457	325, 283	CoX+CoN		
	[Co(NH ₃) ₅ Cl](NO ₃) ₂	487	462	323, 283	(b) CoN str.		
	[Co(NH ₃) ₅ Br]Cl ₂	487	462	324	(c) NCoN		
	[Co(NH ₃) ₅ I]Br ₂	428	473	319	def. and NCoX def.		
(45-47)	trans-Pt(NH ₃) ₂ Cl ₂	326			MN str.	KBr disc	CsBr
	deuterated	332					
	cis-Pt(NH ₃) ₂ Cl ₂	322					
	deuterated	322					
	trans-Pt(NH ₃) ₂ Cl ₄	300					
	Cu(NH ₃) ₂ Cl ₂	481, 306					
	Hg(NH ₃) ₂ Cl ₂	471					
	Zn(NH ₃) ₂ Cl ₂	442, 306					
	Cd(NH ₃) ₂ Cl ₂	374					
	Pt(NH ₃) ₄ Cl ₂ ·H ₂ O	497					
	Zn(NH ₃) ₄ I ₂	410					
	Cu(NH ₃) ₄ Cl ₂	420					
	Cd(NH ₃) ₄ Cl ₂	381					
	Hg(NH ₃) ₄ (ClO ₄) ₂	408					
	(48)	trans-Pt(NH ₃) ₂ Cl ₂	507			MN	mull
cis-Pt(NH ₃) ₂ Cl ₂		508			str.		
cis-Pt(NH ₃) ₂ (NO ₂)Cl		510					
trans-Pt(Et ₂ S)Cl ₂ (NH ₃) ₂		493					
trans-Pt(C ₂ H ₄)(NH ₃)Cl ₂		481					
cis-Pd(NH ₃) ₂ Cl ₂		477, 497					
[Co(NH ₃) ₅ Cl]Cl ₂		493					
deuterated		487					
trans-Pd(NH ₃) ₂ Cl ₂		496					
deuterated		493					
cis-Pd(NH ₃) ₂ (NO ₂) ₂		496, 463					
Id(NH ₃) ₄ Cl·H ₂ O		498					
[Co(NH ₃) ₆]Cl ₃		502					
It(NH ₃) ₄ Cl ₂		511					
(48)		FgNH ₂ Br cubic	510			HgN str.	KBr disc
	FgNH ₂ Br rhombic	560					

Class and Reference	Compound	Vibration (cm ⁻¹)	Assignment	Phase	Optics	Structure	Comments
Carbonyls (51,52)	Ni(CO) ₄	(a)			(a)NiC str.	Vapor	CsBr
		(b)	422	459	(b)NiCO bend		
Fe(CO) ₅		472	CO bend	gas, liquid	KBr	trig. bipy.	No 377
		522	-				364
		93	-	?	?	?	No 377
		72	-				364
(55)		364	apical FeC str.	gas	KBr	tetr. bipy.	364, 377
		377	Fe-C str.				crucial to structure
		430	in phase FeCO bend				
		475	out of phase FeCO bend				
(56)		364	Comb.	evaluates		trig. bipy.	
		377	eq FeC str.	old data			
		430	eq CO bnd				
		472	ax FeC str.				
(57)		68	γ ₉	vapor	grating	trig. bipy.	* = calc. from comb.
		114	γ ₁₄				
		104	γ ₁₅				
(57)		93	γ ₁₆	liquid	grating	trig. bipy.	
		72	γ ₉				
Fe ₃ (CO) ₁₂		575	C-Fe stretches	solution	CsBr	?	
		594					
Co ₂ (CO) ₈		531	Co-C-bend	solution	?	2 bridge carbonyls	
		641	Co-C str.			Co-Co bond	
Cr(CO) ₆		436	CrC str.	gas	KRS-5	octahedral	
		661	CO wag				
		98	CrCrC bend	gas	grating	octahedral	

Reference	Bands Found	Assignment	Phase	Optics	Structure	Comments
$\text{Mo}(\text{CO})_6$	(60) 368 595 (57) (61) 81 agrees with others	MoC str. CO wag CMoC band -	gas	KRS-5	Octahedral	
$\text{Mn}_2(\text{CO})_{10}$	(62) 625	none	solid, solution	?	bioccta- hedral (X-ray)	with M-M bond
$\text{Re}_2(\text{CO})_{10}$	(62) 455	none	same	?	same	
Carbonyl Hydrides	(59) 548	CoCO bend	solution	?	basically tetrahedral	
$\text{HCo}(\text{CO})_4$	(63) 704 330 403 467 541 703 1934 1400	CoC str. sym. CoCstr. CoC str. + CoCO bend Co bend Co bend CoC str. Co-H str. 2(CoC str.)	gas	CsBr	basically tetrahedral	
$\text{DCo}(\text{CO})_4$	(64) 600 296	Co-C str. sym. CoC str.	gas	?	basically tetrahedral	proves that C and/ O partici- pate with H or D.
$\text{HMn}(\text{CO})_5$	(65) 1396 1193	Co-D str. 2(Co-C str.)	gas	?		
$\text{HMn}(\text{CO})_5$	(64) 615 664 735 458 1782	MnH bend + CO bend CO bend MnH bend + CO bend none Mn-H str.	?	?		?

Reference	Bands Found	Assignment	Phase	Optics	Structure	Comments
$\text{DMn}(\text{CO})_6$ (64)	676 doublet 1287	CO bend + CO bend MnD str.	?	?	?	
$[\text{Co}(\text{CO})_4]^-$ (59)	555 1833	CoCo bend CO str.	pyridine	?	tetrahedral	
(67)	530	none	THF	CsBr	-	
$\text{Mn}(\text{CO})_5^-$ (67)	467 511 564	none	THF	CsBr	-	
$\text{Co}(\text{CO})_3\text{NO}$ (68)	390 442 467 483 528 565	CoC symm str. deg. CO bend CO bend symm. CO bend CoN symm str. CoC asymm. str.	gas	CsBr	tetrahedral	diffraction agrees
(69)	confirms above	-	-	-	-	tetrahedral
Class and Reference	Compound	Vibration (cm^{-1})	Assignment	Phase	Optics	
carbonyl-amine (70)	trans-Pt(CO)(p-tol)Cl ₂	488	Pt-N str.	mull	CsBr	
	trans-Pt(CO)(p-tol)Br ₂	492				
	trans-Pt(CO)(p-tol)I ₂	493				
	trans-Pt(CO)(an)Br ₂	481				
	trans-Pt(CO)(an)I ₂	482				
	trans-Pt(Et ₂ S)(NH ₃)Cl ₂	493				
Cyanide Complexes	trans-Pt(NH ₃) ₂ Cl ₂	507				
	$\text{Ni}(\text{CN})_4^{=}$	(a) 543 (b) 448 (c) 430	(a)NiC str. (b)out plane NiCN bend (c)inplane NiCN bend	KBr discs, mull, single crystal	CsBr	
(75)		(a) 304 (b) 306	(a)CuC str. (b)CuCN bend	single crystal	CsBr	
(76)	$\text{K}_3\text{Cu}(\text{CN})_4$					

Class and Reference

Class and Reference	Compound	Vibration (cm ⁻¹)		Assignment	Phase	Optics
(77)	Pt(CN) ₄ ⁼	(a) 300	(b) 505	(a)CN wag (b)PtC str.	KBr disc	CsBr
(78)	Fe(CN) ₆ ⁻³	(a) 505	(b) 2125	(a)M-C str. (b)CN str.	?	?
	Fe(CN) ₆ ⁻⁴	584	2047			
	Cr(CN) ₆ ⁻³	462	2127			
	Mn(CN) ₆ ⁻³	514	2125			
	Mn(CN) ₆ ⁻⁴	-	2060			
	Mn(CN) ₆ ⁻⁵	-	2043			
(71-73)	[Ag(CN) ₂] ⁻	390		M-C str.	mull	CsBr
	[Au(CN) ₂] ⁻	427				
	[Hg(CN) ₂] ⁻	442				
(74)	K ₂ Zn(CN) ₄	358	315	both MC str. + NCM bend	mull, crystal	CsBr
	K ₂ Cd(CN) ₄	316	-			
	K ₂ Hg(CN) ₄	330	-			
(79)	Co(CN) ₆ ⁻³	422		M-C str.	?	?
	Cr(CN) ₆ ⁻³	457				
	Mn(CN) ₆ ⁻³	443				
	Fe(CN) ₆ ⁻³	416				
(80)	K ₃ Co(CN) ₆	416		CoCN bend CoC str.	single crystal	CsBr
		564				
(81)	K ₃ [Co(CN) ₆].3H ₂ O	(a) 565	(b) 414	(a)MC str. (b)MCN bend	mull, KBr disc	CsBr fore- prism, grating
	K ₄ [Fe(CN) ₆].3H ₂ O	585	414			
	K ₃ [Fe(CN) ₆].3H ₂ O	511	387			
	K ₄ [Ru(CN) ₆].3H ₂ O	550	376			
	K ₄ [Os(CN) ₆].3H ₂ O	554	360			
	[Co(NH ₃) ₆][Co(CN) ₆]	(a) 562	(b) 328	(a)MC str. (b)NCoN def.	same	same
	[Co(NH ₃) ₆][Fe(CN) ₆]	511	330	(c)MCN bend		
			(c) 455			
			459			

Class and Reference	Compound	Vibration (cm^{-1})	Assignment	Phase	Optics
(82)	$\text{K}_4\text{Mo}(\text{CN})_8 \cdot 2\text{H}_2\text{O}$	402	none	KBr disc	CsBr
	dehydrated	388			
	$\text{K}_3\text{Mo}(\text{CN})_8 \cdot 2\text{H}_2\text{O}$	402			
	dehydrated	388			
	$\text{K}_4\text{W}(\text{CN})_8 \cdot 2\text{H}_2\text{O}$	407			
	dehydrated	400			
	$\text{K}_3\text{W}(\text{CN})_8 \cdot 2\text{H}_2\text{O}$	407			
	dehydrated	400			

References 83-99 contain material on halo compounds of transition metals. The material is omitted due to space limitations.

References

1. F. A. Cotton, "The Infra-red Spectra of Transition Metal Complexes", Modern Co-ordination Chemistry, R. J. Lewis and R. G. Wilkens, eds., Interscience, London (1960), pp. 301-400.
2. F. F. Bentley, E. F. Wolfarth, N.E. Srp, and W. R. Powell, Spectrochim. Acta, 13, 1 (1958).
3. V. W. Meloche and E. K. Kalbus, J. Inorg. Nuclear Chem., 6, 104 (1958).
4. L. A. Duncanson, J. W. Eddell, M. B. Lloyd., and W. T. Moore, Spectrochim. Acta, 15, 64 (1959).
5. S. F. D. Orr, Spectrochim. Acta, 14, 86 (1959).
6. G. Herzberg, Infra-red and Raman Spectra of Polyatomic Molecules, Van Nostrand, New York (1945), pp. 168-71.
7. K. Purcell, Inorganic Seminar, University of Illinois, July 12, 1962.
8. K. E. Lawson, Infra-red Absorption of Inorganic Substances, Reinhold, New York (1961).
9. F. F. Bentley and E. F. Wolfarth, Spectrochim. Acta, 15, 165 (1959).
10. F. A. Miller, G. L. Carlson, F. F. Bentley, and W. H. Jones, Spectrochim. Acta, 16, 135 (1960).
11. J. Lewis, R. S. Nyholm, and F. W. Smith, J. Chem. Soc., 1961, 4590.
12. J. M. Goldstein, W. M. McNabb, and J. F. Hazel, J. Am. Chem. Soc., 78, 3543 (1956).
13. J. Fujita, A. E. Martell, and K. Nakamoto, J. Chem. Phys., 36, 324 (1962).
14. M. Judith Schmelz, T. Miyazawa, S. Mizushima, T. J. Lane, and J. V. Quagliano, Spectrochim. Acta, 9, 51 (1957).
15. J. Fujita, A. E. Martell, and K. Nakamoto, J. Chem. Phys., 36, 331 (1962).
16. T. J. Lane, I. Nakagawa, J. L. Walter, and A. J. Kandathil, Inorg. Chem., 1, 267 (1962).
17. N. S. Gill, R. H. Nuttall, D. E. Scaife, and D. W.A. Sharp, J. Inorg. Nuclear Chem., 18, 79 (1961).
18. J. Fujita, A. E. Martell, and K. Nakamoto, J. Chem. Phys., 36, 339 (1962).
19. D. B. Powell and N. Sheppard, J. Chem. Soc., 1961, 1112.
20. D. B. Powell and N. Sheppard, Spectrochim. Acta, 17, 68 (1961).
21. H. D. Megaw, Acta Cryst., 5, 739 (1952).

22. R. T. Mara, G. B. B. M. Sutherland, and H. V. Tyrell, *Phys. Rev.*, 96, 801 (1954).
23. R. D. Waldron, *Phys. Rev.*, 99, 1727 (1955).
24. S. Dasgupta and A. P. B. Sinha, *Trans. Faraday Soc.*, 53, 909 (1957).
25. D. B. Powell and N. Sheppard, *Spectrochim. Acta*, 13, 69 (1958).
26. P. J. Hendra and D. B. Powell, *Spectrochim. Acta*, 17, 913 (1961).
27. M. H. Ortner, *J. Chem. Phys.*, 34, 556 (1961).
28. K. Nakamoto, J. Fujita, and H. Murata, *J. Am. Chem. Soc.*, 80, 4817 (1958).
29. I. R. Beattie and H. J. V. Tyrell, *J. Chem. Soc.*, 1956, 2849.
30. E. R. Lippincott and R. D. Nelson, *J. Am. Chem. Soc.*, 77, 4990 (1955).
31. E. R. Lippincott and R. D. Nelson, *Spectrochim. Acta*, 10, 307 (1958).
32. W. K. Winter, B. Curnette Jr., S. E. Whitcomb, *Spectrochim. Acta*, 12, 1085 (1959).
33. R. G. Snyder, *Spectrochim. Acta*, 15, 807 (1959).
34. G. Calvin and G. E. Coates, *J. Chem. Soc.*, 1960, 2008.
35. D. M. Adams, J. Chatt, and B. L. Shaw, *J. Chem. Soc.*, 1960, 2047.
36. D. M. Adams, *J. Chem. Soc.*, 1962, 1220.
37. D. M. Adams and G. Booth, *J. Chem. Soc.*, 1962, 1112.
38. K. Nakamoto, P. J. McCarthy, A. Ruby, and A. E. Martell, *J. Am. Chem. Soc.*, 83, 1066 (1961).
39. K. Nakamoto, P. J. McCarthy, A. E. Martell, *J. Am. Chem. Soc.*, 83, 1272 (1961).
40. K. Nakamoto, Y. Morimoto, A. E. Martell, *J. Am. Chem. Soc.*, 83, 4533 (1961).
41. K. E. Lawson, *Spectrochim. Acta*, 17, 248 (1961).
42. E. P. Bertin, I. Nakagawa, S. Mizushima, T. J. Lane, and J. V. Quagliano, *J. Am. Chem. Soc.*, 80, 525 (1958).
43. T. Shimanouchi and I. Nakagawa, *Spectrochim. Acta*, 18, 89 (1962).
44. G. M. Barrow, R. H. Krueger, and F. Basolo, *J. Inorg. Nuclear Chem.* 2, 340 (1956).
45. D. B. Powell, *Chem. and Ind.*, 1956, 314.
46. D. B. Powell, *J. Chem. Soc.*, 1956, 4495.
47. D. B. Powell and N. Sheppard, *J. Chem. Soc.*, 1956, 3108.
48. I. Nakagawa, R. B. Penland, S. Mizushima, T. J. Lane, and J. V. Quagliano, *Spectrochim. Acta*, 9, 199 (1957).
49. J. R. Ferraro, *J. Chem. Ed.*, 38, 101 (1961).
50. D. M. Adams and J. Chatt, *J. Chem. Soc.*, 1962, 2821.
51. L. H. Jones, *J. Chem. Phys.*, 28, 1215 (1958).
52. L. H. Jones, *J. Chem. Phys.*, 23, 2448 (1955).
53. R. K. Shelton and K. S. Pitzer, *J. Am. Chem. Soc.*, 72, 1107 (1950).
54. F. A. Cotton, A. Danti, J. S. Waugh, and R. W. Fessenden, *J. Chem. Phys.*, 29, 1427 (1958).
55. M. F. O'Dwyer, *J. Mol. Spect.*, 2, 144 (1958).
56. C. W. F. T. Pistorius and P. C. Haarhoff, *J. Chem. Phys.*, 31, 1439 (1959).
57. R. S. McDowell and L. H. Jones, *J. Chem. Phys.*, 36, 3321 (1962).

58. F. A. Cotton and G. Wilkinson, J. Am. Chem. Soc., 79, 752 (1957).
59. R. A. Friedel, I. Wender, S. L. Shufler, and H. W. Sternberg, J. Am. Chem. Soc., 77, 3951 (1955).
60. N. J. Hawkins, H. C. Mattraw, W. W. Sabol, and D. R. Carpenter, J. Chem. Phys., 23, 2422 (1955).
61. L. H. Jones, J. Chem. Phys., 36, 2375 (1962).
62. E. O. Brimm, M. A. Lynch, Jr., and W. J. Sesny, J. Am. Chem. Soc., 76, 3831 (1954).
63. W. F. Edgell, C. B. Magee, and G. J. Gallup, J. Am. Chem. Soc., 78, 4185 (1956).
64. W. F. Edgell, G. Asato, W. Wilson, and C. Angell, J. Am. Chem. Soc., 81, 2022 (1959).
65. W. F. Edgell and R. Summitt, J. Am. Chem. Soc., 83, 1772 (1961).
66. F. A. Cotton, J. L. Dawn, and G. Wilkinson, J. Chem. Soc., 1959, 833.
67. W. F. Edgell, J. Huff, J. Thomas, H. Lehman, C. Angell, and G. Asato, J. Am. Chem. Soc. 82, 1254 (1960).
68. R. S. McDowell, W. D. Horrocks Jr., and J. T. Yates, J. Chem. Phys., 34, 530 (1961).
69. C. B. Magee, Ph.D. thesis, Purdue University, through (1).
70. R. J. Irving and E. A. Magnusson, J. Chem. Soc., 1958, 2283.
71. L. H. Jones, J. Chem. Phys., 26, 1578 (1957).
72. L. H. Jones, J. Chem. Phys., 27, 468 (1957).
73. L. H. Jones, J. Chem. Phys., 27, 665 (1957).
74. L. H. Jones, Spectrochim. Acta, 17, 188 (1961).
75. R. L. McCullough, L. H. Jones, and G. A. Crosby, Spectrochim. Acta 16, 929 (1960).
76. L. H. Jones, J. Chem. Phys. 29, 463 (1958).
77. D. M. Sweeny, I. Nakagawa, S. Mizushima, and J. V. Quagliano, J. Am. Chem. Soc., 78, 389 (1956).
78. V. Caglioti, G. Sartori and M. Scrocco, J. Inorg. Nuclear Chem., 8, 87 (1958).
79. V. Caglioti, G. Sartori, and C. Furlani, J. Inorg. Nuclear Chem., 13, 22 (1960).
80. L. H. Jones, J. Chem. Phys., 36, 1209 (1962).
81. I. Nakagawa and T. Shimanouchi, Spectrochim. Acta, 18, 101 (1962).
82. E. G. Brame Jr., F. A. Johnson, E. M. Larson, and V. W. Meloche, J. Inorg. Nuclear Chem., 6, 99 (1958).
83. T. G. Burke, D. F. Smith, and A. H. Nielson, J. Chem. Phys., 20, 447 (1952).
84. J. Gaunt, Trans. Faraday Soc., 50, 209 (1954).
85. H. C. Mattraw, N. J. Hawkins, D. R. Carpenter, and W. W. Sabol, J. Chem. Phys., 23, 985 (1955).
86. M. D. Myers and F. A. Cotton, J. Am. Chem. Soc., 82, 5027 (1960).
87. C. L. Chernick, H. H. Claasen, and B. Weinstock, J. Am. Chem. Soc., 83, 3166 (1961).
88. B. Weinstock, H. H. Claasen, and J. G. Malm, J. Chem. Phys., 32, 181 (1960).
89. J. L. Boston and D. W. A. Sharp, J. Chem. Soc., 1960, 908.
90. H. H. Claasen, H. Selig, and J. G. Malm, J. Chem. Phys., 36, 2888 (1962).
91. H. H. Claasen, J. G. Malm, and H. Selig, J. Chem. Phys., 36, 2890 (1962).
92. R. D. Peacock and D. W. A. Sharp, J. Chem. Soc., 1959, 2762.
93. J. R. Geichman, E. A. Smith, S. S. Trond, P. R. Ogle, Inorg. Chem., 1, 661 (1962).

94. A. Buchler, J. B. Berkowitz-Mattuck, and D. Dugre, J. Chem. Phys. 34, 2202 (1961).
95. J. K. Wilmhurst, J. Mol. Spect., 5, 343 (1960).
96. F. A. Miller and G. L. Carlson, Spectrochim. Acta, 16, 6 (1960).
97. M. F. A. Dore, J. A. Creighton, and L. A. Woodward, Spectrochim. Acta, 18, 267 (1962).
98. A. Buchler, W. Klemperer, and A. G. Emslie, J. Chem. Phys. 36, 2499 (1962).
99. G. E. Leroi, T. C. James, J. T. Hougen and W. Klemperer, J. Chem. Phys., 36, 2879 (1962).
100. E. D. Palik, Private Communication, through (31).
101. L. J. Bellamy, The Infra-red Spectra of Complex Molecules, Methuen, London (1958).

A MASS-SPECTROMETRIC STUDY OF SULFUR VAPOR
from 81°C to 1,737°C.*

J. R. Marquart

October 25, 1962

INTRODUCTION

Although a number of experiments have been performed in order to ascertain the composition of sulfur vapor, there has been a great deal of disagreement as to interpretation of results. Vapor-density measurements were made as early as 1909(1) and again in 1941(2). In both instances, the data were explained on the basis of only S_8 , S_6 , and S_2 species being present. More recent work by Braune, Peter and Neveling(3) indicated that S_4 must be included in the species. This conclusion has received support from optical absorption studies(4). Knudsen cell effusion studies (5,6,7) showed that S_8 is predominate at all temperatures up to the melting point. Fouretier(8), measuring the pressure exerted on a piston, and Taillade(9), incorporating entrainment in a carrier gas, obtained vapor-pressure values somewhat larger than those found by the Knudsen experiments, but the interpretation remained unchanged. West and Menzies(10), in a careful experiment, measured the total vapor pressure above liquid sulfur by a static isoteniscope. Their data agreed well with the determination of the sulfur point (11).

Guthrie, et. al.,(12) calculated the thermodynamic properties of S_8 gas and found that K_{eq} for the reaction $4S_2(g) \rightleftharpoons S_8(g)$, as calculated from the ΔH° and ΔS° values, disagreed with that calculated directly from vapor composition results (1,3). They concluded that other vapor species than S_8 , S_6 , S_4 , and S_2 are present.

Three mass-spectrometric studies of sulfur (13,14,16) have shown that S_2 predominates over all other species, whereas a fourth study(15) has indicated that only S_8 , S_7 , S_6 , and S_5 are primaries in the vapor.

Predissociation in the emission spectrum of sulfur sets the upper limit for the dissociation energy of diatomic sulfur at 4.4-ev. However, 3.6 and 3.3 are also possible values depending upon the state of excitation of atomic sulfur(17). Gaydon(17) accepted the predissociation limit as the probable value of the dissociation energy, whereas Herzberg and Mundie(18) preferred the 3.6-ev value. Various experiments have given support to each of these values: 3.3ev. (19,20,21), 3.6 ev. (22,23,24,14), and 4.4 ev. (25,26,27,28,29,30,31).

* Work performed by J. R. Marquart and J. Berkowitz at Argonne National Laboratory, Argonne, Illinois

Large discrepancies have also occurred in the values of appearance potentials, particularly for S_2^+ . These range from about 10.7 eV (32,33) to about 8.3 eV (13).

Therefore, a study was undertaken at this laboratory to attempt to resolve the uncertainty arising in the composition and thermodynamic properties of sulfur vapor. It was decided to extend the temperature range of mass-spectrometric studies by observing the vapor composition of a number of metal sulfides which decompose to sulfur upon heating, as well as by vaporizing rhombic sulfur itself.

EXPERIMENTAL ARRANGEMENT

The arrangement of the Knudsen cell and ion source was virtually identical to that described by Berkowitz and Chupka (34). A study was made to determine materials for use as a Knudsen cell which remain inert to each specific sulfide. It was found that platinum was satisfactory for ZnS and CdS; fused-alumina lined molybdenum for S, Na_2S_2 , Sb_2S_3 , FeS, and HgS; and molybdenum for CaS and SrS. Temperature of the Knudsen cell was determined by a Pt, Pt-10%Rh thermocouple for temperatures up to 1000°C and by an optical pyrometer for higher temperatures.

EXPERIMENTAL RESULTS

In each experiment, it was found that the major ion peaks could be identified as due to multiples of sulfur and in a few instances to the ions of the metal or metal sulfide. The experimental results are summarized in Tables I through IV.

DISCUSSION

On the basis of the extrapolated vapor composition obtained in Table IV, it appears that at 400°K. and about 6.56×10^{-5} atmospheres pressure S_8 is the predominate species whereas S_6 and S_7 are also significant. The value of the appearance potential for S_2^+ is only about 0.7 eV higher for the sulfur experiment than for the higher temperature experiments, but the source of S_2^+ appears to be fragmentation of higher polymers in the electron beam. Furthermore, contrary to the results of other experimentors (13,14,16), S_2^+ was not found to be the predominate species. It should be noted, however, that Hagemann (16) reports that his data were taken by a double oven arrangement with the source at 523°K. and 10^{-5} mm. pressure. If this was the case, then the added tendency toward lower polymers at higher temperature and lower pressure makes his data agree with this experiment.

The enthalpy of dissociation of each polymer was calculated from the slope of the $\log K_{eq}$ vs. $1/T$ plot for the reaction and also, where entropy data are available, from the ΔF° of the reaction which was based on the equilibrium concentration of

Table III. It is noted that enthalpies determined by the second method are higher by one to two kcal. for the S_8 , S_6 , and S_4 reactions. But it should be remembered that the literature values of the entropies are somewhat doubtful. The dissociation energy for S_2 appears to be about 107.16 kcal., which lends support to the spectroscopically proposed values of 4.4 ev. or 101 kcal.

REFERENCES

1. G. Preuner and W. Schupp, Z. physik. Chem., 68, 129(1909).
2. W. Klemm and H. Kilian, Ibid., B49, 279 (1941).
3. H. Braune, S. Peter, and V. Neveling, Z. Naturforsch., 6a, 32 (1951).
4. H. Braune and E. Steinbacker, *ibid.*, 7a, 486 (1952).
5. K. Neumann, Z. physik. Chem., A 171, 399 416 (1934).
6. R. S. Bradley, Proc. Royal Soc. (London), A205, 553 (1951).
7. E. Rideal and P. M. Wiggins, *ibid.*, A210, 291 (1952).
8. G. Fouretier, Compt. rend., 218, 194 (1944).
9. M. Taillade, *Ibid.*, 218, 836 (1944).
10. W. A. West and A. W. C. Menzies, J. Phys. Chem., 33, 1880 (1929).
11. E. F. Mueller and H. A. Burgess, Natl. Bur. Standards Sci. Papers, 15, 163 (1919).
12. G. B. Guthries, Jr., D. W. Scott, and G. Waddington, J. Am. Chem. Soc., 76, 1488 (1954).
13. P. Bradt, F. L. Mohler, and V. H. Dibeler, J. Research Natl. Bur. Standards, 57, 223 (1956).
14. P. Goldfinger, M. Ackerman, and M. Jeunhomme, "Vaporization of Compounds and Alloys at High Temperature", ASTIA, AFOSR 5932, Brussels, January 1959, Final Technical Report, Contract AF 61(052)-19.
15. M. C. Zietz, "A Mass-Spectrometric Investigation of Sulfur Vapor as a Function of Temperature", University of California Radiation Laboratory Report UCRL-9277, June 1960.
16. R. Hagemann, "Etude de l' Hemioxyde de Soufre S_2O par Spectrometrie de Masse", Commissariat a l' Energie Atomique (France), Dept. de Physico-Chimie Report DPC-IS/SSM/62-691RH, June 1962.
17. A. G. Gaydon, "Dissociation Energies and Spectra of Diatomic Molecules", (Chapman and Hall, Ltd., London, 1953), second edition.
18. G. Herzberg and L. C. Mundie, J. Chem. Phys., 8, 263(1940).
19. W. Nernst, Z. Elektrochem., 9, 622 (1903).
20. H. von Wartenberg, Z. anorg. Chem., 56, 320 (1908).
21. R. M. Reese, V. H. Dibeler, and J. L. Franklin, J. Chem. Phys., 29, 880 (1958).
22. G. St. Pierre and J. Chipman, J. Am. Chem. Soc., 76, 4787 (1954).
23. C. L. McCabe, C. B. Alcock, and R. G. Hudson, J. Metals, 8, aIME Trans., 206, 693 (1956).
24. E. W. Dewing and F. D. Richardson, Trans. Faraday Soc., 54, 679 (1958).
25. R. Colin, P. Goldfinger, and M. Jeunhomme, Nature, 187, 408 (1960).

26. P. Goldfinger, "Vaporization of Compounds and Alloys at High Temperature", Annual Summary Report No. 3 Contract AF 61(052)225, Brussels, 28 February 1962.
27. A. D. Walsh, "The Threshold of Space", in Proc. Conf. on Chemical Aeronomy in Cambridge, Massachusetts, 1956, M. Zelikoff editor (Fergamon Press, New York, 1957).
28. L. Brewer, J. Chem. Phys., 31, 1143 (1959).
29. D. G. H. Marsden, Ibid., 31, 1144 (1959).
30. R. G. W. Norrish and G. A. Oldershaw, Proc. Royal Soc. (London), A249, 498 (1959).
31. G. St. Pierre and J. Chipman, J. Metals, 8, 1474 (1956).
32. H. D. Smyth and J. P. Blewett, Phys. Rev., 46, 276(1934).
33. L. P. Blanchard and P. LeGoff, Can. J. Chem., 35, 89 (1957).
34. J. Berkowitz and W. A. Chupka, Ann. New York acad. Sci., 79, 1073 (1960).
35. R. Horbe and O. Knacke, Z. Erzbergban u. Metallhüttenw., 8, 556 (1955).
36. C. L. McCabe, J. Metals, 6, AIME Trans., 200, 969 (1954).
37. E. Kordes and B. Rackow, Z. physik. Chem., 200, 129(1952).
38. J. R. West, Ind., Eng. Chem., 42, 713 (1950).
39. W. H. Evans and D. D. Wagman, J. Research of National Bureau of Standards, 49, 141 (1952).
40. N. W. Luft, Monatsh., 86, 474 (1955).
41. J. W. Otvos and D. F. Stevenson, J. Am. Chem. Soc., 78, 546 (1956).

Table I. Appearance Potentials (a) (electron volts) of Sulfur and Sulfide Vapor

Sample and Temp.	Sulfur	HgS	Na ₂ S ₂	CdS	ZnS	FeS	SrS	CaS
Ion	383°K	507°K	814°K	953°K	1116°K	1388°K	1934°K	1962°K.
S ⁺ (mass 32)	17.7±.5	13.7±.5	14.2±.5	14.7±.5	14.0±.2	11.5±1.0	10.357 ^(b)	10.357 ^(a)
S ₂ ⁺ (mass 64)	10.2±.5	9.7±.5	9.6±.5	9.5±.5	9.6±.4	8.5±1.0	9.31±.2 ^(b)	9.45±.2 ^(b)
S ₃ ⁺ (mass 96)	13.5±.5	10.7±.5	10.5±.5	10.0±.5				
S ₄ ⁺ (mass 128)	13.5±.5	10.7±.5						
S ₅ ⁺ (mass 160)	10.0±.5	9.7±.5						
S ₆ ⁺ (mass 192)	9.5±.5	9.5±.5						
S ₇ ⁺ (mass 224)	9.5±.5	9.7±.5						
S ₈ ⁺ (mass 256)	9.0±.5	9.7±.5						
M ⁺ (metal ion)			8.0±1.0	8.5±.5	9.6±.4		5.4±.5	5.9±.4
MS ⁺ (metal sulfide ion)							5.7±.5	6.4±.4

(a) All appearance potentials normalized by comparison to appearance potential of Hg⁺ as 10.43 except (b).

(b) Normalized relative to appearance potential of S⁺ as 10.357.

Table II. Estimated Partial Pressures of Sulfides (a) (atm)

Species	HgS 480°K	Na ₂ S ₂ 814°K	CdS 986°K	ZnS 1203°K	FeS 1318°K	SrS 1809°K	CaS 1972°K
S							
S ₂	9.03x10 ⁻⁸	1.08x10 ⁻⁵	2.21x10 ⁻⁵	3.24x10 ⁻⁵	1.22x10 ⁻⁵	4.91x10 ⁻⁶	5.61x10 ⁻⁶
S ₃	3.43x10 ⁻⁹	1.34x10 ⁻⁸	8.61x10 ⁻⁹	6.08x10 ⁻⁹			1.24x10 ⁻⁶
S ₄	4.38x10 ⁻⁹	3.45x10 ⁻⁹	3.46x10 ⁻¹⁰				
S ₅	3.66x10 ⁻⁹						
S ₆	1.64x10 ⁻⁸						
S ₇	1.56x10 ⁻⁹						
S ₈	1.96x10 ⁻⁹						
M(metal)	2.68x10 ⁻⁸	9.97x10 ⁻⁹	1.31x10 ⁻⁶	2.30x10 ⁻⁵	<5.9x10 ⁻¹⁵	1.33x10 ⁻⁷	2.17x10 ⁻⁵
MS(metal sulfide)	<1.2x10 ⁻⁹	4.27x10 ⁻¹⁰	<9.6x10 ⁻⁹	<1.5x10 ⁻¹⁰	<2.8x10 ⁻¹⁵	1.03x10 ⁻⁹	2.25x10 ⁻⁸
Total	1.50x10 ⁻⁷	1.10x10 ⁻⁵	3.55x10 ⁻⁵	5.58x10 ⁻⁵	1.22x10 ⁻⁵	4.39x10 ⁻⁷	7.85x10 ⁻⁵

(a) All partial pressures estimated by applying the constant 3.0299×10^{-12} atm/unit grid intensity deg. obtained for S₂ by evaporating a known amount of ZnS to dryness while continually monitoring mass 64. All other species were corrected for ionization cross-section on the basis of 1 for S, 1.5 for S₂, 2.0 for S₃, 2.5 for S₄, --- 4.5 for S₄ estimated on the basis of the results of Berkowitz, Tasman, and Chupka, J. Chem. Phys., (in Press) and using Otvos and Stevensons ratios (41) for the metal and metal sulfide.

Table III

RELATIVE ION INTENSITY AND COMPOSITION OF SULFUR VAPOR

Species	Relative Ion Intensity 375°K.	Estimated Partial Pres. 400°K	% 400°K
S	9.22	2.05×10^{-30} atm.	3.1×10^{-24}
S ₂	75.32	4.74×10^{-8}	7.2×10^{-2}
S ₃	17.08	1.79×10^{-8}	2.7×10^{-2}
S ₄	38.37	7.09×10^{-8}	1.08×10^{-1}
S ₅	32.82	3.04×10^{-7}	4.63×10^{-1}
S ₆	27.82	1.02×10^{-5}	15.48
S ₇	3.69	3.12×10^{-6}	4.76
S ₈	100	5.19×10^{-5}	79.09

TABLE IV. THERMODYNAMIC FUNCTIONS FOR DISSOCIATION OF SULFUR VAPOR SPECIES

gas reaction	These data		(3)		Braune et al Literature		Calc from this data and Literature	
	ΔH_{400}° kcal	$\frac{\text{Keq}}{\text{units of atm.}}$ (b)	ΔF_{400}° kcal	ΔS_{400}° cal/deg. (c)	$\Delta H_{623-1273}^{\circ}$ kcal	Literature ΔS_{400}° cal/deg.	ΔF_{400}° kcal	Literature ΔK_{400}° kcal
1/4 S ₈ -S ₂	19.89	5.58x10 ⁻⁷	11.44	21.13	23.05	25.75 ^(e)	21.74	
2/7 S ₇ -S ₂	17.63	1.77x10 ⁻⁶	10.53	17.75				
1/3 S ₆ -S ₂	17.22	2.19x10 ⁻⁶	10.36	17.15	21.25	22.43 ^(f)	19.33	
2/5 S ₅ -S ₂	14.63	1.92x10 ⁻⁶	8.63	15.00				
1/2 S ₄ -S ₂	13.07	1.78x10 ⁻⁴	6.86	15.53	14.20	18.55 ^(f)	14.28	
2/3 S ₃ -S ₂	10.23	6.92x10 ⁻³	3.95	15.70				
1/2 S ₂ -S	54.64	9.42x10 ⁻²⁷	47.63	17.52	53.58 ^(d)	13.31 ^(f)	52.95	

- (a) From slope of log Keq vs 1/T Plots.
- (b) From Partial Pressures of Table III.
- (c) Calculated from H_{400}^{\circ}} and F_{400}^{\circ}}.

Thermodynamic functions used in cal:

- (d) Evans and Wagman⁽⁵⁹⁾ : S_{400}^{\circ}} for S(g) = 41.7358, S_{400}^{\circ}} for S₂(g) = 56.85, H_{400}^{\circ}}-H_{0}^{\circ}} for S(g) =
- (e) Guthrie et al⁽¹²⁾ : H_{400}^{\circ}}-H_{0}^{\circ}} for S₂(g) = 2162.5, H_{400}^{\circ}}-H_{0}^{\circ}} for S₈(g) = 2952.
- (f) Interpolated from Luft⁽⁴⁰⁾.

STUDIES OF THE HAEMOPROTEINS


B. W. Baylis

October 30, 1962

INTRODUCTION

Perhaps the most useful introduction to this material, as presented to inorganic chemists, would be an explanation of terms. Therefore, there follow definitions, which, even if not entirely generally accepted, at least are adhered to in this paper.

Phosphine



Derivatives of phosphine found naturally are indicated on Table I on page 54.

Haem group: protoporphyrin complex of iron (II).

Haemochrome: haem with variable donor molecules above and below the plane of the ring.

Haemoglobin: haem with its specific protein attached at one end of the z axis and usually with water at the other. This often refers to that naturally occurring material which has four sub-units of this type.

Haemoprotein: the group of naturally occurring iron(II) or iron(III) complexes with various porphyrins and various donor molecules.

Donor molecule is used here only for molecules or ions along the z axis.

Substituent is used here to refer only to side chains on the porphyrins or model chelating groups.

Methaemoglobin: haemoglobin the iron of which is oxidized to +3.

Attention should be drawn to the seminar given last fall (1) on the metalloporphyrins. None, or at least very little, of the material covered in that seminar is mentioned here.

OXIDATION-REDUCTION POTENTIAL STUDIES

Studying the oxidation potential of iron in haemoproteins has proved quite complicated. These systems display poor reversibilities and worse solubility properties. In addition, the potentials vary with pH, steric and electronic requirements of both substituents on the porphyrin and ligands at the 1:6 sites of the iron, and spin states of the iron(II) and iron(III). Of course these variables are all interconnected. The experiments in this area represent an attempt to sort out each set of variables and examine it alone. When this was not done, the interpretation of the data was subject to considerable error by oversimplification.

October 20, 1952

J. V. Kelly

... to the ... in this paper.

... of ...



... of ...

... of ...

... of ...

... of ...

... of ...

... of ...

... of ...

... of ...

... of ...

... of ...

... of ...


... of ...

... of ...

... of ...

In addition to the experimental difficulties involved, the use of the naturally occurring haemoproteins brings in too many unknown factors for an interpretation to be possible. Hence model systems are devised and studied and the results compared with results of similar studies on the naturally occurring materials. If the two sets of results bear enough resemblance, or if major differences can be rationalized, the interpretation developed from the model system is used to make understandable the data on the natural system.

Martell and Calvin(2) report the following data showing the variation of oxidation potential with pH for several haemochromes.



Lines A,B,F,G, and H indicate that these derivatives change with pH in a way that stabilizes the iron(III). It is doubtful that the second amine remains in the coordination sphere. This hypothesis allows for the following explanation:



An anionic ligand can stabilize a higher oxidation state; thus the hydroxide will stabilize the iron(III). Calculation of the variation of the observed E° with pH based on this equilibrium gives a curve with a slope of 0.60. This is noted to be the slope observed for lines A,B,F,G and H. The iron-cyanide bond is too strong to be affected by the hydroxide ion. Hence, there is no variation in E° with pH for line C. The slopes of lines D and E are explained by a proposed removal of less than one hydrogen ion per iron. This could be accomplished either by removing a hydrogen ion from some of the imidazole rings or by displacing some of the rings by hydroxyl ions.

Changing the substituents on the pyrrole rings changes the potential. R. J. P. Williams (3), using 8-hydroxy quinoline-iron complexes as his models, studied the effects on E° of change in pH while varying the substituents. Comparison of E_p with acid constants of the ligands led to no simple relationship. With the 5-cyano and 5-formyl derivatives anomalously low potentials were obtained, indicating stabilization of the iron(II). These substituents are π acceptors, a fact which is not reflected in the pK_a of the ligand. The increased π acceptance of the ligand will stabilize the iron(II), which is 10-30% diamagnetic.

The oxidation potentials of derivatives of the 1,10-phenanthroline complexes(3) are all about -1 volt, indicating stabilization of the iron(II). The reduced form is diamagnetic, and the oxidized form is a mixture of spin-paired and spin-free complexes. The reagent 2-methyl phenanthroline provides an exception, with an oxidation potential of 0.585 volt. Its iron(II) complex is paramagnetic, indicating that the ligand cannot interact as strongly with the metal as before. This may be caused by steric interference by the methyl group. These studies refute the earlier assumption (2) that the stability of iron(III) complexes, and hence the oxidation potential, is increased by increasing electron-donor properties of the substituents. It was here shown that this is only true if both oxidation states are either spin-free or spin-paired. In any case, iron(II) is always stabilized by electron acceptance. The biologically important complexes contain diamagnetic iron(II) and spin-free or a mixture of spin-free and spin-paired iron(III). Therefore, in these σ donors and π acceptors will stabilize the iron(II).

Williams (4) has recently studied iron-dipyridyl complexes as a model system. Comparing E_p with pK_a values of the substituted ligands indicates that the variation in oxidation potential is dominated by the spin state. This information can be applied to explain the data available on porphyrins. Other pertinent information follows:

- 1) The iron(II)-nitrogen distance is 0.1-0.2 Å. longer than the iron(III)-nitrogen distance. Therefore, steric considerations are of greater importance in spin-paired iron(III) complexes.
- 2) Sigma donor power of the nitrogen atoms in the porphyrins decreases in the order meso or haemato > proto > rhodo > dihydro. These porphyrins are described in the Table on page 54.
- 3) Pi acceptor power of the porphyrins decreases in the order rhodo > proto > meso or haemato > dihydro.

The first part of the report is a summary of the work done during the year. It is followed by a detailed account of the work done during the year, and a list of the publications of the author.


The second part of the report is a detailed account of the work done during the year. It is followed by a list of the publications of the author.

The third part of the report is a detailed account of the work done during the year. It is followed by a list of the publications of the author.

The fourth part of the report is a detailed account of the work done during the year. It is followed by a list of the publications of the author.

The fifth part of the report is a detailed account of the work done during the year. It is followed by a list of the publications of the author.

The sixth part of the report is a detailed account of the work done during the year. It is followed by a list of the publications of the author.




The observed chart⁽⁵⁾ could result from reduced steric restrictions around the iron and a change from imidazole to primary amine ligands as one proceeds from left to right. Properties of the porphyrins prove less important. Considerations of this type have led to the assignments of many of the donor groups listed on the Table on page 54.

Williams (6) gives the order of increasing oxidation potential of iron(II)- dimethyl glyoxime base₂ as pyridine < imidazole < water < ammonia < hydroxide. Falk(7) summarizes the considerations involved in the change in oxidation potential with change in donor molecule as follows:

- 1) Electrostatic forces. If the donor is an anion, the higher valence state will be favored, raising the potential.
- 2) Back-double-bonding. This favors the lower oxidation state, lowering the potential.
- 3) Ligand-field stabilization energy. This varies with the cations and spin states involved.
- 4) The pKa values of the ligands. In a series of related ligands, increasing pKa causes an increase in oxidation potential.

It is worth noting that the curves of E° versus pH and the log of oxygen absorption by haemoglobin are quite similar. From this Martell and Calvin(2) conclude that the processes involved are similarly affected by changes made on the molecule. The oxidation potential of haemoglobin at the pH of blood is -0.1 volt. Thus the effect of the donor groups is to decrease the tendency towards oxidation but not so much as to prevent oxygenation. Oxygen further stabilizes the spin-paired complex.

1. 1911
2. 1912
3. 1913
4. 1914
5. 1915



The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

The following table shows the percentage of the population in each of the five years mentioned above. The percentage of the population in each year is as follows: 1911, 10%; 1912, 20%; 1913, 30%; 1914, 40%; 1915, 50%.

SPECTRAL STUDIES

Spectra have been studied for haemoglobin over the IR, visible, and UV regions (8). Most important are the α and β bands, the Soret or γ band, and the charge-transfer bands. The first three are considered to be $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ transitions by some (9). Platt (10) has carried out molecular-orbital calculations to arrive at expressions for the energy states of the porphine molecule. The lowest energy transitions possible correspond to the α and β bands.


In porphine there are four of the longer wavelength bands, I, II, III, and IV (9,10). Bands III and IV arise from transitions involving electron density moving away from the amine hydrogens, and I and II arise from electron density moving away from the nitrogens that are not bound to a hydrogen. Bands I and III do not involve any change in vibrational state (10). Bands II and IV arise from coupling of the electronic transition with a change in vibrational state. Hence I and III are strengthened by substituents increasing the asymmetry, whereas II and IV do not change. This behavior has also been observed for the α and β bands of the metal porphyrins. Therefore I and III seem related to the α and II and IV to the β bands.

Upon inserting several divalent cations, Williams (9) observed that the wavelengths for α and β decreased with increasing electronegativity of the cation. The greater the ability of the cation to donate π density, the greater the decrease in the wavelengths of α and β .

Haem is paramagnetic and its α band is weak and appears around 575 m μ . After accepting either pyridine or ammonia the complex becomes diamagnetic, α shifts to 558 m μ ; and its intensity increases markedly. Additional ligands change the haemochrome only slightly. Oxygen, carbon monoxide, cyanide, etc., do change the magnetic moment, increase the intensity of α and move α and β to shorter wavelengths. The shift of α and β to lower wavelength is less with neutral bases than with anions (6). In general, α , β , and γ bands do not move far unless there occurs a change in the spin state. The wavelength of α goes up with a change from spin-free to spin-paired in iron(III), but it goes down with iron(II) complexes (5). A high $\Delta\lambda$ ($\lambda_{Fe II} - \lambda_{Fe III}$) indicates a spin free complex in spin-paired cases, the wavelength of α and β are longer the stronger the donor character of the ligand. If the donor molecule is a double-bonding ligand, the more electron-attracting the substituents on the porphyrin, the greater the shift to longer wavelength of the α and β bands (7).

The intensity of the Soret band does not vary unless attack occurs on the porphyrin ring (6). The α and β bands are more intense with low than with high spin complexes. The intensity of α varies with substituents, but β remains constant. The intensity ratio of α to β increases with increasing π donation

ELECTRONIC TRANSITIONS



GROUND STATE

character of the donor molecule. Reasons for the variance of intensity and position of the π and π^* bands as the substituents are changed are given by Gouterman(11).

Many spin-free iron(III) complexes with unsaturated anions have absorbance bands between 600 and 650 m μ and 450 and 500 m μ (6). These are also present in methaemochromes. Because charge transfer involves a change in dipole moment these bands are intense. Low-spin methaemochromes give no charge-transfer bands. The charge-transfer bands move to shorter wavelength as π donation of the donor molecules increase. Unsaturated ligands raise the wavelength more than their π donation ability would indicate. Obviously π interaction is involved here.

In spectra of spin-paired haemoproteins there is a band at 500 m μ . This is also in iron(II) complexes of unsaturated nitrogen bases (9).

OTHER PHYSICAL METHODS

Orgel(12) used electron spin resonance to study hydroxy haemoglobin. His data indicate anisotropy even in the xy plane. This might be caused by approach of the d_{xy} cloud of the imidazole group. This would be in a position to interact with one of the degenerate d_{xz} and d_{yz} orbitals but not the other. He found the differences in energy to be 800 cm.⁻¹ between d_{xy} and d_{xz} and the same splitting between the d_{xz} and d_{yz} .

X-ray methods have been used to study the structure of myoglobin and haemoglobin. The structure of myoglobin has been determined to 2 Å (13). This shows an imidazole residue pointing down the z axis of the haem group. The rest of the protein is wound around, affording a hydrogen bond site for a carboxylic acid group of the porphyrin. In addition, there is a second imidazole residue pointing up the z axis, but it is far enough away to be only mildly affected by the iron.

The haemoglobin study has been completed to a resolution of 5.5 Å. (14). This shows that each of the four units is quite similar to the myoglobin. It also gives an orientation for the four haem groups. These studies indicate that the four units are really similar pairs of 2 identical units.



Orientation of the Haem Groups

Since the protein is attached to a side chain on the porphyrin, asymmetry about each iron is possible. The optical rotary dispersion of catalase(15) indicates that the group of four units exhibits dispersion at 600 m μ but there is no Cotton effect from the single units.

CHEMICAL STUDIES

Hydrolysis studies of these compounds have proved enlightening(6). Complexes with zero or negative charge are difficult to hydrolyze. Thus the pH necessary increases with greater negative charge on the complex. Peroxidase, methaemoglobin, and metmyoglobin hydrolyze at similar pH, indicating similar negative charge. Catalase requires strong base, indicating that the donor molecules are two carboxylates.

Oxygen appears to be absorbed only if there are no active hydrogens on an atom bound to the iron(6). Haemoglobin absorbs oxygen and carbon monoxide reversibly. Peroxidase containing iron(II) absorbs CO but is oxidized by O₂, indicating that there is a hydrogen on one of the atoms bound to the iron. If the iron-nitrogen distance is increased, as by heat, the iron-oxygen bond is also weakened(9). The spin multiplicity of the oxygen is changed by the interaction with haemoglobin(5). All ferro-porphyrin complexes of unsaturated bases can carry oxygen, but no complex of a saturated base has this ability.

The oxygen is labile, even though the complex is covalent. Therefore, the iron is in an energy state only slightly more stable than its high-spin states.

It has been noted that the fourth oxygen to be absorbed shows the greatest affinity(1). Another interesting phenomenon is the Bohr effect. Haemoglobin is a stronger acid than oxyhaemoglobin between pH 6.1 and 4.5, whereas the reverse is true between pH 6.1 and 9.0 (1). This is accounted for (16) by the presence of two independent oxygen-linked acid groups per haem, one of which is strengthened and the other weakened when haem combines with oxygen. The Bohr effect is lessened at high ionic strength. It is thought (17) that haemoglobin dissociates into two molecules at high ionic strength in neutral solution.

Wang(18) has developed a film having the properties similar to haemoglobin, in an attempt to determine the structural requirements for this behavior.

ROLE OF IRON IN REACTIONS

Catalase is known to be involved in oxidation of substrate by peroxides. This behavior has been observed with metmyoglobin, methaemoglobin, and peroxidase(6). All these are iron(III) complexes, and George(19) presents the intermediates as containing iron(IV), with oxygen bound in the sixth coordination position. The other alternative would be for the porphyrin to support the

charge and the iron(III) to be low spin. Both hypotheses are supported (6) by the magnetic moment corresponding to two unpaired electrons, oxidation equivalent of one, charge-transfer bands at 600 m μ and 450 m μ , and Soret band displaced to the red. However, the fact that the Soret band retains its usual intensity indicates that the porphyrin ring has not been attacked. Kinetic studies further support this(20). Electron spin resonance work(21) suggested that the electron was removed from the porphyrin but later work (22) showed the radical concentration to be too low to explain the amount of oxidation. George has argued that the radicals were merely a by-product(23).

Dwyer has contended(24) that too much emphasis is being placed on the iron, which should be thought of as only stabilizing the oxidation product of the organic group. Many reactions involving these complexes appear to be outer-sphere, since no more groups can get close to the iron. Electron exchanges would then appear to occur at the outer fringes of the organic group rather than at the iron. Perhaps the possibility of hydrogen-atom transfer should also be considered as a mechanism.

An example of outer sphere reduction with a haemoprotein as catalyst was suggested by Williams(9).

REFERENCES

1. Howe, W. B., "Properties of the Metalloporphyrins", U. of I. Inorganic Seminars, 1961-62, p. 70.
2. Martell, A. E., and Calvin, M., "Chemistry of the Metal Chelate Compounds", Prentice-Hall, Inc., Englewood Cliffs, N. J., 1952, pp. 359-380.
3. Williams, R. J. P., J. Chem. Soc., 1958, 2010.
4. Williams, R. J. P., Biochemistry, 1, 379 (1962).
5. Williams, R. J. P., "A Symposium on Haematin Enzymes," J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 41-55.
6. Williams, R. J. P., Biochem. J., 78, 246 (1961).
7. Falk, J. E., and Perrin, D. D., "A Symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 56-79.
8. Drabkin, D. L., "A Symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 142-172.
9. Williams, R. J. P., Chem. Revs., 56, 299 (1956).
10. Longuet-Higgins, H. C., Rector, C. W., and Platt, J. R., J. Chem. Phys., 18, 1174 (1950).
11. Gouterman, M., J. Chem. Phys., 30, 1139 (1959).
12. Orgel, L. E., "A Symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 1-18.
13. Kendrew, J. C., Dickerson, R. E., Strandberg, B. E., Hart, R. G., Davies, D. R., Phillips, D. C., and Shore, V. C., Nature, 185, 422 (1960).
14. Perutz, M. F., Rossmann, M. G., Cullis, A. F., Muirhead, H., Will, G., and North, A. C. T., Nature, 185, 416 (1960).

15. Eichhorn, G. L., and Oshahr, A. J., "Advances in the Chemistry of the Coordination Compounds", S. Kirschner, Ed., The Macmillan Company, New York, 1961, pp. 216-223.
16. Antonini, E., Wyman, J., Jr., Rossi-Fanelli, A., and Caputo, A., J. Biol. Chem., 237, 2773(1962).
17. Rossi-Fanelli, A., Antonini, E., and Caputo, A., J. Biol. Chem., 236, 391 (1961).
18. Wang, J. H., "A symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 98-104.
19. George, P., and Irvine, D. H., Biochem. J., 52, 511 (1952).
20. Westheimer, F. H., "Enzymes", Boyer, P. D., Lardy, H., and Myrback, K., Ed., Academic Press, New York (1959), pp. 291-295.
21. Gibson J. F., and Ingram, D. J., Nature, 178, 871(1956).
22. Gibson, J. F., Ingram, D. J., and Nicholls, P., Nature 181, 1398 (1958).
23. George, P. and Irvine, D. H., Biochem. J., 60, 596 (1955).
24. Dwyer, F. P., "A symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 19-28.
25. George, P., Beetlestone, J., and Griffith, J. S., "A Symposium on Haematin Enzymes", J. E. Falk, R. Lemberg, and R. K. Morton, Ed., Pergamon Press, New York, 1961, pp. 105-141.

Table I STRUCTURES OF THE HAEMOPROTEINS

Haemoprotein	Porphyrin	Substituent Placement	Function	Donor Molecules
Haemoglobin	Proto (2) 4 haems per unit (15)	(15) CH ₃ , 1,3,5,8 CH=CH ₂ 2,4 CH ₂ CH ₂ COOH 6,7	(2) O ₂ transport(2)	2 imidazoles, 1 weak (5) or 1 imidazole, 1 H ₂ O
Myoglobin	Proto (2) 1 haem (15)	CHO 2 (2) CH=CH ₂ 4 CH ₂ CH ₂ COOH 7,3	O ₂ storage (2)	1 imidazole, 1 H ₂ O (5)
Chlorocruorin	Spirographis (2) or Rhodo (4)	<u>CHOHCH₂R 2 (7)</u> CH=CHR 4 CHO 8	O ₂ transport(?) (2)	
Catalase	Proto(FeIII) (2) 4 haems (15)	CH=CHR (4) CHOHCH ₂ R (4) 1 partly saturated pyrrole	Catalytic decomp. (2) of H ₂ O ₂	2 carboxylates (5)
Peroxidase	Proto(FeIII) (2) 1 haem		Oxidize substrate(2) with H ₂ O ₂	amine-carboxylate(5) or carboxylate-H ₂ O
Cytochrome a	Rhodo (4)		Cellular Respiration (2)	2 imidazoles, 1 weak(5)
Cytochrome a ₂	Dihydro (4)			
Cytochrome a ₃	Rhodo (4)		Cellular Respiration ⁽²⁾	1 imidazole, 1 H ₂ O
Cytochrome b	Proto (2)		Cellular Respiration(2)	1 amine, 1 imidazole(5)
Cytochrome c	Similar to (4) Meso or Hemato C: 1 haem (15) C ₁ : 4 haems	CH ₃ 1,3,5,8 (7) CH ₂ SHCH ₃ 2,4 CH ₂ CH ₂ COOH 6,7	Cellular Respiration(2)	2 imidazoles (5)
Cytochrome C ₃	Haemato-type(4)			
Cytochrome d	Haemato-type(4)			
Cytochrome f	Haemato-type(4)	No unsaturates (5)		1 amino, 1 H ₂ O (5)

Case No.	Case Name	Case Description	Case Status	Case Date
1001	John Doe	Case 1001: John Doe vs. Jane Smith	Open	2023-10-27
1002	Jane Smith	Case 1002: Jane Smith vs. John Doe	Open	2023-10-27
1003	John Doe	Case 1003: John Doe vs. Jane Smith	Open	2023-10-27
1004	Jane Smith	Case 1004: Jane Smith vs. John Doe	Open	2023-10-27
1005	John Doe	Case 1005: John Doe vs. Jane Smith	Open	2023-10-27
1006	Jane Smith	Case 1006: Jane Smith vs. John Doe	Open	2023-10-27
1007	John Doe	Case 1007: John Doe vs. Jane Smith	Open	2023-10-27
1008	Jane Smith	Case 1008: Jane Smith vs. John Doe	Open	2023-10-27
1009	John Doe	Case 1009: John Doe vs. Jane Smith	Open	2023-10-27
1010	Jane Smith	Case 1010: Jane Smith vs. John Doe	Open	2023-10-27

Table II DATA ON THE HAEMOPROTEINS

Haemoprotein	Spin of Fe(II,III)	E ₀	ν, ε	ν, ε	Soret, ε	ClB, F ₀ (III)
Haemoglobin	high (4) 5.65 BI (6)	-0.15(4)	590,5	550,13 (9)	430(5)	500,10.0(6) 641,4.4(6)
Myoglobin	high (4) 5.73 B M (6)	-0.02(4)	(9) FeIII580(6)	555,15 (9) 520(6)	435(5) 405,170(5)	500,10.2(6) 630,3.9(b)
Chlorocruorin	high (4)	-0.20(4)	605?,12 (9)	565,12 (9)		
Catalase	5.09 B M (6)		585,9.5 (6)	537,13.2 (6)	407,138 (6)	490,17(6) 622,10.5(6)
Peroxidase	(OH ⁻)2.66 (25) (H ₂ O)5.43 (6)		580,2.5(6)	570,7.5 (6)	470(5) 405,91(6)	497,10.0(6) 641,2.8 (6)
Cytochrome a	mostly low(4)	-0.28(4)	605,hi (9)	560-70,10 (9)	444?(5)	
Cytochrome a ₂	high/low(4)	-0.30(4)	605,hi (9)	560-70,10 (9)	448(5)	
Cytochrome a ₃	high/low(4)	-0.30(4)	558,29 (9)	528,16 (9)	425(5)	
Cytochrome b	low (4)	0.00±0.08(4)	550,29 (9)	520,29 (9)	415(5)	
Cytochrome c	mostly low(4)	-0.15±0.25(4)				
Cytochrome c ₂	2.14B M (25)	-0.262 (2)				
Cytochrome c ₃	low (4)	+0.20 (4)				
Cytochrome d	high (4)	0.00 (4)				
Cytochrome f	high/low (4)	-0.36(4)			423(5)	

Note: ϵ is a millimolar extinction coefficient

ANTIFERROMAGNETISM IN SOME FIRST-ROW TRANSITION-METAL FLUORIDES

J. C. Schapiro

November 6, 1962

INTRODUCTION

The chemistry of simple and complex transition-metal fluorides has recently been discussed in two review articles (1,2); the major portion of the material contained in these articles will not be further considered. For the most part, we shall be interested in the behaviour of the first-row transition-metal fluorides in the solid state. Guggenheim(3) has described the preparation of single crystals of some transition-metal fluorides, which has facilitated x-ray and NMR studies.

Fluoride ion is considered to be a weak-field ligand as is evidenced by its position in the spectrochemical series(4), and as such should give rise to high-spin (paramagnetic) complexes. It should be noted that Ferguson, Knox and Wood(5) have found in a study of the absorption spectra of solid K_2NaCrF_6 and CrF_3 (both contain CrF_6 octahedra but, next nearest neighbors differ) that the calculated Dq for these compounds differ by 900 cm^{-1} . Therefore, the environment surrounding the ligand has a pronounced effect on the splitting of the d -orbitals. The field in CrF_3 , which contains Cr-F-Cr bridges, is weaker. It is generally found that fluoro complexes are paramagnetic, the $[NiF_6]^{2-}$ ion is the only known fluoro complex of a first row transition-metal ion which is diamagnetic(6). However, many of these complexes, the magnetic susceptibility curves of which show typical paramagnetic behaviour at room temperature and above, show a decrease in magnetic susceptibility (antiferromagnetic behaviour) as the temperature is decreased.

ANTIFERROMAGNETISM

The unusual properties of an antiferromagnetic compound, below its Néel point, are due to interionic interactions which are comparable to the thermal energies at the Néel point and which become greater than thermal energies with further decrease in temperature. In addition to the maximum point in the magnetic susceptibility curve, the transition from a disordered paramagnetic to an ordered antiferromagnetic structure is also indicated by Δ -type peaks in the specific heat and thermal expansion (7). It is evident, in some materials, that magnetic interactions can not be direct dipole-dipole interactions and that coupling of the dipoles takes place through intermediate ions. Cooperative magnetism of this type is found in many transition-metal fluorides and oxides.

EXCHANGE MECHANISMS

The exchange mechanisms proposed by various investigations have been qualitatively reviewed by Blackman(7). Kanamori(8) has considered the superexchange interaction in terms of the symmetry properties of electron orbitals. Early development of

THE UNIVERSITY OF CHICAGO LIBRARY

CHICAGO, ILL.

1950

The University of Chicago Library is pleased to announce the acquisition of a new volume in the series of the University of Chicago Press. The book, entitled "The History of the University of Chicago" by [Author Name], is a comprehensive study of the institution's development from its founding in 1837 to the present. It covers the early years of struggle, the growth of the faculty and student body, and the university's role in the development of modern higher education. The book is available in paperback for \$12.50 and in hardcover for \$25.00.

This volume is a significant contribution to the history of the University of Chicago. It provides a detailed account of the university's early years, from its founding in 1837 to the present. The author, [Author Name], is a leading expert on the subject and has conducted extensive research into the university's history. The book is written in a clear and engaging style, making it accessible to a wide range of readers. It is a must-read for anyone interested in the history of higher education in the United States. The book is available in paperback for \$12.50 and in hardcover for \$25.00.


The University of Chicago Library is pleased to announce the acquisition of a new volume in the series of the University of Chicago Press. The book, entitled "The History of the University of Chicago" by [Author Name], is a comprehensive study of the institution's development from its founding in 1837 to the present. It covers the early years of struggle, the growth of the faculty and student body, and the university's role in the development of modern higher education. The book is available in paperback for \$12.50 and in hardcover for \$25.00.

The University of Chicago Library is pleased to announce the acquisition of a new volume in the series of the University of Chicago Press. The book, entitled "The History of the University of Chicago" by [Author Name], is a comprehensive study of the institution's development from its founding in 1837 to the present. It covers the early years of struggle, the growth of the faculty and student body, and the university's role in the development of modern higher education. The book is available in paperback for \$12.50 and in hardcover for \$25.00.


exchange mechanisms excluded the possibility of non-orthogonal orbitals. In the crystal, the orbital of the cation and those of the anions overlap considerably; therefore, they are not orthogonal, except when the orbitals have certain different symmetry properties. Yamashita and Kondo(9) have studied the superexchange interaction in MnO using the Heitler-London model and have included overlap (non-orthogonality) among the electron orbitals.

1. SUPEREXCHANGE-ANDERSON MODEL


The first superexchange mechanism was developed by Anderson(10) and was initially applied to the MnO system. Consider a system of two magnetic ions, such as the Mn^{2+} ion separated by an O^{2-} ion, which has the following colinear configuration



where \underline{d}_1 and \underline{d}_2 represent the total (five) unpaired electrons in the $3d$ levels of the Mn^{2+} ions and \underline{p} and \underline{p}^1 represent the two electrons in the bonding $2p$ orbital of the oxygen ion. There is a finite possibility that one electron will jump from the O^{2-} ion into the $3d$ level of an Mn ion, becoming \underline{d}^1 (or alternatively into an s -state, which is coupled by some strong spin-dependent interaction with \underline{d} electrons on the magnetic ion. According to the Pauli principle, the spin of the transferred electron will be antiparallel to the resultant spin of the electrons (\underline{d}_2) already present. A mixture of the original ground state with the following state



will be of lower energy than the original ground state. The electron (\underline{p}) remaining on the oxygen ion can couple with the magnetic electron on the magnetic ion to which the electron has not been transferred. Since the electrons originally on the oxygen ion had antiparallel spins there will appear to be an indirect negative exchange (antiferromagnetic) coupling between the spins of the two Mn ions.



It should be emphasized that the Anderson mechanism is a combination of two processes: (1) electron transfer, and (2) a

coupling process. Therefore, the Anderson model gives rise to antiferromagnetic interaction between cations with more than half-filled d shells and a ferromagnetic interaction between cations with less than half-filled d-shells. In the case of a less than half-filled d-shell, the spin of the transferred electron would be parallel (Hund's rule) to the resultant spin of the electrons on the magnetic ion. The conclusion for the less than half-filled d-shell is, however, not always supported experimentally, e.g., CaMnO_3 is antiferromagnetic.

2. SEMI-COVALENT EXCHANGE-GOODENOUGH MODEL

Goodenough and Loeb(11) and Goodenough (12) have proposed a model for superexchange which takes into account the mixing of 4s and 4p orbitals on the cation. Again, we shall use MnO as an example. In the crystal both lobes of an O^{2-} p orbital may overlap with an empty cation orbital. For coordinate covalence the O^{2-} orbital and empty cation orbital should be nearly degenerate. In ionic crystals, this is not the case for the ground state of the ion. However, if the cation has a net spin, exchange energy is gained if one of the two electrons in the oxygen p orbitals is excited to become part of the cation, thus forming a one electron "semi-covalent" bond. "Semi-covalence is defined as a bond due to the coupling of a single anion electron to the net spin of a cation (parallel spins). The other anion electron, therefore, has spin antiparallel to the spin of that cation. This other electron may participate in semi-covalence with a cation on the other side of the anion, and the spins of the two cations would be coupled antiparallel to each other. If the spins of the cations are parallel, semi-covalent bond formation can not occur.



SEMI-COVALENT EXCHANGE IN COLINEAR M-O-M SYSTEM

3. SLATER MODEL

The Slater model, as described by Yamashita and Kondo(9), takes into account the polarization of the electron cloud of the anion through the exchange energy with the cations. Again, consider the O^{2-} ion between two Mn^{2+} ions in a ^6S state, the spins of which are antiparallel. Since exchange interaction and overlap interaction are effective only between the electrons of the same spin orientation, the charge cloud of one (2p₀) electron of the O^{2-} ion with the positive spin orientation may polarize in such a way as to approach the one Mn^{2+} ion with opposite spin and to go away from the other Mn^{2+} ion with the same spin. In this case the internal energy (promotion of the electron from 2p to 3s level) of the O^{2-} ion increases, but the loss will be compensated by the gain of the interaction energy. As for the other (2p₀) electron with the negative spin orientation, the situation is the same. The other four electrons (2p_x, 1, -1, α , β) may also polarize but to a lesser extent. Here α refers to an electron with $+1/2$ spin and β to an electron with $-1/2$ spin. The configuration corresponding to the Slater model is the following $[\text{Mn}^{2+}, \text{O}^{2-}(2p)^6] + [\text{Mn}^{2+}, \text{O}^{2-}(2p_{\alpha})^2(3s^{\uparrow}) - (2p_{\beta})^3] + [\text{Mn}^{2+}, \text{O}^{2-}, (2p_{\alpha})^3(3s^{\downarrow})(2p_{\beta})^2]$

A fourth mechanism, designated as the double-exchange(7) interaction and pertaining to ferromagnetic behaviour in mixed oxides will not be discussed.

Wollan and co-workers(13) have considered the exchange mechanism in relationship to the structure observed in perovskite lattices (MF_3 and KMnF_3 compounds) The $d_{x^2-y^2}$ and d_{z^2} orbitals are the ones which overlap the p orbitals of the intervening anions in this type of structure, and therefore these are the orbitals which would be expected to be associated with the indirect magnetic exchange in compounds having a perovskite structure. The observed magnetic data(13,14) can be accounted for on the basis of the following types of indirect exchange:

(a) Whenever half-filled orbitals of two magnetic cations overlap, respectively, the two lobes of a given p orbital, the magnetic exchange will be antiferromagnetic.

(b) Whenever empty orbitals of two magnetic cations overlap, respectively, the two lobes of a given anion p orbital, the magnetic exchange is likewise antiferromagnetic.

(c) Whenever an empty orbital of one magnetic ion overlaps one lobe of an anion p orbital and the other lobe of the same anion p orbital overlaps a half-filled orbital of another magnetic ion, the magnetic exchange is ferromagnetic.

TECHNIQUES FOR INVESTIGATION

The two major physical methods which are employed in the experiments to be considered in the following sections are neutron diffraction and paramagnetic resonance (NMR and ESR). Angular variation in the paramagnetic resonance spectra reflects the local symmetry(15). The three types of symmetry to be considered are (1) symmetry of the crystal (2) symmetry of the paramagnetic susceptibility, and (3) local symmetry of the paramagnetic complex. Hyperfine structure may be observed if a ligand Y has a nuclear magnetic moment. If the electron orbits of the X ion overlap those of the Y ions, the order of magnitude of the hyperfine structure is given by the product of the hyperfine structure of the free atom X and the probability of finding the unpaired electron on Y. Therefore, this splitting is observed in the ESR spectra of the magnetic ions and in addition is also observed in the NMR spectra of diamagnetic ligands. Tinkham(16) first applied this technique to fluoride by observing the ESR spectrum of Mn^{2+} in a ZnF_2 crystal.

MAGNETIC BEHAVIOUR OF THE FLUORIDES

1. MF_2 COMPOUNDS

Neutron-diffraction studies(17) of MnF_2 , FeF_2 , CoF_2 , and NiF_2 , at low temperatures showed the super lattice structure characteristic of antiferromagnetism. These compounds crystallize with a rutile lattice, and the magnetic cell was found to have the same dimensions as the chemical unit cell; the moments of the magnetic ions at the lattice corners being antiparallel to the

moments of the body-centered ions. The alignment of the magnetic moments for MnF_2 , FeF_2 , and CoF_2 was found to be along the tetragonal axes.

Shulman and Jaccarino(18,19) investigated the F^{19} resonance in an oriented single crystal of MnF_2 between $68^\circ K.$ and $300^\circ K.$ They attribute the large shift observed from the F^{19} resonance in a diamagnetic field to the promotion of an electron from the F^- ion to the Mn^{2+} ion. It is thus a measure of the superexchange interaction. However, in MnF_2 the F^- ions have C_{2v} site symmetry, and the Mn^{2+} ion has D_{2h} site symmetry. This low symmetry introduces uncertainty into the direction of the orbitals containing unpaired spins and complicates the interpretation of the data in terms of F^- ion orbitals .

2. KMF_3 COMPOUNDS

Okazaki and Suemune(20,21) have examined the crystal structures of $KMnF_3$, $KFeF_3$, $KCoF_3$, $KNiF_3$, and $KCuF_3$ above and below their Néel temperatures(20). All except $KCuF_3$ are of the cubic perovskite type at room temperature.

Compound	(T_N °K)	Crystal Structure Above T_N	Crystal Structure Below T_N
$KMnF_3$	88	Cubic	Monoclinic
$KFeF_3$	113	Cubic	Rhombohedral
$KCoF_3$	114	Cubic	Tetragonal($a > c$)
$KNiF_3$	275	Cubic	Cubic
$KCuF_3$	243	Tetragonal($a > c$)	Tetragonal

Scatturin, Corliss, Elliott and Hastings(14) have studied the antiferromagnetic structures of $KCrF_3$, $KMnF_3$, $KFeF_3$, $KNiF_3$, and $KCuF_3$ via neutron diffraction. The Mn, Fe, Co, and Ni double fluorides exhibit an ordering in which the divalent $3d$ ion is coupled antiferromagnetically to its six nearest neighbors. In the Cr double fluoride, the magnetic structure consists of ferromagnetic (001) sheets, coupled antiferromagnetically along the [001] direction. The authors were not able to observe any magnetic scattering in the neutron-diffraction pattern of $KCuF_3$. The compound $KMnF_3$ is an example of a substance which has been studied by both EPR and NMR techniques. Shulman, Knox, and Wyluda(23) and Shulman and Knox(24) have studied the NMR of F^{19} and measured the hyperfine interactions between the F nucleus and the magnetic electrons. The isotropic hyperfine interaction constant A_s was found to be $(16.26 \pm 0.4) \times 10^{-4} \text{ cm.}^{-1}$ which corresponds to $(0.52 \pm 0.02)\%$ $2s$ character for the unpaired electron, whereas the anisotropic constant $(A_{\parallel} - A_{\perp})$ is equal to $(0.17 \pm 0.1) \times 10^{-4} \text{ cm.}^{-1}$ which corresponds to $(0.18 \pm 0.1)\%$ $2p$ character. An antiferromagnetic transition evidenced by changes in the resonance pattern was observed at $88.5^\circ K.$ Ogawa(25) measured the EPR of $^{55}Mn^{2+}$ ion as a dilute substitutional

1940-1941

...

...

...

...

...
...
...
...
...

...

impurity in single crystals of KMgF_3 (essentially have RMnF_3 environment), KCdF_3 , K_2MgF_4 , and NaMgF_3 . The A_s in KMnF_3 was estimated to be somewhat smaller than $16.7 \times 10^{-4} \text{ cm.}^{-1}$ from the observed results.

Shulman and Knox(26) have also measured the NMR of F^{19} in single crystals of KNiF_3 . Since this crystal has high symmetry (perovskite structure), the resonance shifts observed were interpreted directly in terms of the fraction of unpaired spins in the F^- ion $2s$, $2p_\sigma$, and $2p_\pi$ orbitals. Large amounts of both P_{σ} and P_{π} interactions with the d electrons was indicated.

3. K_2MF_4 COMPOUNDS

The most interesting of these is the Cu^{2+} compound, which has been shown by Knox(27) to have two short Cu-F bonds and four long Cu-F bonds. It is isostructural with K_2NiF_4 . However, whereas the Ni^{2+} compound is strongly antiferromagnetic the Cu^{2+} compound is not(28). This has been shown to be a result of the peculiar structure and the electronic configuration in this compound, which rules out any superexchange interaction.

CONCLUSION

Antiferromagnetic behaviour in the first-row transition-metal fluorides considered is the result of some type of superexchange mechanism. The mechanism of superexchange in transition-metal fluorides is strongly dependent upon the electronic configuration and site symmetry of the metal ions and the crystal symmetry of the compounds.

REFERENCES

1. A. G. Sharpe, *Adv. in Fluorine Chem.*, 1, 29(1960).
2. R. D. Peacock in *Progress in Inorganic Chemistry*, Vol. II, F. A. Cotton, ed., Interscience, New York(1960), p. 193.
3. H. Guggenheim, *J. Phys. Chem.*, 64, 938(1960).
4. T. M. Dunn in *Modern Coordination Chemistry*, J. Lewis and R. G. Wilkins, ed., Interscience, New York (1960), p.266.
5. J. Ferguson, K. Knox, and D. L. Wood, *J. Chem. Phys.*, 35, 2236(1961).
6. W. Klemm, *Angew.Chem.*, 66, 468(1954).
7. L. C. F. Blackman, *Research*, 12, 164 (1959).
8. J. Kanamori, *J. Phys. Chem. Solids*, 10, 87, (1959).
9. J. Yamashita and J. Kondo, *Phys. Rev.*, 109, 730(1958).
10. P. W. Anderson, *ibid.*, 79, 350 (1950).
11. J. B. Goodenough and A. L. Loeb, *ibid.*, 98, 391(1955).
12. J. B. Goodenough, *ibid.*, 100, 565, (1955).
13. E. O. Wollan, H. R. Child, W. C. Koehler, and M. K. Wilkinson, *ibid.*, 112, 1132 (1958).
14. V. Scatturin, L. Corliss, N. Elliott, and J. Hastings, *Acta Cryst.*, 14, 19 (1961).
15. W. Low, *Paramagnetic Resonance in Solids*, Academic Press, New York (1960), pp. 52-123.
16. M. Tinkham, *Proc. Roy. Soc., London*, A236,535(1956).
17. R. A. Erickson, *Phys. Rev.*, 90, 779 (1953).
18. R. G. Shulman and V. Jaccarino, *ibid.*, 103, 1126 (1956).

19. R. G. Shulman and V. Jaccarino, ibid., 103, 1126 (1956).
20. A. Okazaki and Y. Suemune, J. Phys. Soc. Japan, 16, 176(1961).
21. A. Okazaki and Y. Suemune, ibid., 671.
22. K. Hirakawa, K. Hirakawa, and T. Hashimota, ibid., 15, 2063 (1960).
23. R. G. Shulman, K. Knox, and B. J. Wyluda, Bull. Am. Phys. Soc. II, 4, 166 (1959).
24. R. G. Shulman and K. Knox, Phys. Rev., 119, 94 (1960).
25. S. Ogawa, J. Phys. Soc. Japan, 15, 1475 (1960).
26. R. G. Shulman and K. Knox, Phys. Rev. Letters, 4, 603(1960).
27. K. Knox, J. Chem. Phys., 30, 991 (1959).
28. W. Rüdorff and D. Babel, Naturwissenschaften, 49, 230(1962).

This leads to the energy levels

$$E_+ = H_0 + \frac{\alpha}{2} + \left[\left(\frac{\alpha}{2} \right)^2 + 2\beta z \right]^{1/2}$$

$$E_0 = H_0 - \alpha$$

$$E_- = H_0 + \frac{\alpha}{2} - \left[\left(\frac{\alpha}{2} \right)^2 + 2\beta z \right]^{1/2}$$

Only one bonding orbital is available for all values of α/β .
When $\alpha/\beta = 0$, the bonding orbital is

$$\psi_+ = \frac{1}{2} (\phi_A + \sqrt{2} \phi_B + \phi_C)$$

When $\alpha/\beta = 1$,

$$\psi_+ = \left(\frac{1}{\sqrt{3}} \right) (\phi_A + \phi_B + \phi_C)$$

When $\alpha/\beta = 0$, an open type bond results; when $\alpha/\beta = 1$, a central bond forms.

Hamilton(10) has performed a four-center, four-electron calculation for diborane, and Yamazaki(11) has treated the problem as a twelve-electron system. Comparison of these results with that of the simple localized molecular orbital indicates that the three-center bond description is very accurate for diborane (12).

SEMI-TOPOLOGICAL TREATMENT

In an attempt to develop a systematic method of describing the electronic structure of the boron hydrides, Eberhardt, Crawford, and Lipscomb(2) presented a set of valence postulates and equations of balance. By applying semi-topological arguments together with the equations of balance, Dickerson and Lipscomb(13) have been able to limit the possible structures for the hydrides and make predictions as to existence and structure of the hydrides and hydride ions. (Platt (14) attempted a symbolic structure treatment, but his results are incompatible with the results of reference (2) and with the correct structure of B_9H_{15} (7)).

The equations of balance (15) are simple to derive. (See also 16, 3, 2). $[B_r B_p H_{p+q}]^c$ (general boron hydride formula) where r = number of boron atoms with no terminal hydrogen atom attached.

Let s = number of B-H-B hydrogen bridge bonds,
 t = number of three-center B-B-B bonds,
 y = number of two-center B-B bonds,
 x = number of single B-H bonds in excess of one per boron.

Hydrogen atom balance: $s + x = q + c$
 Orbital balance: $s + t = p + r + c$

Electron balance: $t + y = p + \frac{3r}{2} - c - \frac{q}{2}$

Boron hydride structures are tabulated in terms of s , t , y , and x . The following principles are the basis of the topological approach:

1. Assume that each boron atom must be chemically bonded to each adjacent boron atom by at least one two-center or three-center bond.
2. Define the connectivity of a boron atom as the number of other boron atoms to which it is adjacent in such a position as to allow bonding.




Define excess connectivity (EC) as connectivity minus 3.

3. Define the excess connectability of a bond arrangement at a boron site as the number of other boron atoms to which it connects the given one, minus the number of orbitals which it requires from the given boron atom.

Note that a B-B single bond or a B-H-B three-center bond produces no change in the EC of a framework.

The procedure to be followed in looking for feasible boron hydride structures is:

1. Assume a boron framework and make a topological sketch of it, indicating the EC at each boron site.
2. Add extra B-H bonds to every boron atom of $EC \neq -1$, and possibly to certain boron atoms of $EC = 0$ or 1 , but not to a boron atom of $EC = 2$.
3. Add three-center B-B-B bonds or arrangements of them in such a way that the EC at every boron site is reduced to zero.
4. Complete the molecule with B-B and B-H-B bonds, subject to the equations of balance.



Central bond

open bond

central and
hydrogen bridge

adjacent central
bond

Fig. 1. Excess connectabilities of various bonds.

To eliminate from consideration certain structural features not present among the known boron hydrides, these principles are applied in order of decreasing reliability:

- (a) A new hydride would probably have at least a plane of symmetry or some other twofold symmetry element.
- (b) No two adjacent boron atoms can be connected by both a central and an open three-center bond.
- (c) Boron atoms of $EC = 2$ do not contribute to a B-H-B bridge or to an extra B-H, thus restricting B-H-B bridges to the edge of the framework, and BH_2 groups to the outer boron atoms.
- (d) Each boron atom of $EC = -1$ will probably be bonded by one hydrogen atom to at least one of its two adjacent boron atoms.
- (e) An edge boron atom of $EC = 1$ will probably not be connected to two other boron atoms by hydrogen bridges. This is the weakest assumption.

The justification of these five principles is ultimately the fact that they are generalizations from the presently known hydrides.

The importance of this topological extension of the valence theory is that it so dramatically reduces the number of likely structures in a given case. For example, B_4 yields the true B_4H_4 structure uniquely. The composition B_7 yields no satisfactory structures based on an icosahedral fragment, and B_{10} yields only the established $B_{10}H_{14}$. (See (13),(16) for other results and predictions.)

This topological method can be extended to various other situations. It is invariant (15) with respect to replacement of H^- by an electron pair donor ligand L; replacement of H by D, a halogen, an alkyl group, or an aryl group; replacement of a bridge hydrogen by XR_2 to be bonded then by two X-B bonds, where X is N or P and R is H, an alkyl group, or an aryl group; replacement of a three-center bonded H^- by a three-center bonded CH_3^- ; substitution of hetero-atoms in the boron framework, e.g., C for B, Be for B^+ , N for B^{2-} , Al for B. The topological treatment has also been used to study reactions and predict products (17). The boron hydride ions, the boron halides, and $B_xH_yO_z$ compounds have been treated in the same manner(16).

MOLECULAR-ORBITAL DESCRIPTIONS

Although the three-center bond concept is useful, the electrons in the more complex and symmetrical boron hydrides must be much more delocalized. Indeed, the localized bond theory does not and cannot predict accurate charge distributions. The molecular orbital approach is necessary for this.

In some of the simpler boron hydrides, the molecular orbitals can be formed readily by assuming the hybridization of the borons in the framework and combining these hybridized orbitals to fit the symmetry of the molecule(2,9,18). In compounds with low symmetry, it is not so obvious which combination will yield proper symmetry orbitals; so group theoretical treatment must be worked out in detail.

The basic assumptions employed in the molecular-orbital method are those used in the simple three-center bond treatment, but now there are many atoms involved. Also, $S_{jk} = S_{kj}$ is not a valid assumption for the boron polyhedra. All the overlap integrals must be evaluated, because the assumption of only nearest-neighbor interactions is not a reasonable assumption and leads to inconsistencies (19).

Longuet-Higgins (20) and Lipscomb(13) both predicted by molecular-orbital theory that a B_{12} hydride should exist as the $B_{12}H_{12}^{-2}$ ion. Just recently, the existence and structure of this ion has been determined(21).

An LCAO treatment of the boron framework orbitals in $B_{10}H_{14}$ yields a charge distribution which agrees with that expected from chemical behavior (22). An investigation of boron orbitals in the B_5H_5 molecule of D_{3h} symmetry indicates expected stability for $B_5H_5^{-2}$, but not for the neutral molecule (22).

The ion $B_{10}H_{10}^{-2}$ has been prepared. With D_{4d} symmetry postulated, its closed shell structure was predicted by molecular orbital theory (23).

The existence of $B_4H_7^-$ and $B_6H_{11}^+$, both related to B_6H_9 , has been predicted because molecular-orbital treatment suggests these ions should be stable (24).

CONCLUSION

Many new hydride ions which have been predicted to be stable by both molecular-orbital and topological methods have been isolated. Their structures confirm the predictions and support the validity of these treatments (25, 26, 27). A crystal-field model, which has been developed for finding the molecular orbitals of identical atoms, is applicable to boron polyhedra (28). The molecular-orbital and semi-topological approaches are complimentary methods for the study of the bonding and structure of the boron hydrides. These principles can be extended to other electron-deficient systems and to metallic bonding (29).

REFERENCES

1. H. C. Longuet-Higgins: J. chim. phys., 46, 275 (1949).
2. W. H. Eberhardt, B. C. Crawford, Jr., and W. N. Lipscomb: J. Chem. Phys., 22, 989 (1954).
3. W. N. Lipscomb: J. Chem. Phys., 22, 985 (1954).
4. B. Siegel and J. L. Mack: J. Chem. Ed., 34, 314 (1957).

5. F. L. Hirshfeld, K. Eriks, R. E. Dickerson, E. L. Lippert, Jr., and W. N. Lipscomb: J. Chem. Phys., 28, 56(1958).
6. A. Stock, Hydrides of Boron and Silicon, Cornell University Press, New York (1933).
7. R. E. Dickerson, P. J. Wheatley, P. A. Powell, and W. N. Lipscomb: J. Chem. Phys., 27, 200 (1957).
8. R. Grimes, F. E. Wang, R. Lewin, and W. N. Lipscomb: Proc. Natl. Acad. Sci., U.S.A., 47, 996 (1961).
9. H. C. Longuet-Higgins: Quart. Revs., 11, 121(1957).
10. W. C. Hamilton: Proc. Roy. Soc. (London), A235, 395 (1956).
11. M. Yamazaki: J. Chem. Phys., 27, 1401 (1957).
12. W. C. Hamilton: J. Chem. Phys., 29, 460 (1958).
13. R. E. Dickerson and W. N. Lipscomb: J. Chem. Phys., 27, 212 (1957).
14. J. R. Platt: J. Chem. Phys., 22, 1033 (1954).
15. W. N. Lipscomb: Proc. Natl. Acad. Sci., U.S.A., 47, 1791 (1961).
16. W. N. Lipscomb: "Advances in Inorganic Chemistry and Radiochemistry", Vol. I, p. 117 (1959).
17. W. N. Lipscomb: J. Inorg. Nuclear Chem., 11, 1 (1959).
18. H. C. Longuet-Higgins: J. Roy. Inst. Chem., 77, 179 (1953).
19. R. Hoffmann and W. N. Lipscomb: J. Chem. Phys., 36, 2179 (1962).
20. H. C. Longuet-Higgins and M. de V. Roberts: Proc. Roy. Soc. (London), A230, 110 (1955).
21. J. A. Wunderlich and W. N. Lipscomb: J. Am. Chem. Soc., 82, 4427 (1960).
22. E. B. Moore, Jr., L. L. Lohr, Jr., and W. N. Lipscomb: J. Chem. Phys., 35, 1329 (1961).
23. W. N. Lipscomb, A. R. Pitochelli, and M. F. Hawthorne: J. Am. Chem. Soc., 81, 5833 (1959).
24. W. N. Lipscomb: J. Chem. Phys., 28, 170 (1958).
25. J. Reddy and W. N. Lipscomb: J. Chem. Phys., 31, 610 (1959).
26. C. R. Peters and C. E. Nordman: J. Am. Chem. Soc., 82, 5758 (1960).
27. A. Kaczmarczyk, R. D. Dobrott, and W. N. Lipscomb: Proc. Natl. Acad. Sci., U.S.A., 48, 729 (1962).
28. R. Hoffmann and M. Gouterman: J. Chem. Phys., 36, 2189 (1962).
29. D. W. Dickerhoof, U. of Illinois Inorganic Seminars, 1958-59, pg. 53.

NUCLEAR QUADRUPOLE RESONANCE IN INORGANIC CHEMISTRY

Ronald H. Carlson

November 20, 1962

INTRODUCTION.

Nuclear quadrupole interactions in gaseous molecules were first discovered in an examination of the radiofrequency spectra of HD and D₂ using molecular beam spectroscopic methods (1). The first evidence for quadrupole interactions in the microwave spectra of gaseous molecules was obtained by Coles and Good (2) in their study of NH₃. Dehmelt and Krüger (3) in their investigation of trans-dichloroethylene were the first to observe direct radiofrequency "pure" quadrupole spectra of solids.

Since its early beginning in the past decade, nuclear quadrupole resonance (NQR) spectroscopy has proven itself an extremely useful tool for gaining additional insight into the chemical bonding of many inorganic compounds. Inasmuch as most of the current NQR literature deals with investigations of solid compounds, this seminar will be concerned primarily with data obtained from pure quadrupole resonance spectroscopy. However, some mention will be made of earlier data obtained from the microwave spectra of gaseous molecules. Several excellent reviews dealing with nuclear quadrupole resonance in both the solid and gaseous state are available (4,5,6,7,8,9,10).

THEORY

Nuclei, the spins of which are greater than 1/2, have a non-spherical charge distribution with respect to the spin axis. The magnitude of this departure of nuclear charge from spherical symmetry is measured by the nuclear quadrupole moment Q . The usual definition of an electric quadrupole moment implies units of charge x distance squared. However, in defining the nuclear quadrupole moment it has been the convention to factor out the protonic charge and to express Q simply in terms cm² (6). Some of the more common nuclei possessing quadrupole moments are (12): H²(D), Li⁶, Li⁷, B¹⁰, B¹¹, N¹⁴, O¹⁷, Al²⁷, S³³, Cl³⁵, Cl³⁷, Br⁷⁹, Br⁸¹ and I¹²⁷.

When placed in an inhomogeneous molecular electric field, a quadrupolar nucleus is capable of undergoing a reorientation under the influence of an oscillating magnetic field of suitable radiofrequency. This reorientation phenomenon is called "nuclear quadrupole resonance". Each possible orientation of a nucleus with respect to the inhomogeneous molecular electric field is quantized, i.e. has a discrete characteristic potential energy. The energy of reorientation or interaction is directly proportional to the "nuclear quadrupole coupling constant" eQq , where e is the protonic charge, Q the quadrupole moment, and q the average value of the electric field gradient. The quadrupole coupling constant has dimensions of energy, but by convention is divided by Planck's constant h to give it dimensions of frequency (usually megacycles per sec.).

Nuclear quadrupole resonances can be measured both in the solid state and in the gaseous state. In the former, they are measured directly in the long wave-length radiofrequency region, whereas in the latter they are measured indirectly as perturbations of rotational lines in the microwave region. From a theoretical standpoint at least, it is much easier to observe quadrupole resonances directly from radiofrequency spectra than indirectly from microwave spectra.

The field gradient q comprises a symmetric tensor having 9 components q_{ij} in the Cartesian coordinate system, where $q_{ij} = \frac{eQ}{4\pi\epsilon_0} \frac{\partial^2 V}{\partial x_i \partial x_j}$ with i, j representing the X, Y or Z Cartesian axis (4).

V is the time-averaged electrostatic potential observed at the nucleus as arising from all charges residing outside of the nucleus. From symmetry considerations and from Laplace's equation: $q_{XX} + q_{YY} + q_{ZZ} = 0$, it is possible to reduce the number of q_{ij} components to 5. Further simplification of the system may be obtained by choosing a special set of axes in which all cross terms (q_{XZ} , q_{YZ} , etc.) become zero. This particular axis system is known as the "principal axis system", and in it the field gradient components are designated by small letter subscripts q_{xx} , q_{yy} , q_{zz} (6). The principal axis system is chosen by convention such that not only the cross terms become zero, but also $|q_{xx}| \leq |q_{yy}| \leq |q_{zz}|$ (I). One can go a step further in the simplification process and define a quantity η called the asymmetry parameter such that $\eta = \frac{q_{xx} - q_{yy}}{q_{zz}}$ (II). η can take on

any value between 0 and 1. Where $\eta = 0$ there is axial symmetry around the z axis and $q_{xx} = q_{yy} = -1/2 q_{zz}$. When $\eta \neq 0$, there is asymmetry around the z axis and the system becomes more complicated.

The quadrupole Hamiltonian may be expressed as (6)

$$\underline{H} = \frac{eQ}{4\pi\epsilon_0} \frac{1}{2I(2I-1)} (q_{xx} \underline{I}_x^2 + q_{yy} \underline{I}_y^2 + q_{zz} \underline{I}_z^2) \quad \text{(III)}$$

where \underline{I}_x , \underline{I}_y , and \underline{I}_z are the projections of the nuclear spin angular momentum upon the 3 principal axes. By constructing the Hamiltonian matrix elements and diagonalizing the resulting matrix, the eigenvalues of the Hamiltonian (the energy levels of the quadrupolar system) for axially symmetric fields ($\eta = 0$) can be obtained (13):

$$\underline{E}_m = \underline{A} [3m^2 - I(I+1)] \quad \text{(IV)}$$

Where I is the nuclear spin quantum number and m is the magnetic quantum number representing the projection of I upon the z axis. The term m can take on values ranging from $I, I-1, \dots, -I$. The quantity \underline{A} in the above expression is equal to $\frac{eQq}{4\pi\epsilon_0}$.

Since the two states $\pm m$ have the same energy, the above energy levels will be doubly degenerate.

The frequencies for quadrupolar transitions are given by the familiar relation $\Delta E = h\nu$ and the selection rule $\Delta m = \pm 1$. For spins of 1 and 3/2 there is only 1 transition frequency, for spins of 2 and 5/2 there are 2 transition frequencies, for spins of 3 and 7/2 there are 3 transition frequencies, etc. For nuclei such as Br^{79} , Br^{81} , Cl^{35} , Cl^{37} with $I = 3/2$, it can readily be shown that

$$\underline{E} \pm 1/2 = -1/4 \underline{eQq} \text{ and } \underline{E} \pm 3/2 = 1/4 \underline{eQq}$$

and that $\underline{\nu} = \Delta E/h = 1/2 \underline{eQq}/h$ (V).

For N^{14} with $I = 1$, $\underline{\nu} = 3/4 \underline{eQq}/h$ (VI). However, I^{127} with $I = 5/2$ has two resonance frequencies with


$$\begin{aligned} \underline{\nu}_1 &= 3/20 \underline{eQq}/h \\ \underline{\nu}_2 &= 6/20 \underline{eQq}/h \end{aligned} \quad (\text{VII})$$

Measurement of the transition frequencies thus allows one to calculate the coupling constants directly in an axially symmetric field.

For nonaxially symmetric fields, the additional parameter η appears in the energy level expressions. For $I = 3/2$ ($\eta \neq 0$), measurement of the transition frequency alone is not sufficient to establish either \underline{eQq} or η . However, by applying a magnetic field and observing the Zeeman splittings, these quantities may be readily determined. For $I = 1, 2, 5/2$ or greater, \underline{eQq} and η may be determined directly from transition frequencies without having to resort to application of a magnetic field (6).

CORRELATION OF \underline{eQq} WITH MOLECULAR STRUCTURE

As previously mentioned, the electric field gradient q in the principal axis system is composed of the three components $q_{xx} = q_{yy} = -1/2 q_{zz} = -1/2 \partial^2 V / \partial z^2$. Now since $\underline{V} = -e/r$, $\frac{\partial^2 V}{\partial z^2} = -2e/r^3$ and $q_{zz} = e/r^3$. Thus q is directly proportional to $1/r^3$ for a charge e at a distance r from the quadrupolar nucleus. This can be illustrated by Figure I.



By convention, the principal axis coordinate system is so chosen that the z axis coincides with the line of centers between the two charges along \underline{r} . When discussing molecules, one is interested in the case where the valence electron ($\ominus e$) participates in a chemical bond with another atom. The z axis then often lies in the direction of the valence bond joining the two atoms, and hence q_{zz} is the component most sensitive to field gradient fluctuations \mathcal{E}^2 brought about by variations in chemical bonding. The quantity q_{zz} has been shown to be inversely proportional to $\frac{1}{r^3}$; thus it is to be expected that only those

electrons which penetrate sufficiently close to the nucleus will have an effect on q_{zz} . Townes and Dailey (14) have shown that the magnitude of q_{zz} at the nucleus is almost entirely dependent upon the electronic distribution in the valence p electronic shell. They state that any s orbital or any closed valence electronic shell is spherically symmetrical and hence gives no contribution to q_{zz} and that valence d and f orbitals do not penetrate sufficiently close to the nucleus to have an effect on q_{zz} .

The contribution to the electric field gradient at the nucleus may arise from the following sources (9):

- 1) Valence electrons of the nucleus in question
- 2) Distortion of the closed shells of electrons around the nucleus
- 3) Charge distributions associated with adjacent atoms or ions.

Generally, to evaluate q at a particular nucleus in a molecule it is necessary only to consider contributions of type 1. However, when dealing with highly ionic molecules, contributions 2 and 3 become important and should be taken into account.

The first formal attempt to interpret observed nuclear quadrupole coupling constants and to correlate them with molecular electronic structure was made by Townes and Dailey in their discussion of some simple Cl compounds (14). They argued that since a neutral Cl atom has 5 valence p electrons and thus lacks one electron of acquiring a symmetrical closed-shell configuration, it would be expected to have a rather large quadrupole coupling constant. Townes and Dailey further state that for a Cl atom involved in a covalent bond, it would be a good approximation to say that it still has a defect of one valence p electron and hence should have a quadrupole coupling constant nearly as large as that for an isolated Cl atom. On the other hand, a Cl atom involved in a chemical bond with a highly electropositive atom such as Na or K has acquired a good share in another p electron and hence has a very nearly symmetrical closed valence shell. Under these circumstances, Cl would be expected to have an eQq value of zero or very close to zero. This effect can best be

represented by the following table:

Table I

<u>Molecule</u>	<u>Structure</u>	<u>eQq (Mc/sec)</u>
atomic Cl		-110.4
ClCN	Cl-C≡N	-82.5
CH ₃ Cl	H ₃ C-Cl	-75.13
NaCl	Na ⁺ Cl ⁻	<1

From the very low value of eQq for NaCl, it was concluded that its bond was almost completely ionic in character. The table also shows that the eQq values for covalently bonded Cl are approximately 25% lower than the corresponding value for atomic Cl. Townes and Dailey attribute this, not to fractional ionic character, but to existing s-p hybridization of the covalent bond of an amount approximating 18%.

In a later paper (15), Townes and Dailey revised their estimates of s-p hybridization in halogens to include Br and I as well as Cl. They put forth a simple rule for the estimation of the amount of s character in a hybrid halide bond: Cl, Br and I bonds are taken to have 15% s character whenever the halogen is bonded to an atom which is more electropositive than the halogen by 0.25 of a unit, otherwise no hybridization is allowed. The authors defend this rule on the basis of observed quadrupole coupling data for halogens and on the basis of promotional energy considerations. Furthermore, they limit the amount of existing d character in halide bonds to 5% or less and in most cases consider it negligible.

Townes and Dailey (14) formulated the equations

$$\underline{eQq}_{(mol)} = [(1-\underline{s}+\underline{d})(1-\underline{i})]\underline{eQq}_{(at)} \quad \text{(VIII)}$$

$$\underline{eQq}_{(mol)} = [(1-\underline{s}+\underline{d})(1-\underline{i})+2(1+\underline{\epsilon})]\underline{eQq}_{(at)} \quad \text{(IX)}$$

for the quadrupole coupling of a halogen forming a hybrid bond with negative and positive ionic character i respectively; s and d are the amount of s and d character in the bond and ε is a constant (9), usually approximated to be 0.25, added to account for the fact that positive ionization increases q by pulling all of the electrons closer to the nucleus. The quantity eQq_(at) is the coupling constant of an isolated atom and has been accurately determined for Cl, Br, and I by atomic beam measurements (9). Using molecule quadrupole coupling constants obtained from microwave spectra and employing equations (VIII) and (IX) and their hybridization rule, Townes and Dailey made a plot of ionicity vs. electronegativity difference for a series of diatomic halides. The curve revealed that for electronegativity differences greater than 2, molecules are essentially 100% ionic.

Gordy (16,17,18) has published a series of papers in opposition to Townes and Dailey's hybridization rule for halogens. He argues, on the basis of dipole moment considerations and observed coupling constants, that pure p orbitals are involved in Br, Cl and I bonds to all atoms except the two first row elements of C and F. Gordy defines a quantity


$$P_q = \left| \frac{eQq_{mol}}{eQq_{at}} \right| \quad (X)$$

where $1 - P_q$ is assumed to be equal to the ionicity of a molecule. He states that quadrupole measurements can be used to evaluate effective electronegativities of atoms bonded in various chemical groups since ionic character is a very sensitive function of electronegativity difference between atoms. For a series of diatomic halides and group (IV) polyatomic halides, Gordy made a plot of $1 - P_q$ against electronegativity difference. From this curve, Gordy concluded that to a good approximation

$$1 - P_q = \frac{X_{hal} - X_m}{2} \quad \text{for } |X_{hal} - X_m| < 2 \quad (XI)$$

where X_{hal} is the electronegativity of the halogen in question and X_m is the effective electronegativity of the atom bonded to the halogen. Gordy also concluded, as did Townes and Dailey, that for electronegativity differences greater than 2, bonding is essentially 100% ionic. Figure II gives a comparison between the Townes and Dailey, Gordy, and Pauling ionicity vs. electronegativity curves. Pauling derived his curve from dipole moment calculations made on hydrohalides (19).

Figure II



As can be seen, both the Townes-Dailey and Gordy curves differ quite markedly from Pauling's curve. This divergence can be explained by the fact that in his dipole moment calculations Pauling did not correct for the large overlap moment, which in the hydrohalides opposes the primary moment, and hence obtained low values of ionicity (18). The divergence between the Townes-Dailey and Gordy curves at lower electronegativity differences can be attributed to the different hybridization rules employed by the two groups of workers.

It is not possible on the basis of quadrupole measurements alone to draw conclusions as to whether Gordy's or Townes and Dailey's hybridization rule is the more nearly correct. Some experimental justification for the latter rule, however, has been obtained by Dailey (20) in his study of the halogen bond character of the alkyl halides. Dailey, using a modified form (4) of equation (VIII)

$$\underline{eQq}_{mol} = (1 - \underline{s} + \underline{d} - \underline{I} - \underline{II}) \underline{eQq}_{at} \quad (XII)$$

and assuming negligible double bond character II and making an independent calculation of the ionicity \underline{I} , was able to determine the sum total of \underline{s} and \underline{d} hybridization for a series of 24 alkyl halides from observed coupling constants. The average values of $(-\underline{s} + \underline{d})$ calculated by Dailey were 13.6% for C-Cl bonds, 8.6% for C-Br bonds, and 1.8% for C-I bonds, in good support of the hybridization rule.

When using quadrupole coupling constants as the basis for correlating the molecular structure of a given family of compounds, it is best to define a quantity,

$$\underline{U}_p = \frac{\underline{eQq}_{(mol)}}{\underline{eQq}_{(at)}} \quad (XIII)$$

called the "number of unbalanced \underline{p} electrons directed along the bond axis" (9). With this quantity, one is able to compare coupling constants obtained from different quadrupolar nuclei. It should be observed that \underline{U}_p has the same meaning as \underline{P}_q defined earlier by Gordy (18). From equation (XII) one can see that

$$\underline{U}_p = \frac{\underline{eQq}_{mol}}{\underline{eQq}_{at}} = (1 - \underline{s} + \underline{d} - \underline{I} - \underline{II}) \quad (XIV)$$

and hence that there are 4 unknown parameters influencing \underline{U}_p . The equation shows that \underline{s} hybridization, ionicity, and double bond character all tend to decrease \underline{U}_p , whereas \underline{d} hybridization tends to increase \underline{U}_p . It is quite apparent therefore that one single measurement of a molecular quadrupole coupling constant will not suffice to establish all 4 of the molecular parameters. Considerable insight can be gained, however, by studying related families of compounds where some of the parameters remain fairly constant or are zero, and by utilizing either Townes and Dailey's or Gordy's hybridization rule and ionicity curve.

Whitehead and Jaffé (46), in their investigation of the halogens and the interhalogens, have attempted to reevaluate the relative contributions of orbital hybridization and ionic character to U_p . They state that neither the assumption of 0% s hybridization as proposed by Gordy, nor the assumption of a set 15% s hybridization as proposed by Townes and Dailey is reasonable or satisfactory when considering halogen bonding orbitals. The authors argue that orbital hybridization will occur only to the extent to match the energies of the bonding orbitals.

APPLICATION OF NQR TO GROUP IV HALIDES

Livingston (21, 22) has obtained the pure quadrupole resonance spectra of a series of chloro-substituted methanes. A regular decrease in eQq was observed in going from CCl_4 to CH_3Cl . Livingston explained this phenomenon by the fact that in going from CCl_4 to CH_3Cl , Cl is being replaced by a less electro-negative element H, with the result that the effective electro-negativity of C is decreased and hence the ionicity of the C-Cl bond is increased. This same trend in coupling constants was also observed by Robinson and coworkers (23) in their investigation of the pure quadrupole resonance spectra of bromo and iodo-substituted methanes. They too ascribed the decrease in eQq as originating from changes in the effective electronegativities of the bonding atoms as H is substituted for Br or I.

Mays and Dailey (24), using microwave spectroscopy, obtained the quadrupole coupling constants of a series of group IV halides of general formula MH_3X , where $M=C, Si$ and Ge and $X=Br$ and Cl . They interpreted the eQq values in terms of the % ionic character and % single bond and double bond character of the M-X bond. Their calculations were based on the following assumptions:

1. Silane and germane derivatives have a double bond structure of the type $H_3M=X$ with metal d orbitals being employed in the bonding.

2. The carbon derivatives resonate only between the single bond covalent structure and the ionic structure i.e.
 $H_3C-X \longleftrightarrow H_3C^+X^-$.

3. In the double bond structure, $H_3M^-=X^+$, Cl and Br contribute a hybrid orbital with negligible d character and with 18% and 8% s character respectively to the σ component of the double bond. A pure p orbital is contributed to the π component of the double bond.

4. The % double bond character can be calculated using the equation: $y = (r_1 - r_0) / (r_1 + 2r_0 - 3r_2)$, where y is the fractional double bond character, r_1 and r_2 are actual M-X single and double distances respectively, and r_0 is the observed internuclear distance of the molecule. Mays and Dailey's results are summarized in Table II. The authors assume an experimental error of 5% in their data.

Table II

Molecule	% double bond character	% single bond character	% ionic character
CH ₃ Cl	0	82	18
SiH ₃ Cl	29	34	37
GeH ₃ Cl	15	44	41
CH ₃ Br	0	82	18
SiH ₃ Br	25	34	41
GeH ₃ Br	15	46	39

In the light of these results, Mays and Dailey (24) were able to rationalize the fact that the dipole moment of GeH₃Cl is considerably greater than that of SiH₃Cl (2.12 > 1.30) despite the fact that the electronegativity difference between Ge and Si is very small. They state that increased π character in Si compounds tends to shorten the Si-X internuclear distance and hence decrease the dipole moment.


Schawlow (25) has interpreted the halogen quadrupole coupling constants obtained by him and others for a series of solid group IV tetrahalides. He used Townes and Dailey's hybridization rule and the same basic method of calculation of bond parameters as did Mays and Dailey (24). Schawlow's results showed increasing ionic character as the electronegativity difference between the group IV metal and the halogen increased. The agreement with Townes and Dailey's ionicity curve, however, was not very good (4), with only the data points obtained from GeI₄, SnI₄, CCl₄, and CBr₄ lying close to the curve. All other points were observed to lie somewhat below the curve, indicative of the fact that probably an excessive amount of double bonding was calculated for these compounds. Schawlow observed that the "number of unbalanced p electrons" U_p was less for the Si-halides than for any of the other M-halides. He interpreted this, not as an ionic effect with Si being the most electropositive group IV element, but rather, indicative of the fact that Si-X bonds have more π character than any of the other M-X bonds. Gordy (18), assuming negligible s and d character in halide bonding orbitals, re-evaluated the molecular structure of the group IV tetrahalides from pure quadrupole spectra obtained by several groups of workers (21,22,23,25,26). His results, summarized in Table III, show the same trend in ionic character as noted above for Schawlow's data, but in general show a significantly smaller % double bond character.

Table III

Molecule	U_p observed	X_m	π character	Ionic Character	Single Bond Character
CCl ₄	0.75	2.58	9	12	79
CBr ₄	0.83	2.55	8	5	87
CI ₄	0.93	2.50	4	1	95
SiCl ₄	0.37	2.04	30	18	52
SiBr ₄	0.46	1.98	26	15	59
SiI ₄	0.58	1.96	20	12	68
GeCl ₄	0.47	2.08	14	32	54
GeBr ₄	0.54	2.03	12	27	61
GeI ₄	0.65	2.00	9	21	70
SnCl ₄	0.44	1.97	9	42	49
SnBr ₄	0.50	1.91	8	37	55
SnI ₄	0.61	1.87	6	31	63

Gordy's calculations of the effective electronegativity, X_m , of the central metal ion are particularly interesting, in that they show that Si is less electronegative than Ge, contrary to popular belief. It should be kept in mind that Gordy's electronegativity values are probably subject to some error since the error limits involved in factoring quadrupole coupling constants into components representative of the amount of π character, ionic character and orbital hybridization in a bond are quite large.

Allred and Rochow (27) interpreted some observed quadrupole coupling constants of group IV halides in terms of the electronegativity of the central metal group IV atom. They studied five series of group IV halides: MCl_4 , MH_3Cl , MBr_4 , MH_3Br , and MI_4 where M is C, Si, Ge, and Sn. For each series they obtained a plot of eQq vs. electronegativity difference of the following type:



From this evidence they concluded, that since Si-X compounds show the smallest halogen coupling constants, Si should be most electropositive of the 4 elements under investigation. Thus the following electronegativity order was postulated by Allred and Rochow as accounting for the observed linear relation between eQq and $[X_x - X_m]$: $C > Ge > Sn > Si$.

Drago (28) argues against Allred and Rochow's order of group IV electronegativities, stating that the quadrupole coupling constants of Si halides are lower than those observed for the corresponding Ge and Sn halides possibly due to the fact that additional π character and s hybridization exist in the Si-X bond. Drago states that the amount of s character in the M-X bond should decrease as one goes from C to Sn and hence that the quadrupole coupling constants of Si compounds should be lower than those for the corresponding Ge and Sn compounds as observed.

APPLICATION OF NQR TO OTHER HALIDE SYSTEMS

Nakamura and coworkers (29, 30, 31, 32) have obtained the pure quadrupole resonance spectra of a series of hexahalo platيناتes, palladates, tellurates, and stannates and have interpreted the data in terms of the covalency of the metal ligand bonds. The authors used equation VIII in their calculations and assumed 15% s character and negligible d character in the halide bonding orbitals. In addition, contributions to eQq from π character were assumed negligible. In each complex, the net charge e on the central metal atom was calculated from

the equation

$$\underline{e} = 4-6(1-\underline{i}) \quad (\text{XV})$$

where the quantity $(1-\underline{i})$ represents the covalency of the molecule. The following table summarizes the results obtained by Nakamura and coworkers:

Table IV

<u>Compound</u>	<u>% covalent character (1-i)</u>	<u>net charge (e)</u>
K ₂ PtCl ₆	56	0.64
K ₂ PtBr ₆	62	0.28
K ₂ PtI ₆	70	-0.20
K ₂ PdCl ₆	57	0.58
K ₂ PdBr ₆	63	0.22
(NH ₄) ₂ TeCl ₆	32	2.08
M ₂ TeBr ₆	42	1.48
M ₂ TeI ₆	52	0.88
K ₂ SnBr ₆	40	1.60
(NH ₄) ₂ SnBr ₆	39	1.66

The coupling constants of the hexahalo-tellurates were found to lie very close to one another regardless of the cation ($M = \text{NH}_4^+$, K^+ , Cs^+). Thus eQq was assumed to be governed by chemical effects rather than by crystal effects due to neighboring ions. The above equation was used in the calculation of \underline{e} because each of the central metal ions in the hexahalides is formally tetrapositive but is surrounded by 6 halogen ions from which charges migrate in along coordinate bonds in accordance with the covalent character of the molecule. As can be noted from the table, the covalency of the stannates and tellurates is less than that of the platinates and palladates by as much as 20%. The authors explained this difference in behavior in terms of the electronic configurations of Te and Sn and the fact that Sn is somewhat less electronegative than Pt and Pd.

Hamlen and Koski (33) obtained the pure quadrupole resonance spectra of TiCl_4 and WCl_6 and interpreted their data using Townes and Dailey's procedure. Values of 87% and 77% were calculated for the ionicity of TiCl_4 and WCl_6 respectively. These values when plotted against electronegativity difference fell appreciably off Townes and Dailey's curve. This deviation was explained in terms of existing $\underline{\pi}$ character in the M-Cl bonds.

Reddoch (34) has investigated the compounds TiCl_4 , ThCl_4 , NbCl_5 and TaCl_5 . He observed four resonance lines in the quadrupole spectra of TiCl_4 and hence postulated it to have a crystal structure similar to that of SiCl_4 , GeCl_4 and SnCl_4 , in the spectra of which four resonance lines were also observed (25). Reddock calculated a value of 87% ionicity for ThCl_4 using Townes and Dailey's procedure. He admitted this value might possibly be high, however, due to admixture of some $\underline{\pi}$ character in the bond. The spectra of NbCl_5 and TaCl_5 were discussed in terms of possible crystal structures for the compounds.

The covalent character of the paramagnetic halides $TiCl_3$, VCl_3 , $CrCl_3$, and $CrBr_3$ was obtained by Barnes and Segel (35) from observed pure quadrupole coupling constants. They state that their values of covalency are in reasonable agreement with those based upon electronegativities of the bonding atoms.

SOLID STATE SHIFTS

The quadrupole coupling constant, observed for a molecule in the solid state is usually 10% to 15% lower than the corresponding coupling constant observed for the molecule in the gaseous state (6). Some examples of this effect are noted in the following table.

Molecule	Nucleus	eQq (Mc/sec)		q_{solid}/q_{gas}
		Gas	Solid	
CH_3Cl	Cl^{35}	78.74	68.06	0.910
CH_2Cl_2	Cl^{35}	78.4	71.98	0.918
CH_3Br	Br^{81}	482	442	0.917
CD_3Br	Br^{81}	479.8	441	0.919
CH_3I	I^{127}	1934	1767	0.914
CD_3I	I^{127}	1929	1764	0.914

No absolute explanation of this solid state shift has as yet been proposed. O'Konski offers a general explanation for the shift observed in the alkyl halides in terms of an increase in ionic character in the crystal lattice resulting from an inductive charge transfer along the C-X bond produced by interaction between the molecular dipoles in the lattice. These solid state shifts are a function of the interaction of a molecule with its surrounding neighbors and hence are of considerable value when a knowledge of intermolecular forces, crystal structure, phase transitions and molecular motions in the solid state is desired.

N^{14} RESONANCES

The first N^{14} resonances obtained from pure quadrupole resonance spectroscopy were reported for $BrCN$ and ICN by Watkins and Pound (36). Casabella and Bray (37) completed the investigation of the series when they reported the N^{14} resonance in $ClCN$. Townes and Dailey (38) postulated an intermolecular structure for solid ICN , in addition to the two resonance structures already known, to account for the fact that the coupling constant of I^{127} increases in going from the gaseous to the solid state. Casabella and Bray (37) extended the Townes and Dailey treatment to include $ClCN$ and $BrCN$ and using both N^{14} and halogen coupling constants were able to calculate the % importance of each resonance structure for the cyanohalogenes in both the solid and gaseous state.

Negita and coworkers (39) have observed an N^{14} resonance in solid HCN. At liquid N_2 temperatures, two resonance lines appeared in the spectra, indicative of an asymmetry in the crystal. From their experiments the authors were able to determine the asymmetry parameter η quite easily without having to resort to the application of a magnetic field.

The pure quadrupole resonance spectra of NH_3 and ND_3 were obtained by O'Konski and Flautt (40). They state that the large decrease in the N^{14} coupling constant observed upon condensing gaseous NH_3 and ND_3 , reflects increased ionic character in the solid state. They explained this enhanced ionicity in terms of resonance structures of the type



involving H-bond formation, as being more important in the solid state.

MISCELLANEOUS RESONANCES

The quadrupole resonance spectrum of Al^{27} in the spinel $MgAl_2O_4$ showed that Al^{+3} ion occupies octahedral sites with rhombic point symmetry D_{3d} (41). Jones and Uehling (42) have obtained the pure quadrupole resonance spectrum of As^{75} in $Na_2HAsO_4 \cdot 7H_2O$. A coupling constant of approximately 44 Mc/sec was reported. They were unable to obtain the spectrum of Na_3AsO_4 and attributed this to the tetrahedral symmetry of the isolated AsO_4 groups. Reddock has obtained a coupling constant of approximately 78 Mc/sec for Nb^{93} in $NbCl_5$. Due to the large asymmetry of the crystal ($\eta = 0.32$), six resonance lines were observed instead of the usual four expected for a spin of 9/2.

The small quadrupole moment of the deuteron (43) and the predominantly spherical charge distribution near the nucleus combine to give deuteron quadrupole coupling constants of less than 1 Mc/sec. Hence any attempt to measure H^2 coupling constants must be done with spectrometers giving extremely high resolution. From measurements of hyperfine structure in the microwave spectrum of D_2O , Posener (44) obtained a coupling constant of 353 ± 4 Kc/sec for the deuteron in the direction of the OD bond. White (45) calculated the D coupling constants in DCN and DCCl from their pure quadrupole spectra and obtained values of 175 Kc/sec. and 290 Kc/sec.; he stated that the field gradient at the nucleus did not arise from any "unbalanced" p electrons, but was primarily due to charges residing on the adjacent C atoms and to overlap effects.

CONCLUSION

Nuclear quadrupole resonance techniques have been utilized to a great extent in the determination of the type of valence bonding in many halide compounds. It must be borne in mind that in the interpretation of the coupling constant one must consider all of the factors influencing it. The value obtained for the coupling constant of a given atom involved in a chemical bond is a function, not only of the ionicity of the bond, but of the degree of hybridization and π character existing in the bond as well.

BIBLIOGRAPHY

1. J. M. B. Kellogg, I. I. Rabi, N. F. Ramsey, and J. R. Zacharias, *Phys. Rev.*, 55, 318 (1939).
2. D. K. Coles and W. E. Good, *Phys. Rev.*, 70, 979 (1946).
3. H. G. Dehmelt and H. Krüger, *Naturwiss*, 37, 111 (1950).
4. T. P. Das and E. L. Hahn, "Nuclear Quadrupole Resonance Spectroscopy", Academic Press, Inc., New York, London (1958).
5. M. H. Cohen and F. Reif, in F. Seitz and D. Turnbull, "Solid State Physics", Vol. V, Academic Press, Inc., New York, London (1957).
6. C. T. O'Konski, in F. C. Nachod and W. D. Phillips, "Determination of Organic Structures by Physical Methods", Vol. II, Academic Press, Inc., New York, London (1962).
7. W. J. Orville-Thomas, *Quart. Revs.*, 11, 162 (1957).
8. W. Gordy, "Microwave Spectroscopy", John Wiley and Sons, Inc., New York (1953).
9. C. H. Townes and A. L. Schawlow, "Microwave Spectroscopy", McGraw-Hill Book Co., Inc., New York (1955).
10. D. K. Roe, *Inorganic Seminars*, 1957-58, 37.
11. R. Livingston, *Record Chem. Prog.*, 20, 173 (1959).
12. H. C. Longuet-Higgins, *Disc. Faraday Soc.* 19, 9 (1955).
13. R. V. Pound, *Phys. Rev.*, 79, 685 (1950).
14. C. H. Townes and B. P. Dailey, *J. Chem. Phys.*, 17, 782 (1949).
15. C. H. Townes and B. P. Dailey, *J. Chem. Phys.*, 23, 118 (1955).
16. W. Gordy, *J. Chem. Phys.*, 19, 792 (1951).
17. W. Gordy, *J. Chem. Phys.*, 22, 1470 (1954).
18. W. Gordy, *Disc. Faraday Soc.*, 19, 14 (1955).
19. L. Pauling, "The Nature of the Chemical Bond", Cornell Univ. Press, Ithaca, New York, 1960, p. 99.
20. B. P. Dailey, *J. Chem. Phys.*, 33, 1641 (1960).
21. R. Livingston, *Phys. Rev.*, 82, 289 (1951).
22. R. Livingston, *J. Phys. Chem.*, 57, 496 (1953).
23. H. Robinson, H. G. Dehmelt, and W. Gordy, *J. Chem. Phys.*, 22, 511 (1954).
24. J. M. Mays and B. P. Dailey, *J. Chem. Phys.*, 20, 1695 (1952).
25. A. L. Schawlow, *J. Chem. Phys.*, 22, 1211 (1954).
26. S. Kojima, K. Tsukada, S. Ogawa, and A. Shimauchi, *J. Chem. Phys.*, 21, 1415 (1953).
27. A. L. Allred and E. G. Rochow, *J. Inorg. Nucl. Chem.*, 5, 269 (1958); *ibid.* 265 (1958).
28. R. S. Drago, *J. Inorg. Nucl. Chem.*, 15, 237 (1960).
29. D. Nakamura, Y. Kurita, K. Ito and M. Kubo, *J. Am. Chem. Soc.*, 82, 5783 (1960).
30. K. Ito, D. Nakamura, Y. Kurita, and M. Kubo, *J. Am. Chem. Soc.*, 83, 4526 (1961).
31. D. Nakamura, K. Ito, and M. Kubo, *J. Am. Chem. Soc.*, 84, 163 (1962).
32. D. Nakamura, K. Ito, and M. Kubo, *Inorg. Chem.*, 1, 592 (1962).
33. R. P. Hamlen and W. S. Koski, *J. Chem. Phys.*, 25, 360 (1956).
34. A. H. Reddoch, *J. Chem. Phys.*, 35, 1085 (1961).
35. R. G. Barnes and S. L. Segel, *Phys. Rev. Letters*, 3, 462 (1959).
36. G. D. Watkins and R. V. Pound, *Phys. Rev.*, 85, 1062 (1952).
37. P. A. Casabella and P. J. Bray, *J. Chem. Phys.*, 28, 1182 (1958).
38. C. H. Townes and B. P. Dailey, *J. Chem. Phys.*, 20, 35 (1952).
39. H. Negita, P. A. Casabella, and P. J. Bray, *J. Chem. Phys.*, 32, 314 (1960).

40. C. T. O'Konski and T. J. Flautt, J. Chem. Phys., 27, 815(1957).
41. E. Brun, St. Hafner, and F. Hartmann, Helv. Phys. Acta, 33, 496 (1960)(C.A. 12045C (1961)).
42. E. D. Jones and E. A. Uehling, J. Chem. Phys., 36, 1690(1962).
43. H. G. Kolsky, T. E. Phipps, Jr., N. F. Ramsey, and H. B. Silsbee, Phys. Rev., 87, 395 (1952).
44. D. W. Posener, Australian J. Phys., 10, 276 (1957).
45. R. L. White, J. Chem. Phys., 23, 253 (1955).
46. M. A. Whitehead and H. H. Jaffe', Trans. Faraday Soc., 57, 1854 (1961).

COMPLEXES OF O-PHENYLENEBIS(DIMETHYLARSINE)

L. V. Interrante

November 27, 1962

INTRODUCTION

Among the species involving the arsenic atom as a donor, the complexes of o-phenylenebis(dimethylarsine)(PDA)(Fig. 1) have received by far the most attention.




Fig. 1

This particular interest in PDA has been largely due to the increase in the stability of its complexes over those of the monodentate tertiary arsines, as a consequence of the chelate effect. The intensity of this interest is illustrated by the range of elements for which PDA complexes have been reported (Table I).

TABLE I - Elements Reported to Coordinate with PDA

IVa	Va	VIa	VIIa	VIII			Ib	IIb	IIIb	IVb	Vb
Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	-	-	P
Zr	-	Mo	Tc	Ru	Rh	Pd	Ag	Cd	-	Sn	As
Hf	-	W	Re	Os	-	Pt	Au	Hg	-	-	Sb Bi

PDA was first synthesized by Chatt and Mann in 1939(1). It was later employed by these workers in a study of the structure of bridged dipalladium derivatives (2). Approximately ten years later, Nyholm (3) began the series of investigations of PDA complexes which has been largely responsible for the extent of knowledge in this area to date.

THEORY

The study of the complexes of PDA is to a large extent the study of spin-paired, covalent complexes to which the valence-bond theory can be applied with considerable success. The low permanent dipole moment and high polarizability of the tertiary arsine group are undoubtedly important factors promoting the formation of covalent complexes and hindering the formation of ionic complexes.

It has been suggested, however, that the major factor responsible for electron pairing and strong bonding in the PDA complexes is the capacity of the arsenic atoms to accept electrons from the metal in $\underline{d} - \underline{d}$ bonding (4). The theoretical feasibility of double bonds in such complexes has been established from overlap integral calculations (4). Further support has been obtained from infrared spectral studies of substituted metal carbonyls (5,6). In particular, the infrared spectrum and dipole moment of the nickel carbonyl derivative, $[\text{Ni}(\text{CO})_2\text{D}]$ [$\text{D} = o$ -phenylenebis(dimethylarsine)], have been examined and interpreted to show that the bond order of the arsenic-nickel bonds in this compound is between 1.4 and 1.7 (6).

The coordination numbers of many of the transition-metal PDA complexes can be understood in terms of the tendency of the metal ion to make use of all nine orbitals (five $(n-1)\underline{d}$, one \underline{ns} and three \underline{np}) for bonding pairs, for non-bonding pairs, or for single electrons (7). This "nine-orbital rule" may be thought of as an extension of the "inert-gas rule" for metal carbonyls. When the carbonyl groups are replaced by less strongly $\underline{\pi}$ -bonding groups, paramagnetism may arise with retention of the nine-orbital configuration for the complexed metal ion.

Table II (7) illustrates the dependence of the coordination number within each electronic configuration upon the number of unpaired electrons, which is, in turn, a function of the ligand field produced by metal ion and ligand.

TABLE II - Illustration of the Nine-Orbital Rule

Configuration	Unpaired Electrons	Coordination Number	Examples
\underline{d}^0	0	9	-
\underline{d}^1	1	8	$\text{K}_3\text{Mo}(\text{CN})_8, [\text{VD}_2\text{Cl}_4]$
\underline{d}^2	0 2	8 7	$\text{K}_3\text{Re}(\text{CN})_8, [\text{ReD}_2\text{Cl}_4]\text{ClO}_4$ $\text{K}_4\text{V}(\text{CN})_7$
\underline{d}^3	1 3	7 6	$\text{K}_4\text{Mo}(\text{CN})_7$ $\text{K}_3\text{Cr}(\text{CN})_6, [\text{CrD}_2\text{Cl}_2]\text{ClO}_4$
\underline{d}^4	0 2 4	7 6 5	$[\text{MoD}(\text{CO})_3\text{I}_2]$ $[\text{Cr}(\text{dipy})_3]^{2+}, [\text{ReD}_2\text{Cl}_2]^+$ $[\text{MnDBr}_2(\text{H}_2\text{O})]\text{Br}$

It should be noted that this rule, although useful for many covalently bonded complexes, is not necessarily obeyed for more ionic complexes.

Provided that bonding orbitals are available, the main factors which seem to govern the tendency to assume a certain coordination number are (i) steric effects and (ii) the resulting charge on the metal ion (8). The second factor is considered to be of major importance in the majority of complexes (9).

The "electroneutrality principle" proposed by Pauling (10) suggests that metal ions tend to have a net charge of zero. Thus a metal ion will add ligands until its net charge, as determined by the degree of electron transfer in the metal-ligand bonds, is close to this value. Important among the factors which determine the degree of electron transfer is double bonding between metal and ligand (9). Consequently, ligands such as PDA which are capable of relieving the negative charge on the metal might be expected to lead to higher coordination numbers for that metal. This is supported in general for those cases in which $d_{\pi} - d$ bonding is important, and is best exemplified by the preparation of five-coordinate PDA complexes of the d^7 and d^8 transition-metal ions (11, 12).

METAL DERIVATIVES OF PDA

The complexes of PDA with the transition metals are summarized in Tables III through VII. Discussion here will be confined only to those derivatives of particular interest.

A. Groups IVa and Va (Ti, Zr, Hf, and V)

Among the complexes of these metals with PDA, most noteworthy are the eight coordinate derivatives of the general formula $[MX_4D_2]$ ($M = Ti, Zr, Hf, V; X = Cl, Br$) (13, 14, 15). The titanium derivative, $[TiCl_4D_2]$, has been shown to have a dodecahedral structure by single-crystal X-ray analysis (14). This is the first example of a complex in which a first transition-series element exhibits eight coordination.

B. Group VIa (Cr, Mo, W)

The ability of PDA to enter into $d_{\pi} - d$ bonding is illustrated by its replacement of carbon monoxide from metal carbonyls with retention of the formal zero-valent oxidation state of the metal.

In the case of the Group VI carbonyls, $[M(CO)_6]$, the derivatives $[M(CO)_4D]$ and $[M(CO)_2D_2]$ ($M = Cr, Mo, W$) (16, 17) have been obtained by reaction with PDA. The action of halogens upon these derivatives leads to further replacement of carbon monoxide and the formation of the complexes $[MoD(CO)_3X_2]$ and $[MoD_2(CO)_2X]X$ ($X = Br, I$) (7). These compounds are diamagnetic and molecular-weight and molecular-conductivity measurements in nitrobenzene indicate that they are seven-covalent derivatives of bivalent molybdenum. Thus with the strongly π -bonding carbonyl groups in addition to PDA, complete spin-pairing can occur,

leading to expansion of the coordination number in accordance with the nine-orbital rule.

C. Group VIIa (Mn, Tc, Re)

The complexes of bivalent manganese with PDA constitute the only known examples of spin-free, outer-orbital complexes involving this ligand. The failure of previous attempts (4) to prepare PDA complexes of manganese and the instability of the derivatives finally obtained (18) illustrate the reluctance of PDA to coordinate under these conditions. Complexes obtained are of the general formula $[MnD_2X_2]$ ($X = Cl, Br, I$).

The complexes of bivalent rhenium and technetium with PDA correspond to the same general formula $[MD_2X_2]$. However, they differ appreciably from those of manganese(II) in that they are spin-paired and of considerably greater stability (19, 20).

This difference is also exhibited by the trivalent derivatives of these metals, which include the spin-free, apparently five-coordinate $[MnDCl_2(H_2O)]ClO_4$ (18) and the spin-paired, six-coordinate $[MD_2X_2]Y$ ($M = Tc, Re; X = Cl, Br, I; Y = ClO_4, X$) (19, 20). In this case, the reluctance of PDA to bond in an outer-orbital situation apparently results in a limitation of the coordination number of the manganese (III) derivative.

Oxidation of the trivalent rhenium and technetium derivatives yields a new series of diamagnetic, eight-covalent complexes of the type $[MD_2X_4]Y$, ($M = Re, Tc; X = Cl, Br; Y = ClO_4, X$), presumably containing the pentavalent metals (21, 22).

The partial replacement of carbon monoxide from the compound $Mn_2(CO)_{10}$ by PDA yields the complex, $[Mn(CO)_3D]$. Partial oxidation may be effected with halogen, leading, in the case of I_2 , to the spin-paired derivative of univalent manganese, $[MnD(CO)_3I]$ (23).

D. Group VIII


1. (Fe, Ru, Os)

In contrast to the isoelectronic manganese (II) ion, trivalent iron, ruthenium, and osmium, like rhenium(II) and technetium(II), yield six-covalent, spin-paired derivatives of the type $[MD_2X_2]Y$ ($M = Fe, Ru, Os; X = \text{halogen}; Y = ClO_4, X$) (3, 24, 25).

The ability of PDA to stabilize high as well as low oxidation states is illustrated by the oxidation of the trivalent iron and osmium complexes to the quadravalent compounds, $[MD_2X_2]Y_2$ (25, 26). This rather unusual oxidation state for iron is supported by magnetic measurements which indicate a d_c^4 configuration (26). Attempts to obtain the corresponding ruthenium(IV) derivative have as yet been unsuccessful (24).

The fact that complexes of the type $[MD_2X_2]$, as obtained with the iron group metals, should be capable of geometrical isomerism is illustrated by the reported isolation of cis-forms of the PDA and ditertiary phosphine complexes of ruthenium(II), $[RuL_2X_2]$ (X = halogen) (27). Unlike the trans forms previously obtained (24), the cis complexes are reduced by $LiAlH_4$ to the complex hydrides $[RuL_2HX]$, which in the case of L = PDA are unstable and decompose upon standing (27).

Replacement of carbon monoxide from metal carbonyls, with subsequent oxidation by halogens, has been employed in the case of iron pentacarbonyl with the subsequent synthesis of a compound in which iron has a formal oxidation state of +1 (28). The following reaction scheme summarizes the conversions possible in this system:



The coordination number of five for the d^7 configuration of iron(I) is also exhibited by the isoelectronic species, manganese(0), cobalt(II), and nickel(III), in the complexes, $[Mn(CO)_3D]$, $[CoBr_2triars]$, and $[NiBr_32ET_3P]$. By adding one more electron, another series of isoelectronic ions exhibiting five-coordination is obtained. These include iron(0), i.e. $[Fe(CO)D_2]$; cobalt(I), i.e. $[Co(RNC)_5]X$; nickel(II), i.e. $[NiD_2X_2]ClO_4$; and gold(III), i.e. $[AuD_2X][ClO_4]_2$ (28). It should be noted that this is just the coordination number predicted by the nine-orbital rule for the d^7 and d^8 ions.

2. (Co, Rh)

Besides the reported preparation of the completely analogous rhodium(III) complexes (29), only the cobalt derivatives of PDA have been investigated in any detail.

Especially interesting are the tris-PDA complexes obtained for both cobalt(II) and cobalt(III), $[CoD_3][ClO_4]_2$ and $[CoD_3][ClO_4]_3$. These are among the few complexes prepared having more than two PDA molecules per metal ion (4).

Geometrical isomerism in the complexes of general formula $[CoD_2X_2]ClO_4$ has been established by means of absorption spectroscopy. By virtue of the reported proximity of PDA to ethylenediamine in the spectrochemical series, it has been possible to distinguish the cis and trans isomers of these complexes by direct comparison of their visible absorption spectra with those of the corresponding ethylenediamine complexes (30).

3. (Ni, Pd, Pt)

The tendency of the d^8 metals to assume a coordination number of five and the multivalency of nickel when coordinated to PDA may account for the particular attention afforded PDA complexes of this subgroup.

In addition to the derivatives of bivalent nickel of empirical formula NiD_2X_2 ($X = Cl, Br, I, CNS, ClO_4$) (31), complexes containing tervalent $[NiD_2X_2] ClO_4$ (31, 32) and quadravalent nickel $[NiD_2X_2][ClO_4]_2$ (33) have been isolated. The complexes obtained with platinum and palladium are entirely analogous to the nickel compounds except for the fact that no complexes of the tervalent state of either metal could be obtained (2, 8, 34).

The structures of the complexes of bivalent nickel mentioned above were initially assumed to be four-covalent and planar, largely on the basis of their diamagnetism (31). Later work, however, has shown that, except for the di-perchlorates, these complexes, as well as those of palladium(II) and platinum(II), are apparently five-covalent in solution (35, 34, 8). In support of this is the reported isolation of the monoperchlorate salts, $[MD_2X] ClO_4$ ($M = Ni, Pd, Pt$) and the preparation of a similar complex of methylbis(3-dimethylarsinopropyl) arsine, $[Ni(\text{triars})Br_2]$ (36), which has been shown by single-crystal X-ray analysis to have a distorted square-pyramidal structure (37). The fact that this does not necessarily imply that the PDA complexes are similarly coordinated in the solid is indicated by X-ray work on the compound PdD_2I_2 , which crystallizes in the form of a distorted octahedron (8). Furthermore, the possibility that some interaction along the sixth coordinate is provided by the solvent molecules in solutions of these complexes must not be ignored (34, 12).

Replacement of carbon monoxide in the nickel-group metal carbonyls has led to a series of new PDA complexes. Partial replacement yields the nickel complex $[Ni(CO)_2D]$, which may be converted to a previously unknown class of cis-planar nickel complexes, $[NiX_2D]$ ($X = I, Br$) (38) by treatment with halogen. Complete replacement has also been effected by heating $[Ni(CO)_2D]$ with PDA in vacuum, leading to the zerovalent nickel complex $[NiD_2]$ (28). This compound and its palladium analog have also been prepared by direct reduction of the corresponding dipositive cationic species using sodium borohydride (39).

E. Group Ib (Cu, Ag, Au)

Complexes of PDA with the Group Ib metals invariably involve either the univalent or tervalent metals. In no instance, including copper, has a complex of the bivalent metal been obtained. This statement may be extended to include the d^9 configuration in general for which no PDA complex has been reported.

The following information is being furnished to you for your information and is not to be disseminated outside your organization.

In addition to the information provided in the attached report, the following information was obtained from the review of the records of the [redacted] office. The information is being furnished to you for your information and is not to be disseminated outside your organization.

The following information was obtained from the review of the records of the [redacted] office. The information is being furnished to you for your information and is not to be disseminated outside your organization. The information is being furnished to you for your information and is not to be disseminated outside your organization.

The following information was obtained from the review of the records of the [redacted] office. The information is being furnished to you for your information and is not to be disseminated outside your organization. The information is being furnished to you for your information and is not to be disseminated outside your organization.

The following information is being furnished to you for your information and is not to be disseminated outside your organization. The information is being furnished to you for your information and is not to be disseminated outside your organization.

The complexes of the univalent metals are isomorphous, four-covalent, presumably tetrahedral species of the type $[MD_2]Y$ ($M = Cu, Ag, Au$) (40, 41, 42, 43, 44). This coordination number, although commonly exhibited by copper(I), is rather unusual for gold(I), which as a rule prefers two-coordination. In this case the use of FDA is thought to favor formation of four-covalent structures since the chelate group prevents the use of linear sp bonds and so favors the addition of a second molecule of PDA (42).

In the case of gold these compounds may be oxidized to complexes containing the trivalent, d^8 ion. These include compounds in which gold appears to be four- five- and six-covalent (42). The bonding in these complexes is explained on the basis of valence-bond theory to involve the $6p_z$ and hybrid $6p_6d_z$ orbitals in addition to the $5d_6s_6p^2$ combination in the square, to account for five- and six-coordination, respectively (43, 42).

Due to the diffuse nature of the $6d_z$ orbital, it is expected that highly polar bonds would be formed. This is consistent with the conversion of the six-covalent derivatives to five-covalent complexes in water (43).

The electroneutrality principle has been employed to explain both the tendency of four-covalent gold(III) to expand its coordination number and the increased stability of the five-coordinate gold(III) complexes over the isoelectronic five-coordinate palladium(II) compounds (42).

F. Group IIb (Zn, Cd, Hg)

Reports of complexes involving the Group IIb metals and PDA have been confined to derivatives of the bivalent (d^{10}) metal ions. As expected these are characteristically four-coordinate and presumably tetrahedral, consisting of the non-electrolytes $[MDX_2]$ and the salts $[MD_2](ClO_4)_2$ ($M = Zn, Cd, Hg$; $X = Cl, Br, I$) (44, 45).

G. The Non-Transition Metals

Apparently due to the relatively poor donor ability of PDA toward metal ions in which $d_{\pi} - d_{\pi}$ bonding is not possible, few examples of non-transition metal complexes of PDA have been reported.

Among several non-transition metal tetra-halides tried, including UCl_4 (46), $ThCl_4$, $SiCl_4$, and $GeCl_4$ (14) only $SnCl_4$ was found to add PDA, forming the six-covalent derivatives, $[SnCl_4D]$ (46).

However with the group VI trihalides, FDA is reported to form five-covalent 1:1 addition compounds of the type $[MX_3D]$ ($M = Bi, Sb, As, P$; $X = Cl, Br, I$) (47, 48).

CONCLUSION

This seminar has attempted to summarize the known complexes of PDA as well as to examine the concepts of the nine-orbital rule and the electroneutrality principle.

It is hoped that the examples presented will serve to demonstrate the utility of these concepts and the valence bond theory in general in predicting the coordination numbers of complexes in which the bonding is primarily covalent.

TABLE III - $d^0 - d^5$ Complexes
D=0-phenylenebis(dimethylarsine)

Compound	Configuration	Unpaired Electrons	Coord'n Number	Oxidation State	References
[TiX ₄ D]; X = Cl, Br	d^0	--	6	IV	15, 13, 14
[MX ₄ D ₂]; M = Ti, Zr, Hf; X = Cl, Br	d^0	--	8	IV	13, 14
[TiX ₃ D(H ₂ O)]; X = Cl, Br	d^1	1	6	III	15
[VX ₄ D ₂]; X = Cl, Br	d^1	1	8	IV	13, 14
[MoDBr ₄]	d^2	2	6	IV	7
[ReD ₂ X ₄]Y; X = Cl, Br; Y = ClO ₄ , X	d^2	0	8	V	21
[TcD ₂ Cl ₄]Y; Y = ClO ₄ , Cl	d^2	0	8	V	22
[CrDX ₃ (H ₂ O)]; X = halogen	d^3	3	6	III	49
[CrD ₂ X ₂]Y; X = halogen, Y = ClO ₄ , CrDX ₄	d^3	3	6	III	49
[CrCl ₂ D(H ₂ O)(ETOH)]Cl	d^3	3	6	III	4
[MoD(CO) ₃ X ₂]; X = Br, I	d^4	0	7	II	7
[MoD ₂ (CO) ₂ X] X; X = Br, I	d^4	0	7	II	7
[MoD ₂ X ₂]; X = Cl, Br, I	d^4	2	6	II	50
[MD ₂ X ₂]Y; M = Re, Tc; X = Cl, Br, I; Y = ClO ₄ , X	d^4	2	6	III	19, 20
[FeD ₂ X ₂]Y ₂ ; X = Cl, Br; Y = FeCl ₄ , ClO ₄ , ReO ₄	d^4	2	6	IV	26
[OsD ₂ X ₂][ClO ₄] ₂ ; X = Cl, Br, I	d^4	2	6	IV	25
[MnDCl ₂ (H ₂ O)]ClO ₄	d^4	4	5	III	18
[MD ₂ X ₂]; M = Tc, Re; X = Cl, Br, I	d^5	1	6	II	19, 20
[MnD(CO) ₂ Br ₂]	d^5	1	6	II	23
[FeD ₂ X ₂]Y; X = Cl, Br; Y = ClO ₄ , FeX ₄ , Br	d^5	1	6	III	3
[RuD ₂ X ₂]Y; X = Cl, Br, I; Y = ClO ₄ , Br, RuDX ₄	d^5	1	6	III	24
[OsD ₂ X ₂]Y; X = Cl, Br, I, SCN	d^5	1	6	III	25
[MnD ₂ X ₂]; X = Cl, Br, I; Y = Cl, Br, ClO ₄	d^5	1	6	III	18
[MnD ₂ X ₂]; X = Cl, Br, I	d^5	5	6	II	18

doubtful in light of more recent work (14)
little evidence cited in support of this structure.

TABLE IV. d⁶ Complexes

Compound	Unpaired Electrons	Coord'n Number	Oxidation State	References
[M(CO) ₄ D]; M=Cr,Mo,W	0	6	0	16, 17
[M(CO) ₂ D ₂]; M=Cr,Mo,W	0	6	0	16, 17
[MnD(CO) ₃ I]	0	6	I	23
[MD ₂ X ₂]; M=Fe,Ru,Os X=Cl,Br,I,CNS	0	6	II	3,24,25,27
[Fe(CO) ₂ DX ₂]; X=Br,I	0	6	II	28
[RuD ₂ HCl]	0	6	II	27
[CoD ₂ X ₂]Y; X=Cl,Br,I,SCN Y=1/2 CoX ₄ ,ClO ₄ , X	0	6	III	51, 30
[CoD ₃][ClO ₄] ₃	0	6	III	4
[CoD ₂ (C ₂ H ₃ O ₂) ₂]ClO ₄	0	6	III	4
[RhD ₂ X ₂]X; X=Cl,Br,I	0	6	III	29
[NiD ₂ X ₂][ClO ₄] ₂ ; X=Cl,Br	0	6	IV	33
[MD ₂ X ₂]Y ₂ ; M=Pd,Pt X=Cl,Br,I Y=ClO ₄ , NO ₃	0	6	IV	34

TABLE V. d⁷ Complexes

Compound	Unpaired Electrons	Coord'n Number	Oxidation State	References
[Mn(CO) ₃ D] _n n = 1,2	1	5	0	23
[Fe(CO) ₂ DI]	1	5	I	28
[CoD ₂]Y ₂ Y=Cl,Br,I,CNS	1	4	II	51
[CoD ₃][ClO ₄] ₂	1	6	II	4
[NiD ₂ X ₂]Y X=Cl,Br,SCN Y=X,ClO ₄ , 1/2 PtCl ₆	1	6	III	31,32
[NiDBr ₃]		5	III	34

TABLE VI - d⁹ Complexes

Compound	Unpaired Electrons	Coord'n Number	Oxidation State	References
NiD ₃ [ClO ₄] ₂	0	6	II	31
AuD ₂ X ₂]Y X=Br,I; Y=ClO ₄ ,I	0	6	III	42
Fe(CO)D ₂]	0	5	0	28,16
Fe(CO) ₃ D]	0	5	0	28,16
MD ₂ X]Y M=Ni,Pd,Pt; X=Cl,Br,I, NO ₂ ,NO ₃ ,CNS; Y=ClO ₄ ,X	0	5	II	8,34,35
NiD ₂ ClO ₄]Y Y=ClO ₄ ,PF ₆	0	5	II	52,12
AuD ₂ X]Y ₂ X=Br,I; Y=ClO ₄ ,I	0	5	III	42
NiD ₂]X ₂ X=Cl,Br,I,CNS,	0	4	II	31,32
NiDX ₂] X=I,Br	0	4	II	38,34
PdCl ₂]	0	4	II	2
PdD ₂]Y ₂ Y=Cl, 1/2 PdCl ₄	0	4	II	2
MD ₂][ClO ₄] ₂ M=Ni,Pd,Pt	0	4	II	31,8,34
AuD ₂][ClO ₄] ₃	0	4	III	42

Little evidence cited in support of these structures
Doubtful in light of more recent work (34).

TABLE VII - d^{10} Complexes

Compound	Unpaired Electrons	Coord'n Number	Oxidation State	References
[NiD ₂]	0	4	0	39, 28
[PdD ₂]	0	4	0	39
[PdD(o-C ₆ H ₄ (PEt ₂) ₂)]	0	4	0	39
[Ni(CO) ₂ D]	0	4	0	38, 5
[CuD ₂]Y Y=ClO ₄ , CuX ₂ , Cl, Br, I X=Cl, Br, I	0	4	I	40
[AuD ₂]Y Y=Br, I, ClO ₄ , picrate, AuI ₂ , CuI ₂	0	4	I	41, 42, 43
[AgD ₂]Y Y=AgX ₂ , ClO ₄ X=Cl, Br, I, NO ₂	0	4	I	44
[MDX ₂] M=Zn, Cd, Hg X=Cl, Br, I	0	4	II	44, 45
[MD ₂][ClO ₄] ₂ M=Zn, Cd, Hg	0	4	II	44, 45

BIBLIOGRAPHY

- J. Chatt and F. G. Mann, J. Chem. Soc., 1939, 610.
- Ibid, 1622.
- R. S. Nyholm, J. Chem. Soc., 1950, 851.
- F. H. Burstall and R. S. Nyholm, J. Chem. Soc., 1952, 3570.
- R. S. Nyholm and L. N. Short, J. Chem. Soc., 1953, 2670.
- J. Chatt and F. A. Hart, J. Chem. Soc., 1960, 1378.
- H. L. Nigam, R. S. Nyholm, and M. H. B. Stiddard, J. Chem. Soc., 1960, 1806.
- C. M. Harris and R. S. Nyholm, J. Chem. Soc., 1956, 4375.
- R. S. Nyholm, Proc. Chem. Soc., 1961, 278.
- L. Pauling, J. Chem. Soc., 1948, 1461.
- R. S. Nyholm, Abstracts of Papers, 3N, 140th Meeting ACS, Chicago, Ill., Sept. 1961.
- C. M. Harris and S. E. Livingstone, Revs. Pure and Appl. Chem., 12, 16 (1962).
- R. J. H. Clark, J. Lewis, R. S. Nyholm, P. Pauling, and G. B. Robertson, Nature, 192, 222 (1961).
- R. J. H. Clark, J. Lewis, and R. S. Nyholm, J. Chem. Soc., 1962, 2460.
- G. J. Sutton, Australian J. Chem., 12, 122 (1959).
- H. L. Nigam and R. S. Nyholm, Proc. Chem. Soc., 1957, 321.
- H. L. Nigam, R. S. Nyholm, and M. H. B. Stiddard, J. Chem. Soc., 1960, 1803.
- R. S. Nyholm and G. J. Sutton, J. Chem. Soc., 1958, 564.
- N. F. Curtis, J. F. Fergusson, and R. S. Nyholm, Chem. and Ind. (London), 1958, 625.
- J. E. Fergusson and R. S. Nyholm, Nature, 183, 1039 (1959).
- J. E. Fergusson and R. S. Nyholm, Chem. and Ind. (London) 1958, 1555.
- Ibid., 1960, 347.
- Ibid., 1959, 130.
- R. S. Nyholm and G. J. Sutton, J. Chem. Soc., 1958, 567.
- Ibid., 572.

26. R. S. Nyholm and R. V. Parish, Chem. and Ind. (London), 1956, 470.
27. J. Chatt and R. G. Hayter, Proc. Chem. Soc., 1959, 153.
28. H. Nigam, R. S. Nyholm, and D. V. Ramana Rao, J. Chem. Soc., 1959, 1397.
29. R. S. Nyholm, J. Chem. Soc., 1950, 857.
30. T. M. Dunn, R. S. Nyholm, and S. Yamada, J. Chem. Soc., 1962, 1564.
31. R. S. Nyholm, J. Chem. Soc., 1950, 2061.
32. R. S. Nyholm, Nature, 165, 154 (1950).
33. R. S. Nyholm, J. Chem. Soc., 1951, 38.
34. C. M. Harris, R. S. Nyholm, and D. J. Phillips, J. Chem. Soc., 1960, 4379.
35. C. M. Harris, R. S. Nyholm, and N. C. Stephenson, Nature, 177, 1127. (1956).
36. G. A. Barclay, R. S. Nyholm, and R. V. Parish, J. Chem. Soc., 1961, 4433.
37. G. A. Mair, H. M. Powell, and D. E. Henn, Proc. Chem. Soc., 1960, 415.
38. R. S. Nyholm, J. Chem. Soc., 1952, 2906.
39. J. Chatt, F. A. Hart, and H. R. Watson, J. Chem. Soc., 1962, 2537.
40. A. Kabesh and R. S. Nyholm, J. Chem. Soc., 1951, 38.
41. R. S. Nyholm, Nature, 168, 705 (1951).
42. C. M. Harris and R. S. Nyholm, J. Chem. Soc., 1957, 63.
43. C. M. Harris, R. S. Nyholm, and N. A. Stephenson, Rec. trav. chim., 75, 687 (1956).
44. J. Lewis, R. S. Nyholm, and D. J. Phillips, J. Chem. Soc., 1962, 2177.
45. G. J. Sutton, Australian, J. Chem., 14, 545 (1961).
46. J. A. Allison and F. G. Mann, J. Chem. Soc., 1949, 2915.
47. G. J. Sutton, Australian J. Chem., 11, 415 (1958).
48. Ibid, 420.
49. R. S. Nyholm and G. J. Sutton, J. Chem. Soc., 1958, 560.
50. J. Lewis, R. S. Nyholm, and P. W. Smith, J. Chem. Soc., 1962, 2592.
51. R. S. Nyholm, J. Chem. Soc., 1950, 2071.
52. N. T. Barker, C. M. Harris, and E. D. McKenzie, Proc. Chem. Soc., 1961, 335.

THE ZERO-VALENT COMPLEXES OF PLATINUM

L. Jane Park

December 11, 1962




Studies of possible zero-valent platinum complexes did not proceed with any rapidity until about 1950. Previously, the only interest indicated was a cursory observation of the reaction of $K_2Pt(CN)_4$ with potassium metal in liquid ammonia, the product being apparently $K_4Pt(CN)_4$ (1). As recently as 1961, Nyholm and co-workers demonstrated that the compound $[(\phi_3P)_4Pt]$ was actually a hydride of bivalent platinum, $[(\phi_3P)_4PtH_2]$ (2). They then questioned the existence of any platinum(0) complex. Their work was followed, shortly thereafter, by the preparation of a compound which gave quite good evidence of being zero-valent.

This seminar will attempt to present the evidence for and against the existence of such compounds and to organize the data which has been published.

PREPARATION.

It might be expected that the zero-valent platinum compounds could be prepared by allowing the ligand to react with metallic platinum, analogously to the preparation of $Ni(CO)_4$. This, however, is not the case. In all of the cases observed, the starting material contained platinum in some oxidation state greater than zero, the most common being two. The platinum compound, usually a complex, is allowed to react with a reducing agent. Depending upon the particular reaction, a ligand may or may not be added.

Examples of the above are:



The compound $[Pt(NS)_4]$ has been prepared from H_2PtCl_6 and $4N_4$ (5), and its structure determined by x-ray (6). As there have been formed both $[Ni(CO)_4]$ and $[Ni(NS)_4]$, one might expect to form $Pt(CO)_4$. Although various partially carbonyl-substituted complexes have been prepared, e.g. $Pt(CO)[(p-Cl\phi)_3P]_2$ (7), no pure platinum carbonyl has been isolated. Carbonyl insertion reactions of a methyl platinum complex do not result in a zero-valent platinum complex (8).

Watt and his co-workers have prepared complexes of platinum(0) containing ammonia and ethylenediamine (9,10). Both are obtained by reduction of the corresponding platinum complex ion in liquid ammonia. In the preparation of $[Pt(N_2)_2]$, there are a number of other complexes formed as by-products.

THE UNIVERSITY OF CHICAGO

Page 100

October 15, 1950

Dear Mr. ... I have received your letter of the 10th and am glad to hear that you are interested in the ...

This is a very interesting problem and I would be pleased to discuss it with you if you have time ...

Sincerely,
[Signature]

I am sorry that I cannot give you a more definite answer at this time, but I will try to get back to you as soon as possible ...

Very truly yours,
[Signature]

(1) $\frac{1}{2} + \frac{1}{3} = \frac{5}{6}$

(2) $\frac{1}{4} + \frac{1}{8} = \frac{3}{8}$

(3) $\frac{1}{5} + \frac{1}{10} = \frac{3}{10}$

The above examples illustrate the addition of fractions with different denominators. In each case, the denominators are first made the same by finding a common denominator ...

It is important to note that the sum of two fractions is always less than the sum of their numerators divided by their common denominator ...

The formation of platinum(0) complexes has been studied polarographically (11). Since $[\text{Ni}(\text{CN})_4]^-$ has been reduced to $[\text{Ni}(\text{CN})_4]^{4-}$ rather than to the metal, and since Pt^{++} has the same outer electron configuration as Ni^{++} , it is to be expected that $[\text{Pt}(\text{CN})_4]^{4-}$ can be produced in this manner. Apparently, this experiment has not been attempted.

The reaction of cis- $[(\text{P}\phi_3)_2\text{PtCl}_2]$ with acetylene produces $[\text{Pt}(\text{P}\phi_3)_2\text{ac}]$ (12). Inasmuch as the acetylide ion carries a (-1) charge, the complex might be expected to contain formally univalent platinum. However, the reaction of $[\text{Pt}(\text{P}\phi_3)_2\text{O}-\text{C}_6\text{H}_4(\text{AsMe}_2)_2]$ with $\text{C}_2(\text{p-NO}_2\phi)_2$ produces $\text{Pt}(\text{P}\phi_3)_2\text{C}_2(\text{p-NO}_2\phi)_2$, which might be considered to contain platinum(0).

STABILIZATION OF THE ZERO-VALENT PLATINUM COMPLEXES.

The factors that lead to stabilization of the platinum(0) complexes can be applied to any compound in which the metal is in an abnormally low valence state.

Generally, one expects these complexes to comply with the Effective Atomic Number (E. A. N.) Rule. Thus, the sum of the electrons in platinum and those contributed by the ligand would be 86, corresponding to the radon configuration. An example of this is $[\text{Pt}(\phi_2\text{PCH}_2\text{CH}_2\text{P}\phi_2)_2]$. However, one would not put complete reliance on this rule to determine which compounds will be stable. Cases have been found in which the complexes contain electrons in excess of the rare-gas number, and also there is no account taken of the spatial requirements (7).

The π -bonding dative effect appears to play a large part in the stabilization, by highly electronegative atoms, of the low valence state. The π -type orbitals on the ligand must be vacant for this to occur. The metal electrons of the metal can then be fed into the ligand orbitals. This favors the donation of the ligand electrons to the metal by means of a sigma bond.

It is interesting to note that a ligand such as $[\text{O}-\text{C}_6\text{H}_4(\text{AsMe}_2)_2]$ forms platinum(0) complexes, but will stabilize abnormally high valence states as well. Filling the π -bonding orbitals of a ligand would effectively cause it to stabilize a high oxidation state.

COORDINATION NUMBER.

The coordination number of the platinum zero-valent complexes most often found is four. This complies with the E. A. N. rule. There appear to be, however, examples of coordination number other than four. If $\text{K}_2\text{Pt}(\text{CN})_4$ is allowed to react with a potassium acetylide, (KC_2R) , in liquid ammonia containing metallic potassium, there is formed a complex which analyzes as $\text{K}_2\text{Pt}(\text{C}\equiv\text{CR})_2$ (13). Assuming a negative charge in the hydrocarbon donor portion, the platinum has, at least formally, an oxidation state of zero.

Coordination number three is believed to occur in several platinum zero complexes. For example, the reaction of $[(p\text{-Cl}\phi)_3\text{P}]_2\text{PtI}_2$ with N_2H_4 produces $\text{Pt}[\text{P}(p\text{-Cl}\phi)_3]_3$ (14). The analogous fluoro compound has also been prepared (4). Apparently three halogen-substituted phenyl phosphines are sufficient to stabilize zero valent platinum, for no tetrakis derivatives have been found.

SPECTRAL STUDIES.

Various spectrophotometric techniques have been employed to determine the actual composition of platinum(0) complexes. It has been reported that upon the addition of K_2PtCl_4 to a solution of $\phi_3\text{P}$ in alcohol, $[\text{Pt}(\phi_3\text{P})_4]$ forms (15). However, the infrared spectrum shows bands which appear to be characteristic of the M-H stretch in metal hydrides; hence the compound is probably $[(\phi_3\text{P})_4\text{PtH}_2]$ and does not contain zero-valent platinum (2).

A spectrophotometric study of the amount of bromine consumed by the reaction $[(\phi_3\text{P})_4\text{PtH}_2] + 2\text{Br}_2 \rightarrow (\phi_3\text{P})_4\text{PtBr}_2 + 2\text{HBr}$ indicated two hydride ions in the coordination sphere.

Spectral studies have been used also to confirm the existence of a platinum(0) complex. Infrared and nuclear magnetic resonance spectra show no Pt-H bonds in $[\text{Pt}(\phi_2\text{PCH}_2\text{CH}_2\phi_2)_2]$ (3).

ANALOGOUS COMPOUNDS.

Inasmuch as platinum and palladium are in successive rows in the periodic table, it might be expected that the two metals would exhibit a great deal of similarity in the compounds which they form. This is found generally to be true. There are, however, some exceptions.

As was stated above, the supposed $[\text{Pt}(\phi_3\text{P})_4]$ was found to be $[(\phi_3\text{P})_4\text{PtH}_2]$. The compound $[\text{Pd}(\phi_3\text{P})_4]$ has been produced from $\text{H}_5\text{H}_5\text{Pd}$, C_6H_9 and $\phi_3\text{P}$, and no evidence is given for the possibility of hydride formation (16).

Two- and three-coordinate palladium complexes have been observed. The two-coordinate ones include the isonitriles, $[\text{Pd}(\phi\text{NC})_2]$ (17). The palladium compound, $[(p\text{-Cl}\phi_3\text{P})_3\text{Pd}]$, is quite analogous to the platinum compound (17).

The compound $\text{K}_4\text{Pd}(\text{CN})_4$ is known to exist and is apparently more soluble than the platinum analog (1). Experiments are hampered by the insolubility of the platinum complex.

A very interesting series of palladium compounds has been formed with ditertiary and tritertiary phosphines and ditertiary arsines (18). A novel type of isomerism appears when the ligand is a tritertiary phosphine. This arises from the number of phosphorus atoms coordinated to the platinum. The two types of ligands may be mixed in the complex. No evidence is found for metal-hydrogen bonding.

BIBLIOGRAPHY

1. J. J. Burbage and W. C. Fernelius, J. Am. Chem. Soc., 65, 1484 (1943).
2. J. A. Chopoorian, J. Lewis, and R. S. Nyholm, Nature, 190, 528 (1961).
3. J. Chatt and G. A. Rowe, Nature, 191, 1191 (1961).
4. A. D. Allen and C. D. Cook, Proc. Chem. Soc., 1962, 218.
5. E. Fluck, M. Goehring and J. Weiss, z. anorg. u. allgem. Chem., 287, 51 (1956).
6. I. Lindqvist and J. Weiss, J. Inorg. Nucl. Chem., 6, 184 (1958).
7. G. W. Watt, J. Electrochem. Soc., 1961, 67.
8. G. Booth and J. Chatt, Proc. Chem. Soc., 1961, 67.
9. G. W. Watt, M. T. Walling, Jr., and P. I. Mayfield, J. Am. Chem. Soc., 75, 6175 (1953).
10. G. W. Watt and J. W. Dawes, J. Am. Chem. Soc., 81, 8 (1959).
11. A. A. Vlcek, Nature, 177, 1943 (1956).
12. J. Chatt, G. A. Rowe, and A. A. Williams, Proc. Chem. Soc., 1957, 208.
13. R. Nast and W. Heinz, Ber., 95, 1478 (1962).
14. L. Malatesta and C. Cariello, J. Chem. Soc., 1958, 2323.
15. L. Malatesta and M. Angoletta, Atti. accad. nazl. Lincei, Rend., Classe sci. fis., mat., e nat., 19, 43 (1955).
16. E. O. Fischer and H. Werner, Ber., 95, 703 (1962).
17. L. Malatesta and M. Angoletta, J. Chem. Soc., 1957, 1186.
18. J. Chatt, F. A. Hart, and H. R. Watson, J. Chem. Soc., 1962, 2537.

CHELATE STABILITY AND SUBSTITUTION
ON AROMATIC LIGANDS

E. J. Olszewski

December 18, 1962


INTRODUCTION

As early as 1945, Calvin and Wilson(1) noted that no attempt had as yet been made to determine how the structural features of organic ligands could influence their tendency to form stable chelate compounds. Tate and Jones(2) have proposed a theoretical expression for the difference in stability for a pair of chelate complexes of the same metal ion having related but slightly different ligands. Martell and Calvin(3) cite inductive and steric effects and resonance interactions as the important factors involved when considering changes in chelate stability due to substitution of differing functional groups on aromatic ligands.


Tables of stability constant data have appeared as late as 1960 (4-7), but the majority of the publications dealing with this problem have appeared in the last two years. Previous to this, stability values of a specific class of aromatic ligands had to be obtained from varied sources. Thus, differences in temperature at which values were obtained or variations in solvent and in ionic strength allowed only for very general conclusions, and when different metal ions were used, for none at all.

DYES

A simple illustration of the relationship between stability and ring substitution can be seen by comparing Eriochrome-Black T with Eriochrome-Black B(8). Addition of a nitro group decreases stability due to a negative inductive



effect. Withdrawal of electron density from the coordinating hydroxyl oxygen is borne out by the lower stability values for both calcium and magnesium ions. For Eriochrome-Black A versus Eriochrome-Black R the same consideration of induction maintains, so that the lower log K_1 value for the divalent magnesium chelate of the former seems reasonable.



Eriochrome-Black B

Eriochrome-Black A

Eriochrome-Black R

Snavely, Fernelius, and Block(9) studied the chelate complexes of four pyrazolone dyes and found a direct relationship between the acid strength of the ligand and its power of coordination.

KETO-IMINES

Martin and co-workers(10) have discussed the stability of chelates of β -ketoimines and beryllium ion. The p-nitro group, as a result of its tendency to withdraw electrons, would increase electron density on the nitrogen by the following mechanism:



An electron donating group, such as Cl, also increases the electron density about the nitrogen atom with similar results:




Gould(11) postulates a positive resonance effect for the methoxy group, and a negative inductive influence. Thus the increase in stability of the ortho-methoxy substituted compound would be a natural consequence of resonance enhancement. In the 2,6-dimethyl substituent, steric effects decrease stability significantly. The p-methyl substituent appears as an anomaly. Having both positive inductive and resonance effects (due to hyperconjugation), it is still less stable than the unsubstituted compound.

QUINOLINES

Individual reports (12-19) had to be consulted for data on the derivatives of 8-hydroxyquinoline. Comparisons were difficult because of differences in ionic strength. The zinc(II) and magnesium(II) complexes of the 5-sulfonates are clearly less stable than those of the unsubstituted system. The negative inductive effect of the sulfonate group decreases electron density at the coordination center. Stability is increased by the 5-methyl substituent as a result of hyperconjugation. The same influence is extant in the 6-methylquinoline, but here there is no resonance interaction and as a consequence stability is not enhanced as greatly. Steric hindrance accounts for the lessened stability of the 7-methylquinoline compound, and the same reasoning applies to the 2-methyl substituent. The 4-methyl substitution increases electron density on the nitrogen atom as a result of resonance interaction making the Cu(II), Mg(II), Zn(II), and $UO_2(II)$ chelates more stable. Additional work on the effects of quinoline substitution is reported by Yasuda(20).


Lane and Mandathil(21) considered the series of compounds:



R = H, CH₃, OCH₃, OH

where A is a structural analog of 8-hydroxyquinoline with the same chelating system, -O-C=C-N=, except that the nitrogen atom is not part of a ring. B and C form larger chelate rings, and the effect of an ortho substituent can be observed. The effect of the electron donating groups is not as significant in A as in B and C. In all cases stability is increased, as demonstrated by the log K values. The large increase in stability of C over B is possibly due to an increase in entropy caused by the blocking effect of the methoxy group in the 3-position. The electron donating influence of the 3-methoxy group is also a factor.

The 4-hydroxybenzothiazoles are also similar to the 8-hydroxyquinolines (22).



R = H, NH₂, NHCH₃

R' = H, CH₃

The values obtained for the formation constants of the metal chelates of 2-amino-4-hydroxybenzothiazole, 2-methylamino-4-hydroxybenzothiazole, and 7-methyl-2-amino-4-hydroxybenzothiazole are higher than those obtained with the corresponding metal chelates of 4-hydroxybenzothiazole. The increase in stability is attributed to the increase of electron density in all cases at the centers of coordination. The metal species considered were Cu(II), Pb(II), Ni(II), Zn(II), Co(II), and Cd(II).

Feng and Fernando(23) studied some additional derivatives of this class of ligand. They used as metal species divalent zinc, cadmium, nickel, cobalt, and manganese. Their results supported those of Lane and Sam(22), and in addition they examined 5,7-dibromo-4-hydroxybenzothiazole and 4-hydroxybenzothiazole-7-sulfonate. For the former ligand, stability was diminished as a consequence of the negative inductive effect of the bromine pair. However, the 7-sulfonate derivative gave results which could not be explained.

1945

The first part of the report deals with the general situation in the country at the end of the war. It mentions the economic difficulties and the political changes that have taken place. The second part of the report is devoted to a detailed description of the various regions of the country. It discusses the agricultural situation, the state of the industry, and the social conditions in each of these regions. The third part of the report contains a summary of the main findings and conclusions of the study. It points out the most serious problems that the country is facing and suggests some possible solutions. The report ends with a list of references and an appendix containing some statistical data.

The following table shows the results of the survey in the various regions of the country.

Table 1

The results of the survey in the various regions of the country are shown in the following table. The table is divided into three main sections: Agriculture, Industry, and Social Conditions. Each section contains a list of the various regions and the results of the survey in each of them. The results are given in terms of the number of people surveyed, the main problems identified, and the suggestions made for improvement. The table shows that there are significant differences in the situation between the various regions, and that there are many common problems that affect a large number of the regions.

The following table shows the results of the survey in the various regions of the country. The table is divided into three main sections: Agriculture, Industry, and Social Conditions. Each section contains a list of the various regions and the results of the survey in each of them. The results are given in terms of the number of people surveyed, the main problems identified, and the suggestions made for improvement. The table shows that there are significant differences in the situation between the various regions, and that there are many common problems that affect a large number of the regions.

SALICYLALDEHYDES

Work has been done on many of the divalent copper derivatives of salicylaldehydes(24,25). The 3-ethoxy and 3-propyl substituents both increase stability. The steric factor does not seem to be operative here. Increased stability is observed for the 4-methoxy species also, but unlike the 3-ethoxy derivative, lack of resonance interaction diminishes its effectiveness. Small stabilization occurs with the 4-methyl, but more significantly with the 3,5-dimethyl where hyperconjugation can be more greatly felt in addition to the positive inductive effect. The negative inductive effect of the 5-sulfonate lessens the $\log \beta_2$ as do both the 3- and 4-nitrosalicylaldehydes.

The strong $-I$ effect of fluorine accounts for the lower stability of the 3-fluoro derivative. In the 3- and 5-chloro compounds a competition exists between the $-I$ and $+R$ effects. The lower $\log \beta_2$ value for the 3-chloro analog is due to the favored inductive effect of the closer chloride.

Jones and co-workers(26) studied the nitro-, chloro-, and methoxy- substituents each on the positions ortho, meta, and para to the OH group. The nitro group weakens stability the greatest in the ortho position whereas the chelate of the m-nitro is the most stable of the three as a result of negligible resonance interaction. For both the chloro and methoxy derivatives, the meta position again gives the most stable species, which is completely unexpected because of the unfavorable resonance interaction.

SALICYLATES

Of the salicylic acid type chelates, the iron(III) derivatives of the 5-sulfo and 5-amino species have been examined(27,28). The stability values decrease for the sulfo species and increase with the amino group. The negative inductive effect of the latter is not so effective due to its distance from the coordination center. Foye and Turcotte(29) in a more recent work reported similar results and extended their study of the 4-amino compound in addition to various nitro, chloro, and alkyl derivatives. They reported small variations in stability constants among the alkyl derivatives but stabilities which were generally slightly higher than that for the parent compound. The 5-chloro species gave slightly lower values whereas the greatest decrease in stability was found for the 3- and 5-nitro species. Tervalent iron and aluminum were the metal ions involved.

ANTHRANILATES

Harris and Sweet(30) considered the divalent copper and cadmium chelates of 5-sulfo, 3,5-diiodo, and 3-methyl anthranilic acids. Stability constant values varied slightly from those of the unsubstituted chelate, and the tendency was toward decreasing stability resulting from steric interaction. The work was expanded by Young and Sweet(31), who studied bulkier derivatives of anthranilic acid and attributed the decrease in stability to steric interactions.

PHENOLS

Sims(32) and Perrin(33) compared the stabilities of 3-hydroxyanthranilic acid and o-aminophenol chelates. For Mn(II), Fe(II), Co(II), Ni(II), and Cu(II), Sims found the o-aminophenolates to be more stable in all cases. Perrin studied just the iron ion and found the opposite to be true. The 3-amino-4-hydroxybenzenesulfonic acid gave less stable iron(II) chelates than the o-aminophenol because of withdrawal of electron density by the sulfo group.

The 2-nitrosophenol chelates of divalent cobalt and copper were studied by Shimura(34). The trend in stability is given as $5\text{-CH}_3 > 5\text{-OCH}_3 > 4\text{-CH}_3 > \text{Nph}, 4\text{-Cl} > 4\text{-Br}$. He also found the 4-methyl derivative to be consistently more stable than the 4-chloro for the divalent copper, nickel, lead, zinc, and cadmium ions.

AMINES

Silver complexes of aniline derivatives (35,36) comprise the next class of ligands. The negative inductive effect of the meta-nitro group diminishes $\log \underline{K}_2$ from 3.17 (for aniline) to 1.7. The additional factor of resonance enables the para-nitro constituent to cause even greater lowering of $\log \underline{K}_2$ to 1.6.

Positive inductive effects exert a stabilizing influence in all cases of substitution by methyl groups. The order of stability here is para > ortho > meta, with $\log \underline{K}_1$ values of 1.56, 1.51, and 1.47 respectively. The disubstituted methyl anilines give even more stable chelates.

Fyfe(37) compared the stability of aniline complexes of silver(I) with those of ortho-, meta-, and para-substituted toluidine, bromoaniline, chloroaniline, and nitroaniline. Stability was enhanced for the toluidines but lowered for the last three classes.

Armeanu and Luca(38) reported that the stability of complex silver ions was increased by the introduction of the methyl group into the aromatic aniline nucleus, and lowered by

substitution of -Cl, -NO₂, and -I. The authors claimed that the lower stability of the complex ions formed with o-toluidine and dimethyl-o-toluidine supported the hypothesis that the lower basicity of the methyl-o-toluidines has its cause in steric hindrance.

In ortho, meta, and para amine-substituted pyridine compounds (35, 39-43) stability is increased. The para substituent produces the greatest stability whereas the steric factor at the ortho position and the poor resonance interaction meta account for their less stable silver(I) complexes.

The positive resonance and inductive effects of the methyl group explain the stability increase of the o-, m-, and p-pyridine derivatives.. The effect of hyperconjugation by -CH₃ groups is not as great as the resonance interaction of -NH₂; thus the pyridyl amine chelates are the more stable. The three methyl groups of 2,4,6-trimethylpyridine cause an even greater increase in stability, but the 2,4-dimethyl member of the series gives the most stable chelates because of reduced steric hindrance.

Both resonance interaction and inductive influence are negative for the -CN group, explaining the lessened stability of these pyridyls.

For the nicotinamides and the methyl nicotinate there is a stability decrease, with the para compounds proving the least stable. The para position is more effective than the meta in raising the stability when the methoxy substituent is introduced.

Divalent cobalt, copper, and nickel complexes of picolinic acid and its 6-methyl derivative have been examined. The lower log K values for the metal chelates of the latter derivative is attributed to a steric factor.

DIPYRIDYLS

Chelate stability decreases in the case of both 5-chloro and 5-nitro substitution on the dipyridyls(44-46). Little change is observed in the 5-phenyl derivative, while with the 5-methyl substituent stability is increased. The log B₂ value of 7.9 for the 2-methyl iron(II) species is extremely low and is an example of the destabilizing effects of steric interaction. This effect is multiplied for the 2,2'-disubstituted dipyridyls (47), and a similar destabilization is found to occur for the 5,5'-disubstituted species by preventing the aromatic rings in the ligand from assuming a coplanar configuration (48).

OTHER LIGANDS

Singh and Banerjee(49) studied a group of violuric acids and found that the introduction of a methyl group in the side benzene ring enhances the stability of the metal chelates. The

effect of the methyl group is most pronounced when it is situated in the meta position and is less when it is in the para position. The low stability of the ortho-methyl chelate is attributed to steric hindrance.

Schwarzenbach and co-workers(50) and Calvin and Bailes(51) have reported studies on a number of metals forming chelates with variously substituted sulfonates and carboxylates of anilinediacetic acid.


In studying a number of related drugs, Andrews et al (52) have remarked that whereas only slight differences in chelate stability were noted, the resulting differences in physiological effects were considerable.

Tate and Jones(2) wanted to determine the usefulness of the Hammett equation to correlate stability constants of cadmium complexes for over twenty substituted benzenesulfonates. They concluded that structural parameters other than those used to represent inductive effects would be required. However they also noted that very few of the expected changes were found as substituents were varied from electron withdrawing to electron donating. They also reported no consistent obvious evidence of steric hindrance arising from ortho substitution.

CONCLUSION

Generally substitution effects on aromatic ligands will correspond with results predicted from theory. In only half of the cases studied is a linear relationship to be found between ligand basicity and chelate stability. A consideration of the relationship between ligand substitution and reaction rates would be a logical extension of this paper. Also interesting would be a study of ligand substitution effects on the oxidation potential of the metal chelates.

TABLE I



M = Be(II), Temp. = 30°C., Ionic Strength = 50% dioxane

Ligand: R' =	log K_1	log K_2	log B_2
Phenyl	10.87	10.49	21.36
4-Nitrophenyl	12.04	10.63	22.67
2-Methoxyphenyl	10.90	10.63	21.53
2,6-Dimethylphenyl	9.98	10.30	20.28
4-Chlorophenyl	11.47	10.75	22.22
4-Methylphenyl	10.43	10.61	21.04

TABLE II

8-Hydroxy Quinolines

Substituted Ligand	Divalent Metal	t	Ionic Strength	log K_1	log K_2	log K_{12}	Reference
8-Hydroxy-Quinoline	Cu	25°	50% dioxane	13.49	12.73	26.22	12
	"	20°	0.01	12.20	11.20	23.40	13
	"	25°	70% dioxane	15.00	14.00	29.00	14
	"	20°	50% dioxane	13.03	12.35	25.38	15
	Mg	25°	0.3M NaClO ₄	6.38	5.43	11.81	12
	"	25°	70% dioxane	6.88	5.96	12.84	14
	"	20°	50% dioxane	5.04	4.29	9.33	16
	Zn	25°	0.3M NaClO ₄	9.96	8.90	18.86	12
	"	25°	70% dioxane	10.91	9.90	20.81	14
	"	20°	50% dioxane	9.34	8.22	17.56	15
UO ₂	20°	0.3M NaClO ₄	11.25	9.64	20.89	15	
5-Sulfo	Cu	25°	→ 0	11.53	10.1	21.6	17
	"	20°	~0.01	12.5	10.6	23.1	13
	"	25°	"	11.5	10.7	22.2	18
	Mg	25°	→ 0	4.79	3.4	8.2	17
	"	20°	~0.01	4.8	3.7	8.5	13
	Zn	25°	→ 0	8.65	7.5	16.2	17
	"	20°	~0.01	8.4	6.7	15.1	13
	"	25°	"	8.7	7.2	15.7	18
5-Methyl	Cu	20°	50% dioxane 0.3M NaClO ₄	13.55	12.35	25.90	15
	Mg	"	"	5.21	4.47	9.68	"
	Zn	"	"	9.46	8.40	17.86	"
	UO ₂	"	"	11.25	9.52	20.77	"
6-Methyl	Mg	"	"	5.09	4.31	9.40	"
	UO ₂	"	"	10.89	9.26	20.15	"
7-Methyl	Mg	"	"	4.64	4.12	8.76	"
	UO ₂	"	"	11.28	9.78	21.06	"
	Zn	"	"	9.31	8.11	17.42	"
4-Methyl	Cu	25°	50% dioxane		13.52		19
	Mg	"	"	6.45	5.46	11.91	"
	Zn	"	"	10.67	9.57	20.24	"
2-Methyl	Cu	20°	50% dioxane 0.3M NaClO ₄	12.48	11.52	23.80	"
	"	25°	50% dioxane	10.22	9.32	19.54	"
	Mg	20°	50% dioxane 0.3M NaClO ₄	5.24	4.48	9.72	"
	"	25°	50% dioxane	3.73	3.13	6.86	"
	Zn	20°	50% dioxane 0.3M NaClO ₄	8.66	8.10	16.76	"
	UO ₂	"	"	~9.4	8	17.4	"

Financial Statement

Account	Balance	Debit	Credit	Balance	Account	Balance	Debit	Credit	Balance
101	100.00			100.00	101	100.00			100.00
102	100.00			100.00	102	100.00			100.00
103	100.00			100.00	103	100.00			100.00
104	100.00			100.00	104	100.00			100.00
105	100.00			100.00	105	100.00			100.00
106	100.00			100.00	106	100.00			100.00
107	100.00			100.00	107	100.00			100.00
108	100.00			100.00	108	100.00			100.00
109	100.00			100.00	109	100.00			100.00
110	100.00			100.00	110	100.00			100.00
111	100.00			100.00	111	100.00			100.00
112	100.00			100.00	112	100.00			100.00
113	100.00			100.00	113	100.00			100.00
114	100.00			100.00	114	100.00			100.00
115	100.00			100.00	115	100.00			100.00
116	100.00			100.00	116	100.00			100.00
117	100.00			100.00	117	100.00			100.00
118	100.00			100.00	118	100.00			100.00
119	100.00			100.00	119	100.00			100.00
120	100.00			100.00	120	100.00			100.00

TABLE III

4-Hydroxybenzothiazoles(22)

Temp. = 25°C., Ionic Strength = 50% dioxane

4-Hydroxybenzothiazole			2-Amino-4-Hydroxybenzothiazole			
	log K_1	log K_2	2 log K_{av} :	log K_1	log K_2	2 log K_{av} .
Cu(II)	9.0	8.0	17.1		9.8	21.1
Pb(II)	7.7	5.0	12.5	9.0	6.3	15.1
Ni(II)	7.3		13.5	8.0		
Zr(II)	6.7			8.0		
Co(II)	7.1			7.6	6.6	14.3
Cd(II)	6.9		12.3	7.9	6.1	14.2

2-Methylamino-4-Hydroxy-Benzothiazole			2-Amino-4-Hydroxy-7 Methylbenzothiazole			
	log K_1	log K_2	2 log K_{av} .	log K_1	log K_2	2 log K_{av} .
Cu(II)	10.9		21.3	11.5	10.5	21.7
Pb(II)	9.8	(9.0)	17.8	9.5	6.9	16.4
Zn(II)	8.3	7.9	16.2	8.5		(16.4)
Ni(II)	8.3	7.4	15.7	8.6	7.6	16.2
Co(II)				8.2	7.1	15.2
Cd(II)				8.2	7.0	15.1

TABLE IV

4-Hydroxybenzothiazoles(23)

Temp. = 25°C., Ionic Strength = 50% dioxane, 0.1 M NaCl

4-Hydroxybenzothiazole			2-Amino-4-Hydroxybenzothiazole	
	log K_1	log K_2	log K_1	log K_2
Zn(II)	6.90	5.92	8.0	7.4
Cu(II)	9.28	8.15	11.4	10.5
Ni(II)	7.35	6.11	8.1	7.2
Co(II)	6.88	6.60	7.8	6.8
Mn(II)	5.36	4.88	6.2	5.2

4-Hydroxybenzothiazole-7-Sulfonic Acid			5,7-Dibromo-4-Hydroxybenzothiazole
	log K_1	log K_2	log K_i
Zn(II)	6.1	4.3	
Cu(II)	9.5	7.8	7.1
Ni(II)	8.1	5.9	5.4
Co(II)	7.9	5.8	4.9
Mn(II)	5.1	3.9	

TABLE V
Salicylaldehydes
M=Cu(II)

Substituted Ligand	t	Ionic Strength	log β_2	Reference
Salicylaldehyde	25°C	50% dioxane	13.31	24
3-Propyl	20°C	50% dioxane \approx 0.1	14.2	1
3-Ethoxy	20°C	50% dioxane \approx 0.1	14.2	1
4,6-Dimethyl	20°C	50% dioxane \approx 0.1	14.4	1
4-Methoxy	20°C	50% dioxane \approx 0.1	13.8	1
5-Methyl	20°C	50% dioxane \approx 0.1	13.4	1
5-Nitro	20°C	50% dioxane \approx 0.1	8.8	1
4-Nitro	20°C	50% dioxane \approx 0.1	9.2	1
3-Nitro	20°C	50% dioxane \approx 0.1	8.2	1
5-Sulfo	25°C 40°C	\approx 0.2 \approx 0.2	9.27 9.79	27 27
3-Floro	20°C	50% dioxane \approx 0.1	11.4	1
5-Chloro	20°C	50% dioxane \approx 0.1	11.4	1
3-Chloro	20°C	50% dioxane \approx 0.1	10.6	1

TABLE VI

Salicylates (29)

Temp. = 20°C., Ionic Strength = 95% ethanol

	Fe(III)			Al(III)		
	log K_1	log K_2	log K_5	log K_1	log K_2	log K_5
4-amino-	16.30	13.91	30.48	15.57	12.45	28.02
4-methyl-	16.26	11.89	29.84	15.76	11.65	27.41
3,5-diisopropyl-	16.23	11.59	29.66	15.62	11.75	27.37
5-tert-butyl-	16.06	12.21	29.44	15.60	11.56	27.16
Salicylic Acid	16.53	12.69	29.34	15.62	11.45	27.07
3-isopropyl-6-methyl-	16.03	11.34	29.08	15.45	11.50	26.95
3-methyl-	15.99	11.85	29.04	15.26	11.53	26.79
5-chloro-	16.07	12.82	28.96	16.02	11.73	27.75
3-nitro-	14.60	12.46	27.54	14.25	11.85	26.10
5-nitro	14.51	12.67	26.86	14.31	12.16	26.47

TABLE VII

Anthranilates(30)

	Divalent Metal	$K_1 \times 10^{-4}$
3-methylantranilic acid	Cd	0.32
	Cu	7.9
N-methylantranilic acid	Cd	0.30
	Cu	3.7
Anthranilic acid	Cd	0.16
	Cu	14
5-sulfoanthranilic acid	Cd	0.26
	Cu	8.7
N-phenylantranilic acid	Cd	0.21
	Cu	0.63
3,5-diiodoanthranilic acid	Cd	0.05
	Cu	0.45

TABLE VIII

Phenols(32)

Ligand	pK _a	pK _a	Ni(II)		Co(II)	Cd(II)	Mn(II)
			K_1	K_2	K_1	K_1	K_1
2-Hydroxyanthranilic Acid	10.09	5.20	5.1	3.6	4.4	4.3	3.4
2-Amino-4,5-dimethylphenol hydrochloride	10.40	5.28	5.9	4.8	5.3	4.9	3.6
2-Aminophenol hydrochloride	9.99	4.86	5.4	-	4.7	4.3	3.6

continued

Phenols (33)

	Mn(II)	Fe(II)	Co(II)	Ni(II)	Cu(II)
3-Hydroxyanthranilic acid	3.4	7.7	4.4	5.1	-
O-aminophenol	3.6	8.0	4.7	5.4	8.8

P. Sims, J. Chem. Soc., 3648 (1959)

		log K_1	log $\frac{K_2}{I}$	Ionic Strength
O-aminophenol	Fe(II)	3.66	6.34	0.006
	Cu(II)	8.77	16.14	0.002
3-amino-4-hydroxy-	Fe(II)	3.32	6.13	0.009
benzenesulfonic acid				
3-Hydroxyanthranilic acid	Fe(II)	3.8	8.3	0.002
4-amino-5-hydroxy	Fe(II)	2.57	5.5	0.018
naphthalene-2,7-	Cu(II)	6.91	11.15	0.020
disulfonic acid (mono-sodium salt)				

TABLE IX

Anilines

Temp = 25°C.,

M = Ag(I)

Ligand	Ionic Strength	log K_1	log $\frac{K_2}{I}$	Reference
Aniline	1M KNO ₃ 50% mol. C ₂ H ₅ OH	1.44		36
m-nitroaniline	"		3.17	35
p-nitroaniline	"		1.7	"
o-methylaniline	"		1.6	"
"	"		3.17	"
m-methylaniline	1M KNO ₃	1.51		36
p-methylaniline	"	1.47		"
"	"	1.56		"
	50% mol. C ₂ H ₅ OH			
3,5-dimethylaniline	1M KNO ₃	1.63	3.48	35
2,6-dimethylaniline	"	1.62		36

Anilines(37)

M = Ag(I), Temp.=25°C., Ionic Strength= 59% Wt. ethanol-H₂O

Ligand	log K_f	Ligand	log K_f
Aniline	3.0		
<u>O</u> -Toluidine	3.65	<u>O</u> -chloroaniline	2.5
<u>m</u> -Toluidine	3.4	<u>m</u> -chloroaniline	2.55
<u>p</u> -Toluidine	3.9	<u>p</u> -chloroaniline	2.5
<u>o</u> -Bromoaniline	2.8	<u>o</u> -nitroaniline	1.9
<u>m</u> -Bromoaniline	2.8	<u>m</u> -nitroaniline	1.7
<u>p</u> -Bromoaniline	2.75	<u>p</u> -nitroaniline	1.7

TABLE X
Pyridines

Temp. = 25°C.

M = Ag(I)

Ligand	Ionic Strength	log K_1	log K_2	log β_2	Reference
Pyridine	→ 0	1.97	2.38	4.35	39
"	→ 0			3.82	40
"	0.5KNO ₃	2.04	2.18	4.22	41
"	50% mol. C ₂ H ₅ OH			3.88	35
o-aminopyridine	?	2.86	2.89	5.75	42
p-aminopyridine	?	2.17	3.04	5.21	"
m-aminopyridine	?	2.80	3.40	6.20	"
o-methylpyridine	0.5M KNO ₃	2.27	2.41	4.68	41
p-methyl pyridine	→ 0	2.00	2.35	4.35	39
m-methyl pyridine	0.5M KNO ₃	2.24	2.46	4.70	41
"	→ 0	2.03	2.36	4.39	"
o-cyanopyridine	"			2.90	39
p-cyanopyridine	"			3.08	"
2,4,6-dimethylpyridine	50% mol. C ₂ H ₅ OH			4.75	35
2,4-dimethylpyridine	0.5M KNO ₃	2.47	2.71	5.18	41
nicotinamide	→ 0			3.22	39
iso-Nicotinamide	"			3.01	"
o-methoxypyridine	"	1.58	2.09	3.67	"
p-methoxypyridine	"	2.28	2.16	4.44	"
ethyl nicotinate	"			2.99	"
ethyl iso nicotinate	"			2.45	"

TABLE XI
Phenanthrolines
M=Fe(II)

Ligand	Temp.	Ionic Strength	log K_1	log K_2	log β_2	log K_3	log β_3	Reference
1:10-phenanthroline	25°C.	→ 0			21.3			45
"	"	"			21.5			"
o-Nitro-1:10-phenanthroline	"	"			17.8			46
p-Chloro-1:10-phenanthroline	"	"			19.7			"
o-Methyl-1:10-phenanthroline	20°C	→	4.2	3.7	7.9	2.9	10.8	47
p-Methyl-1:10-phenanthroline	"	"					22.3	46
o-Phenyl-1:10-phenanthroline	25°C	-			21.1			"

TABLE XII

Anilinediacetic Acids(50,51)
Temp.=20°C., Ionic Strength = 0.1

Ligand	Metal	log K_1
Anilinediacetic acid	Mg(II)	1.15
	Ca(II)	0.6
	Fe(II)	2.01
2-Hydroxyaniline diacetic acid	Fe(II)	7.69
2-Sulfoaniline diacetic acid	Mg(II)	2.68
	Ca(II)	4.57
3-Sulfoaniline diacetic acid	Mg(II)	1.26
	Ca(II)	1.26
4-Sulfoaniline diacetic acid	Mg(II)	1.15
	Ca(II)	0.95
2-Carboxyanilinediacetic acid	Mg(II)	3.91
	Ca(II)	5.06
3-Carboxyanilinediacetic acid	Mg(II)	1.38
	Ca(II)	1.46
4-Carboxyanilinediacetic acid	Mg(II)	1.30
	Ca(II)	1.30

TABLE XIII

Benzenesulfonates(2)

M=Cd(II), Temp. = 25°C.

Ligand	K_1
1. 4-Aminobenzenesulfonate	0.1983
2. 4-Methylbenzenesulfonate	0.3604
3. Benzenesulfonate	0.4185
4. 2-Chloroaniline-4-sulfonate	0.5451
5. Benzene-1,3-disulfonate	0.6960
6. 4-Chlorobenzenesulfonate	0.8540
7. 4-Bromobenzenesulfonate	0.9215
8. 2,4,6-Trimethylbenzene-sulfonate	1.505
9. 2,5-Dimethylbenzenesulfonate	1.586
10. 4-Amino-3-methylbenzenesulfonate	1.743
11. 3-Aminobenzenesulfonate	1.759
12. 2-Aminophenyl-4-sulfonate	2.247
13. 2,5-Dichloro-4-aminobenzenesulfonate	2.411
14. 4-Chloro-3-methylbenzenesulfonate	2.460
15. 2,4-Dimethylbenzenesulfonate	2.700
16. 2-Hydroxy-4-methoxybenzophenone-5-sulfonate	2.748
17. 4-Hydroxybenzenesulfonate	2.822
18. 4-Toluidine-2-sulfonate	2.839

1911
 1912
 1913
 1914
 1915
 1916
 1917
 1918
 1919
 1920
 1921
 1922
 1923
 1924
 1925
 1926
 1927
 1928
 1929
 1930
 1931
 1932
 1933
 1934
 1935
 1936
 1937
 1938
 1939
 1940
 1941
 1942
 1943
 1944
 1945
 1946
 1947
 1948
 1949
 1950
 1951
 1952
 1953
 1954
 1955
 1956
 1957
 1958
 1959
 1960
 1961
 1962
 1963
 1964
 1965
 1966
 1967
 1968
 1969
 1970
 1971
 1972
 1973
 1974
 1975
 1976
 1977
 1978
 1979
 1980
 1981
 1982
 1983
 1984
 1985
 1986
 1987
 1988
 1989
 1990
 1991
 1992
 1993
 1994
 1995
 1996
 1997
 1998
 1999
 2000
 2001
 2002
 2003
 2004
 2005
 2006
 2007
 2008
 2009
 2010
 2011
 2012
 2013
 2014
 2015
 2016
 2017
 2018
 2019
 2020
 2021
 2022
 2023
 2024
 2025
 2026
 2027
 2028
 2029
 2030
 2031
 2032
 2033
 2034
 2035
 2036
 2037
 2038
 2039
 2040
 2041
 2042
 2043
 2044
 2045
 2046
 2047
 2048
 2049
 2050
 2051
 2052
 2053
 2054
 2055
 2056
 2057
 2058
 2059
 2060
 2061
 2062
 2063
 2064
 2065
 2066
 2067
 2068
 2069
 2070
 2071
 2072
 2073
 2074
 2075
 2076
 2077
 2078
 2079
 2080
 2081
 2082
 2083
 2084
 2085
 2086
 2087
 2088
 2089
 2090
 2091
 2092
 2093
 2094
 2095
 2096
 2097
 2098
 2099
 2100

Year	Month	Day	Event
1911	Jan	1	...
1911	Jan	2	...
1911	Jan	3	...
1911	Jan	4	...
1911	Jan	5	...
1911	Jan	6	...
1911	Jan	7	...
1911	Jan	8	...
1911	Jan	9	...
1911	Jan	10	...
1911	Jan	11	...
1911	Jan	12	...
1911	Jan	13	...
1911	Jan	14	...
1911	Jan	15	...
1911	Jan	16	...
1911	Jan	17	...
1911	Jan	18	...
1911	Jan	19	...
1911	Jan	20	...
1911	Jan	21	...
1911	Jan	22	...
1911	Jan	23	...
1911	Jan	24	...
1911	Jan	25	...
1911	Jan	26	...
1911	Jan	27	...
1911	Jan	28	...
1911	Jan	29	...
1911	Jan	30	...
1911	Jan	31	...

1911

Year	Month	Day	Event
1911	Jan	1	...
1911	Jan	2	...
1911	Jan	3	...
1911	Jan	4	...
1911	Jan	5	...
1911	Jan	6	...
1911	Jan	7	...
1911	Jan	8	...
1911	Jan	9	...
1911	Jan	10	...
1911	Jan	11	...
1911	Jan	12	...
1911	Jan	13	...
1911	Jan	14	...
1911	Jan	15	...
1911	Jan	16	...
1911	Jan	17	...
1911	Jan	18	...
1911	Jan	19	...
1911	Jan	20	...
1911	Jan	21	...
1911	Jan	22	...
1911	Jan	23	...
1911	Jan	24	...
1911	Jan	25	...
1911	Jan	26	...
1911	Jan	27	...
1911	Jan	28	...
1911	Jan	29	...
1911	Jan	30	...
1911	Jan	31	...

TABLE XIII (CONTINUED)

Ligand	K_1
19. 1-Methylbenzenesulfonate	3.153
20. 2-Chloroaniline-5-sulfonate	3.213
21. 1-Benzaldehydesulfonate	3.496
22. 3-Toluidine-4-sulfonate	12.76
23. 2,5-Dichlorobenzenesulfonate	13.92
24. Aniline-N-methylsulfonate	29.52

REFERENCES

1. M. Calvin and K. W. Wilson, J. Am. Chem. Soc., 67, 2003 (1945).
2. J. F. Tate and M. M. Jones, J. Am. Chem. Soc., 83, 3024 (1961).
3. A. E. Martell and M. Calvin, "Chemistry of the Metal Chelate Compounds", Prentice-Hall, Inc., Englewood Cliffs, N.J., 1959, p. 171.
4. S. Chaberek and A. Martell, "Organic Sequestering Agents", Wiley and Sons, Inc., New York, 1959.
5. K. B. Yatsimirskii and V. P. Vasil'ev, "Instability Constants of Complex Compounds", Consultants Bureau, New York, 1960.
6. R. L. Smith, "The Sequestration of Metals", Chapman and Hall, Ltd., London, 1959.
7. J. Bjerrum, G. Schwarzenbach, and L. Sillen, "Stability Constants, Part I: Organic Ligands", Special Publication No. 6, The Chemical Society, London, 1957.
8. G. Schwarzenbach and W. Biederman, Helv. Chim. Acta., 31, 678 (1948).
9. F. A. Snavely, W. C. Fernelius and B. P. Block, J. Am. Chem. Soc., 79, 1028 (1957).
10. D. F. Martin, G. A. Janusonis and B. B. Martin, J. Am. Chem. Soc., 83, 73 (1961).
11. E. S. Gould, "Mechanism and Structure in Organic Chemistry", Holt and Co., New York, 1959.
12. W. D. Johnston and H. Freiser, J. Am. Chem. Soc., 74, 5239 (1952).
13. A. Albert, Biochem. J., 54, 646 (1953).
14. L. E. Maley and D. P. Mellor, Austral. J. Sci. Res., 2, 92 (1949).
15. H. Laving and H. S. Rossotti, J. Chem. Soc., 2910 (1954).
16. R. Naanen, Suomen, Kem., 26, 11 (1953).
17. R. Naanen and E. Uisitalo, Acta. Chem. Scand., 8, 112 (1954).
18. L. E. Maley and D. P. Mellor, Acta. Chem. Scand., 8, 579 (1954).
19. W. D. Johnston and H. Freiser, Analyt. Chim. Acta., 11, 201 (1954).
20. M. Yasuda, Z. Physik. Chem., 29, 377 (1961).
21. T. J. Lane and A. J. Kandathil, J. Am. Chem. Soc., 83, 3782 (1961).
22. T. J. Lane and A. Sam, J. Am. Chem. Soc., 83, 2223 (1961).
23. P. K. Peng and Q. Fernando, Inorg. Chem., 1, 426 (1962).

24. L. E. Maley and D. F. Mellor, *Nature*, 159, 370 (1947).
25. M. Calvin and H. C. Melchior, *J. Am. Chem. Soc.*, 70, 3270 (1948).
26. J. G. Jones, J. B. Poole, J. C. Tomkinson, and R. J. P. Williams, *J. Chem. Soc.* 2001 (1958).
27. A. Agren, *Acta. Chem. Scand.*, 9, 39(1955).
28. L. Vareille, *Bull. Soc. Chim. France*, 872 (1955).
29. W. O. Foye, and J. G. Turcotte, *J. Pharm. Sci.*, 51, 329(1962).
30. W. F. Harris and T. R. Sweet, *J. Phys. Chem.*, 60, 509 (1956).
31. A. Young and T. R. Sweet, *J. Am. Chem. Soc.*, 80, 800 (1958).
32. P. Sims, *J. Chem. Soc.*, 3648 (1959).
33. D. D. Perrin, *J. Chem. Soc.*, 2245 (1961).
34. H. Shimura, *Nippon Kagaku Zasshi*, 82, 641 (1961).
35. E. Larsson, *Z. phys. Chem.*, 169A, 207 (1934).
36. C. Golumbic, *J. Am. Chem. Soc.*, 74, 5777 (1952).
37. W. S. Fyfe, *J. Chem. Soc.*, 2018 (1952).
38. V. Armeanu and C. Luca, *Z. phys. Chem.*, 214, 81 (1960).
39. R. K. Murmann and F. Basolo, *J. Am. Chem. Soc.*, 77, 3484(1955).
40. W. S. Fyfe, *Nature*, 169, 69 (1952).
41. R. J. Bruehlmann and F. H. Verhoek, *J. Am. Chem. Soc.*, 70, 1401 (1948).
42. H. B. Jonassen and C. C. Rolland, Report to the Office of Naval Research, Project No. Nnr, 685.
43. F. Holmes and W. R. Crimmins, *J. Chem. Soc.*, 1175 (1955).
44. T. S. Lee, L. M. Kolthoff, and D. Leussing, *J. Am. Chem. Soc.* 70, 2348 (1948).
45. W. W. Brandt and D. K. Gullstrom, *J. Am. Chem. Soc.*, 74, 3532 (1952).
46. H. Irving and R. Williams, *J. Chem. Soc.*, 3417 (1953).
47. F. H. Burstall, *J. Chem. Soc.*, 1662 (1938).
48. F. W. Cagle and G. F. Smith, *J. Am. Chem. Soc.*, 69, 1860(1947).
49. R. P. Singh and N. R. Banerjee, *Jour. Indian Chem. Soc.*, 37, 713 (1960).
50. G. Schwarzenbach, A. Willi, and R. O. Bach, *Helv. Chim. Acta.*, 32, 1175 (1949).
51. M. Calvin and R. H. Bailes, *J. Am. Chem. Soc.*, 68, 949(1946).
52. A. C. Andrews, T. D. Lyons and T. D. O'Brien, *J. Chem. Soc.*, 1776 (1962).

CHEMICAL EFFECTS OF NUCLEAR TRANSFORMATIONS
IN THE GAS PHASE

Agnes Chia Wang

January 8, 1963

I. INTRODUCTION

The product atom from a nuclear reaction usually acquires a large momentum and often possesses kinetic and electronic energies far in excess of chemical bond energies. Such excited atoms then move through the surrounding medium while losing energy and being slowed down by collision. During these collisions, the recoil atom may undergo reaction leading to stable chemical combination. The area of "hot atom" chemistry involves the study of these chemical effects of nuclear transformations.

From certain points of view, we can classify the studies into four groups:

1. Chemistry of atoms formed by induced nuclear reactions.
-Product isotopic with the target (Szilard-Chalmers type of reaction); for example, $C^{12}(\underline{\gamma}, \underline{n})C^{11}$ and $(\underline{n}, \underline{\gamma})$ reactions of halogens.
-Product non-isotopic with the target; for example, $N^{14}(\underline{n}, \underline{p})C^{14}$, $He^3(\underline{n}, \underline{p})T$ and $Li^6(\underline{n}, \underline{\gamma})T$ reactions.
2. Chemistry of atoms formed by isomeric transition; for example, $Br^{80m} \rightarrow Br^{80}$.
3. Chemistry of atoms formed by alpha-decay; for example, the study of Pb^{212} .
4. Chemistry of atoms formed by beta-decay; for example, the study of Bi^{210} and beta-decay product of tritium.

This seminar will be limited to discussions of the gas-phase reactions only. For work in this field as well as on liquid and solid phase reactions, a number of general review articles are available. (1,2,3,4,5)

II. THEORETICAL STUDIES

Theoretical studies in this field are still in their infancy (6). A number of workers have proposed different models in an attempt to describe the mechanisms of the hot-atom reactions, such as the billiard ball model of elastic collision by Libby and coworkers (7); the Estrup-Wolfgang model (8), the simple model for competitive tritium reactions by Rowland and coworkers (9); and the mathematical model formulated by Gordus and Hsuing (10). The last authors also published calculated results concerning momentum transfer to an atom in a molecule (11), and failure to bond-rupture following nuclear recoil from $(\underline{n}, \underline{\gamma})$ activation (12). In all the cases, a number of assumptions have to be made.

III. EXAMPLES OF EXPERIMENTAL RESULTS

1. Halogens, Cl^{38} , Br^{80} , Br^{80m} and I^{128}

Recoil halogen atom reactions in the gas phase, $Cl^{37}(n,\gamma)Cl^{38}$ (13), $Br^{79}(n,\gamma)Br^{80}$ (14), and $I^{127}(n,\gamma)I^{128}$ (15) were studied by Willard and coworkers. They found that the recoil halogen atoms in the gaseous hydrocarbons were able to replace the hydrogen atom. The organic yields ranged from a few percent for chlorine to about 54% in the case of iodine in the presence of methane. These facts indicate that some of the halogen atoms produced by (n,γ) processes can react with a hydrocarbon chain in a one-step reaction to become organically bound.

Recent studies by Rack and Gordus of the same reactions in the presence of inert-gas moderators show that, in the case of bromine, the reaction with methane occurs principally as a result of the recoil kinetic energy acquired by the Br^{80} in the (n,γ) activation process (16). For iodine, their data suggested that of the 54.4% organic yield as a consequence of the (n,γ) activation, about 18% forms as a result of hot I^{128} reactions, 11% as a product of excited iodine atoms or I^+ ions in the 3P_2 , 3P_1 , and/or 3P_0 states, and 25% as a result of reactions of $I^{\frac{3}{2}}(^1D_2)$ ions (17). They also reported that the manner in which molecular additions inhibit the reaction of (n,γ) activated I^{128} with CH_4 also enable them to study indirectly interactions of other molecules with atoms activated by neutron processes (18).

Willard and coworkers also studied the reactions activated by the Br^{80m} isomeric transitions (19). They found that the yields are lower than that from the (n,γ) process.

2. Carbon, C^{11} and C^{14}

C. Mackay and coworkers (20) reported the study of reactions of carbon atoms with a number of gaseous molecules using 20.5-min. C^{11} produced by a neutron stripping reaction of C^{12} . With simple oxidizing agents (e.g., air, O_2 , O_2-O_3 mixtures, CO, CO_2 , NO, NO_2 , N_2O and SO_2) carbon monoxide is always the major product. They explained this in terms of a third body restriction on the further oxidation to CO_2 . With hydrogen, the recoil carbon atom gives half methane and half CO although the only sources of oxygen are trace impurities and the wall. With hydrocarbons the most important single product in all systems is acetylene, amounting to 30-35%. From the results they were able to point to the existence of several reaction mechanisms, involving C-H, C-C bonds.

Yang and Wolf studied the $N^{14}(n,p)C^{14}$ reaction in anhydrous ammonia in the gas phase (21). A uniquely high specific yield of methane- C^{14} has been observed. This high yield of methane- C^{14} as essentially the sole significant product from operations of the recoil carbon in gaseous ammonia was not

affected by the addition of inert moderating gases. The presence of oxygen as radical scavenger did not produce any significant effects. Reaction of the recoil carbon in the presence of methane and methylamine, however, gave a complex mixture of radioactive products, which is common to all carbon-14 systems. Yang and Wolf proposed a process of hydrogen abstraction to account for the apparently unique result from the recoil carbon in the presence of ammonia in the gas phase. They also used a collision complex mechanism to account for the nature of the recoil fragments.

3. Tritium

Hot-atom chemistry of tritium in the gas phase is probably the most extensively studied branch in this field(22,23,24,25). Hot displacement was observed in alkanes in which H, 2H, or an alkyl group is displaced. The types of reactions are:


- A. Hydrogen atom abstraction. This reaction, the only one observed at thermal energies, also occurs with hot hydrogen (although with a much higher collision yield)



- B. Hydrogen atom displacement



- C. Replacement of alkyl group to form a degraded alkane



- D. Replacement of two groups to form a radical. Either two hydrogen atoms or a hydrogen atom and an alkyl group may be displaced by the hot atom.



The resulting radical may combine with iodine scavenger to form alkyl iodide. Alternatively, it may decompose into a smaller radical and an alkene by breaking a C-C bond with rearrangement,



The reactions initiated by beta decay of tritium forming tritiated hydrocarbons in systems initially composed of gaseous propane and tritium are also reported (26).

4. Lead-212 from α -decay

The observations that recoil tritium and halogen atoms undergo substitution reactions in the gas phase have led to an investigation with recoil metal atoms to see whether similar reactions occur. Accordingly, the gas phase reactions of lead-212 recoiling from alpha decay of polonium-216 have been studied (27,28).

These studies have indicated that the Pb^{212} produced by alpha decay in the presence of such gases as CH_4 , Ar, and CO is retained in the gas phase to a large extent and may be transferred as a vapor using standard vacuum line techniques. Further studies regarding the method by which the recoil lead is retained in the vapor phase are still in progress. At the present time, it seems quite likely that aerosol formation may be at least partly responsible for the observed results.

IV. CONCLUSION

The chemical effect of nuclear transformations, or hot atom chemistry is a branch of radiochemistry that is relatively new and not well understood. Although much progress has occurred since observations by Szilard and Chalmers in 1934 (29) opened the field, more effort should be made to explore this area.

LITERATURE CITED

1. J. E. Willard, Ann. Rev. Nuclear Sci., 3, 193 (1953); Ann. Rev. Phys. Chem., 6, 141 (1955).
2. A. C. Mahl and N. A. Bonner, "Radioactivity Applied to Chemistry," John Wiley and Sons, Inc., New York, 1951, Chapter 8.
3. D. L. Baulch and J. F. Duncan, Quart. Revs., 12, 133 (1958).
4. Papers in Proceedings of IAEA Symposium on Chemical Effects of Nuclear Transformations, Prague, 1960, published by International Atomic Energy Agency, Vienna, 1961.
5. S. C. Lind, "Radiation Chemistry of Gases," Reinhold Publishing Corp., New York, 1961, Chapter 11.
6. Section on theoretical aspects in reference 4.
7. L. Friedman and W. F. Libby, J. Chem. Phys., 17, 647, (1949); M. S. Fox and W. F. Libby, J. Chem. Phys., 20, 487 (1952).
8. F. J. Estrup and R. Wolfgang, J. Am. Chem. Soc., 82, 2661, 2665 (1960).

9. A. Sokolowska, L. A. Haskin, and F. S. Rowland, J. Am. Chem. Soc., 84, 2469 (1962).
10. C. H. Hsiung and A. A. Gordus, Unpublished results, Ph. D. thesis of C. H. Hsiung, University of Michigan, 1962.
11. C. H. Hsiung and A. A. Gordus, J. Chem. Phys., 36, 947 (1962).
12. A. A. Gordus and C. H. Hsiung, J. Chem. Phys., 36, 954 (1962).
13. J. C. W. Chieu and J. E. Willard, J. Am. Chem. Soc., 75, 6160 (1953).
14. A. A. Gordus and J. E. Willard, J. Am. Chem. Soc., 19, 4609 (1957).
15. J. F. Horning, G. Levey, and J. E. Willard, J. Chem. Phys., 20, 1556 (1952); G. Levey and J. E. Willard, J. Chem. Phys., 25, 904 (1956).
16. E. P. Mack and A. A. Gordus, J. Phys. Chem., 65, 944 (1961).
17. E. P. Mack, and A. A. Gordus, J. Chem. Phys., 34, 1855 (1961).
18. E. P. Mack and A. A. Gordus, J. Chem. Phys., 36, 287 (1962).
19. Ref. 14, also J. E. Willard, J. Am. Chem. Soc., 62, 3161(1940).
20. C. Mackay, M. Pandow, P. Polark, and R. Wolfgang, Proceedings of IAEA symposium on Chemical Effects of Nuclear Transformations Vol. II. P. 1).
21. J. Y. Yang and A. P. Wolf, J. Am. Chem. Soc., 82, 4488 (1960).
22. F. S. Rowland, J. K. Lee, B. Musgrave and R. M. White, Proc. IAEA Symposium on Chemical Effects of Nuclear Transformations, Prague, Oct. 1960, Published by Internal Atomic Energy Agency, Vienna, 1961, Vol. 2, p. 6).
23. M. Henchman, D. Urch, and R. Wolfgang ibid., p. 83.
24. J. K. Lee, B. Musgrave, and F. S. Rowland, J. Am. Chem. Soc., 82, 3545 (1960).
25. D. Urch and R. Wolfgang, J. Am. Chem. Soc., 83, 2981 (1961).
26. K. Yang and P. L. Gant, J. Phys. Chem., 66, 1619 (1962).
27. J. G. Kay and F. S. Rowland, Paper presented before the Division of Physical Chemistry at the 139th Meeting of the American Chemical Society, St. Louis, Missouri, March, 1961.
28. J. G. Kay and A. C. Wang, unpublished results.
29. L. Szilard and T. A. Chalmers, Nature, 134, 462 (1934).

RECENT CHEMISTRY OF THE ALANES

D. P. Eyman

January 15, 1963


I. INTRODUCTION

The chemistry of the alanes [alane is the name proposed by Wiberg for aluminum hydride and its derivatives (1)] is dominated by a highly acidic character and the tendency of the aluminum to attain tetra coordination. These characteristics are demonstrated by the common occurrence of associated alanes (2,3); amine and ether adducts of alanes, which are more stable than analogous borane adducts (4); and anionic complexes with alkali metal halides and alkyls. The strength of the aluminum-nitrogen coordinate bond in amine alane adducts is such that with the correct substituents on nitrogen and aluminum condensation to amino alanes and imino alanes is possible (5).

II. AUTOCOMPLEXATION

The large tendency for alanes to associate comes from the electron deficiency of aluminum compounds with a coordination number of three. Association leads to dimers, trimers, and polymers with atoms or groups as bridges bound by three-center bonds.

The structure of dimeric trimethylalane, as established by X-ray crystal analysis, involves bridging methyl groups (6). It has been shown that in higher trialkylalanes the α carbon of the alkyl group forms a bridge between aluminum atoms (7). The dimers have a distinct tendency to dissociate to monomers in the vapor phase at temperatures up to 160°C. The degree of association of trialkylalanes decreases in the order (8)



Triisopropylalane, triisobutylalane, and all alanes of the types $\text{Al}(-\text{CH}_2\text{CHR}'_2)_3$ and $\text{Al}(-\text{CH}_2\text{CR}'_3)_3$ are monomeric, as determined cryoscopically in benzene (2). In general the tendency to dimerize diminishes with increasing alkyl chain length or increasing alkyl complexity. This behavior is probably the result of steric hindrance (9).

Dialkylalanes are trimeric according to cryoscopic measurements in benzene (10). There is no difference among the methyl, ethyl, and isobutyl compounds. Infrared studies indicate that association occurs by way of Al-H-Al bonds. The heat of association is estimated to be 15-20 kcal. per hydrogen bridge (11).

Alanes of the type R_2AlX (R=alkyl; X=OR, NR₂, PET₂, Cl) are associated through Al-X-Al bridges. The dimers and trimers

are known in (9,12). $(\text{R}_2\text{AlCl})_2$ $(\text{Et}_2\text{AlOEt})_2$ $(\text{Et}_2\text{AlNEt}_2)_2$ $(\text{Et}_2\text{AlOMe})_2$ $(\text{Et}_2\text{AlPMe}_2)_2$

Deaggregation of some of these polymers can be effected by addition of electron donors. The compounds $\text{Me}_2\text{NAlCl}_2$, $\text{Me}_2\text{NAl}(\text{C}_6\text{H}_5)_2$, and $\text{Me}_2\text{NAl}(\text{C}_4\text{H}_9)_2$ are inert to the attack of bases (13).

Alane "inner complexes" of the type $\text{R}_2\text{Al}(\text{CH}_2)_3\text{Y}$ (R=alkyl, aryl; Y=OEt, SEt, OPr, NEt₂), as well as $\text{R}_2\text{Al}(\text{CH}_2)_4\text{OEt}$ are known (14,15,16). The thermal condensation of triphenylalane and acetylacetone produces diphenylalane acetylacetonate, which has been assigned a chelate structure (17).

III. MOLECULAR COMPLEXES

Alane and trialkylalanes behave as very strong acceptors and form complexes with ethers and amines which are more stable than the analogous borane complexes. The following order of donor character towards trimethylalane has been established (4,18).




Wiberg (19) has reported that triethylamine alane is largely dimeric in benzene whereas tripropylamine alane is only slightly dimerized and tributylamine alane is monomeric. All of these are monomeric in diethyl ether. The values for the degree of association (\underline{n}) of some amine alane adducts in benzene are listed in Table I (20).

Table I
Degree of Association for Amine Alane Adducts

<u>Compound</u>	<u>n</u>	<u>Compound</u>	<u>n</u>
$\text{Me}_3\text{N}:\text{AlH}_3$	1.40	$\text{Me}_3\text{N}:\text{AlHCl}_2$	1.46
$\text{AllylMe}_2\text{N}:\text{AlH}_3$	1.37	$\text{Me}_3\text{N}:\text{AlCl}_3$	1.36
$\text{Et}_2\text{MeN}:\text{AlH}_3$	1.33	$\text{Et}_3\text{N}:\text{AlH}_3$	1.10
$\text{Me}_3\text{N}:\text{AlH}_2\text{Cl}$	1.32	$\text{Pr}_3\text{N}:\text{AlH}_3$	1.05


Association is believed to occur through hydrogen bridges or through dipole-dipole interaction in solution (13).

Tensiometric titrations by Ruff (21) confirmed an earlier report of penta coordinated aluminum in bis-trimethylamine alane (22). These studies also demonstrated the existence of bis-complexes for Me_2NAllyl and Et_2NMe , but not for Et_3N and Pr_3N . The order of stabilities for bis-trialkylamine alane complexes is



Dipole moment studies (11) and single crystal X-ray analysis (23) both indicate that bis-trimethylamine alane is trigonal bipyramidal.

A novel adduct of alane and N,N,N',N' -tetramethylethylenediamine is reported to be a chelated dimer having the structure (24)




Dautel and Zeil (25) have reported an alane dioxane 1:1 adduct which behaves as a polymer. Apparently aluminum is penta-coordinate, and the dioxane is behaving as a bridging bifunctional ligand. Fetter and co-workers (26,27,28) recently have reported 1:1 trimethylalane adducts of trimethylhydrazine and tetramethyltetrazine. Proton resonance studies of the trimethylalane tetramethyltetrazine complex indicated that the tetramethyltetrazine was acting as a bidentate ligand.

Molecular complexes of alanes with other organo-metallic compounds are known. These are non-electrolytes and presumably electron deficient compounds. Among such complexes are LiAlEt_4 , EtMgAlEt_4 , and $\text{Mg}(\text{AlEt}_4)_2$ (29). Ruff (30) has recently prepared a 1:2 adduct of bis-dimethylamino alane and methylamino borane.

IV. ANIONIC COMPLEXES

Anionic complexes of the alanes are usually obtained by adding certain alkali metal compounds (MX), especially salts, to aluminum compounds of the general formula $\text{AlR}_3 - \text{X}_n$ ($n=2$ to 0). Typical compounds for addition are: alkali halides, pseudohalides, hydrides, alkyls, aryls, and alkoxyls. The stabilities of these anionic complexes follow certain rules. Large alkali metal ions and small halide ions produce the most stable metal halide alane complexes. The stability decreases with increasing size of the alkyl attached to aluminum: all trialkylalanes from methyl to butyl add NaF; $\text{Al}(\text{C}_6\text{H}_{13})_3$ does not (31). The stability of tetraethyl aluminate is demonstrated by the fact that the ethide ion displaces diethyl ether from the triethylalane diethyl ether adduct (32).

Bonitz (33) attributed the appreciable conductance of triethylalane to self-ionization of the type



In addition to the type MAlR_4 , there exists the 1:2 type complexes in which two molecules of the alane are added to the alkali metal compound (34). The 1:2 complex, $\text{NaF}:2\text{AlEt}_3$, is a liquid at room temperature and is a much better electrical conductor than $\text{NaF}:\text{AlEt}_3$ (35).

V. CONDENSATION REACTIONS AND ALUMINUM POLYMERS

Alane addition compounds of ammonia, primary amines, and secondary amines normally do not exist at room temperature due to an internal stabilization by splitting off RH . This condensation leaves an amino alane. The N-H and Al-R bonds are replaced by additional Al-N bonds to maintain tetra coordination for both aluminum and nitrogen (36). This leads to polymeric structures which are cyclic. If the amino alane has both N-H and Al-R bonds, it undergoes a second condensation step when the temperature is sufficiently high. This step produces insoluble, non-volatile, cross-linked, highly polymeric frameworks (37).

Wiberg and Gosele (38) reported that 1:1 and 1:2 alane pyridine adducts undergo an exothermic hydrogen migration at elevated temperatures. In the presence of excess pyridine, the product is tris-pyridino alane.

From the reaction of alane with excess ammonia, Wiberg has isolated, at different temperatures, $\text{AlH}_3:\text{NH}_3$, AlH_2NH_2 , $\text{AlH}(\text{NH}_2)_2$, and $\text{Al}(\text{NH}_2)_3$ (39). Stepwise thermal decomposition of $\text{AlH}_3:\text{NH}_3$ gave AlH_2NH_2 , $(\text{AlHNH})_x$, and AlN (40). By means of the same procedure, the analogous derivatives of primary amines (41) and secondary amines (42) were prepared.

Ruff (43) has prepared bis- and tris-dialkylamino alanes by the reaction of trialkylamine alane adducts with dialkylamines. As discussed in section II, dialkylamino alanes are usually dimers or trimers. The compounds $(\text{Me}_2\text{N})_3\text{Al}$ and $(\text{Me}_2\text{N})_2\text{AlH}$ are associated in benzene, but $(i\text{-Pr}_2\text{N})_3\text{Al}$ is monomeric. Longi and co-workers (44) have prepared $(\text{Me}_2\text{N})_3\text{Al}$ and $[(\text{C}_6\text{H}_5)_2\text{N}]_3\text{Al}$ from the appropriate secondary amine and $(\text{Me}_2\text{CHCH}_2)_3\text{Al}$.

Nielsen and Sisler (45) reported the preparation of diethyl-2,2-dimethylhydrazino alane but did not isolate an adduct of triethylalane and N,N -dimethylhydrazine. Infrared data favor a six-membered ring structure. When heated,

the hydrazino alane condensed to a polymer having the composition $(\text{MeH}_2\text{NNAEt})_x$. Laubengayer and co-workers (36) have allowed the adduct $\text{MeH}_2\text{H}:\text{AlEt}_2\text{Cl}$ to undergo successive condensation steps. The first condensation, at 50°C ., yields $(\text{MeHNAEtCl})_x$ and C_2H_6 . The second step, at 210°C ., yields $(\text{MeNAlCl})_x$, which is a high polymer. The adducts $\text{MeH}_2\text{N}:\text{AlEtCl}_2$ and $\text{Me}_2\text{HN}:\text{AlEt}_2\text{Cl}$ undergo only one condensation step. These same workers have prepared the imino alane $(\text{MeNAlC}_6\text{H}_5)_x$, as the product of two successive condensation steps, from the adduct $\text{MeH}_2\text{N}:\text{Al}(\text{C}_6\text{H}_5)_3$. This imino alane and $(\text{MeNAlCl})_x$ are both stable at 350°C . in a nitrogen atmosphere. However, both are hydrolytically unstable.

Fetter and his associates (26,27,28) have prepared polymers by the reactions of trimethylalane and substituted hydrazines and tetrazines. A polymer, having the composition $(\text{MeAlNHNMe})_x$, prepared by the reaction of trimethylalane with methylhydrazine, decomposed at 200°C . Another polymer, $[\text{Al}(\text{MeNNMe})_{1.5}]_x$, was prepared by the reaction of trimethylalane with 1, 2-dimethylhydrazine. Both of these polymers are hydrolytically unstable.

VI. CONCLUSIONS

Much of the recent published research dealing with alane chemistry is concerned with the alane amine adducts and their condensation products. Most of these investigations are designed to elucidate the character of the aluminum-nitrogen bond and the effects upon this bond of substitution on aluminum or nitrogen. The thermal stabilities of the aluminum-nitrogen polymers prepared thus far are quite encouraging. The major problem for future workers in this field is the preparation of a hydrolytically stable aluminum-nitrogen polymer.

REFERENCES

1. Wiberg, E., *Angew. Chem.*, 65, 16 (1953).
2. Zeiss, H., *Organometallic Compounds*, p.207, Reinhold Publishing Co., New York (1960).
3. Coates, G. E., *Organo-Metallic Compounds*, p.126, John Wiley & Sons, Inc., New York (1960).
4. Stone, F. G. A., *Chem. Revs.*, 58, 101 (1958).
5. Bähr, G., *F.I.A.T. Reviews of German Science, 1939-1946, Inorganic Chemistry, Part II*, p.157.
6. Lewis, P. H., and Rundle, R. E., *J. Chem. Phys.*, 21, 986 (1953).
7. Rundle, R. E., *J. Phys. Chem.*, 61, 45 (1957).
8. Eisch, J., and Gilman, H., *Adv. Inorg. Chem. and Radiochem.*, 2, 61 (1960).
9. Hoffman, E. G., *Ann.*, 629, 104 (1960).
10. Wiberg, E., *F.I.A.T. Reviews of German Science, 1939-1946, Inorganic Chemistry, Part I*, p.148.
11. Schomberg, G., and Hoffmann, E. G., *Z. Elektrochem.*, 61, 1110 (1957).
12. Davidson, N., and Brown, H. C., *J. Am. Chem. Soc.*, 64, 316 (1942).
13. Ruff, J. K., *J. Am. Chem. Soc.*, 83, 1798 (1961).
14. Bähr, G., and Müller, G. E., *Chem. Ber.*, 88, 251 (1955).
15. Zakharkin, L. I., and Savina, L. A., *Izvest. Akad. Nauk S.S.S.R., Otdel, Khim. Nauk*, 444 (1959); *C. A.*, 53, 2162d (1959).
16. Zakharkin, L. I., and Savina, L. A., *Izvest. Akad. Nauk S.S.S.R., Otdel, Khim. Nauk*, 1039 (1960); *C. A.* 54, 24345i (1960).

17. Costa, G., and Calcinari, R., *Gazz chim. ital.*, 89, 1415 (1959); *C. A.*, 54, 22006i (1960).
18. Wiberg, E., May, A., and Nöth, H. *Z. Naturforsch.*, 10b, 239 (1955).
19. Wiberg, E., *Z. Naturforsch.*, 10b, 238 (1955).
20. Ruff, J. K., and Hawthorne, M. F., *J. Am. Chem. Soc.*, 82, 2141 (1960).
21. *Ibid.*, 83, 535 (1961).
22. Wiberg, E., *Z. anorg. u. allgem. Chem.*, 272, 221 (1958).
23. Parry, R. W. Nordman, C. E., and Heitsch, C. W., Abstracts 142nd Meeting A. C. S., Sept., 1962, p.21N.
24. Davidson, J. M., and Wartik, T., *J. Am. Chem. Soc.*, 82, 5506 (1960).
25. Dautel, R., and Zeil, W., *Z. Elektrochem.*, 64, 1234 (1960).
26. Fetter, N. R., and Bartocha, B., *Can. J. Chem.*, 39, 2001 (1961).
27. *Ibid.*, 40, 342 (1962).
28. Fetter, N. R., Brinckman, F. E., and Moore, D. W., *Can. J. Chem.*, 40, 2184 (1962).
29. Ziegler, K., and Holzkamp, E., *Ann.*, 605, 93 (1957).
30. Ruff, J. K., *Inorg. Chem.*, 1, 612 (1962).
31. Ziegler, K., Köster, R., Lehmkuhl, H., and Reinert, K., *Ann.*, 629, 33 (1960).
32. Baker, E. B., and Sisler, H. H., *J. Am. Chem. Soc.*, 75, 5193 (1953).
33. Bonitz, E., *Chem. Ber.*, 88, 742 (1955).
34. Schultz, R. F., Metal Organic Compounds, *Advances in Chemistry Series #23*, p.163, American Chemical Society, Washington, D. C. (1959).
35. Ziegler, K., International Conference on Coordination Chemistry, Special Publication No. 13, The Chemical Society, London (1961).
36. Laubengayer, A. W., Smith, J. D., and Ehrlich, G. G., *J. Am. Chem. Soc.*, 83, 542 (1961).
37. Laubengayer, A. W., Inorganic Polymers, Special Publication No. 15, The Chemical Society, London (1961).
38. Wiberg, E., and Gösele, W., *Z. Naturforsch.*, 10b, 236 (1955).
39. Wiberg, E., and May, A., *Z. Naturforsch.*, 10b, 230 (1955).
40. *Ibid.*, p.229.
41. *Ibid.*, p.232.
42. *Ibid.*, p.234.
43. Ruff, J. K., *J. Am. Chem. Soc.*, 83, 2835 (1961).
44. Longi, P., Mazzanti, G., and Bernardini, F., *Gazz. chim. ital.*, 90, 180 (1960); *C. A.*, 65, 11339d (1961).
45. Nielsen, R. P., and Sisler, H. H., *J. Chem. Soc.*, 3818, 1962.
46. Laubengayer, A. W., Wade, K., and Lengnick, G., *Inorg. Chem.*, 1, 632 (1962).

ROLL CALL

RATE OF A SUBSTITUTION REACTION IN CHLOROFORM

Ref: S. D. Ross and M. Finkelstein, J. Am. Chem. Soc.
79, 6547 (1957)

George Shier

February 19, 1963

A detailed study of the rate of reaction of 2,4-dinitrochlorobenzene with n-butyl ammine showed that the rate constant was not constant for different initial concentrations of ammine. The reaction showed a positive neutral salt effect but no common ion effect with n-butylammonium salts. It was found that the reaction obeyed the rate law $K_2 = K_1 + K_7(n\text{BaNH}_2)$ where the K_1 term represents a path first order in both ammine and chloride and K_7 represents a path first order in chloride and second order in ammine. The mechanistic implications were discussed.

ROLL CALL

INFRARED STUDIES OF TRIMETHYLTIN COMPOUNDS

R. C. Burrows

February 19, 1963

In an attempt to determine whether the free trimethyltin cation exists in the solid state, Okawara, Hathaway, and Webster(1) have examined the infrared spectra of the compounds $(\text{CH}_3)_3\text{SnX}$, where X represents ClO_4 , NO_3 and $(\text{CO}_3)_{1/2}$. By comparing the number of bands predicted for various symmetry species and the number found for the trimethyltin and anion residues, they have shown that the cation $(\text{CH}_3)_3\text{Sn}^+$ is not present as such in these compounds, but rather that this group is covalently bound to the anion. The perchlorate is thought to have a polymeric structure involving five-coordinate tin, planar trimethyltin residues, and bidentate perchlorate groups. Although the nitrate and carbonate groups are probably also bidentate in their respective compounds, a non-planar trimethyltin residue was inferred here since both the symmetric and the antisymmetric C-Sn stretches are infrared active. Further structural details could not be derived for these two compounds on the basis of the infrared evidence alone.

An investigation(2) of trimethyltin tetrafluoroborate was also conducted by these workers along similar lines. The infrared spectrum of this compound corresponded to that which is theoretically predicted for a planar trimethyltin residue and a tetrafluoroborate group of C_{2v} symmetry. From these results the structure was assumed to be analogous to that deduced for trimethyltin perchlorate with the tetrafluoroborate group, generally considered to be a non-complexing anion, behaving as a bidentate ligand, as shown.



References

1. R. Okawara, B. J. Hathaway and D. E. Webster, Proc. Chem. Soc., 1963, 13.
2. B. J. Hathaway and D. E. Webster, Proc. Chem. Soc., 1963, 14.

ROLL CALL

BRIDGE ADDUCTS-THE INTERACTION OF BORON TRIFLUORIDE
WITH TRANSITION-METAL CYANIDE COMPLEXES

Gerald Feistel

February 19, 1963

A new class of compounds containing the $M-C\equiv N-BF_3$ linkage has been synthesized. The compounds, $K_2Ni(CN)_4 \cdot BF_3$, $K_2Ni(CN)_4 \cdot 4BF_3$, $K_4Mo(CN)_8 \cdot 7.3 BF_3$, and $K_4Fe(CN)_6 \cdot 3.7 BF_3$ have been formed by the reaction of BF_3 gas with the solid metal cyanide complexes at various temperatures and pressures.

A shift in the infrared band of the cyanide group and the appearance of new bands indicate the formation of the adduct. There is no change in the magnetism of the compounds of the Ni cyanides. Lack of simple metal cyanide- BF_3 ratios is attributed to a kinetic effect.

1. D. F. Shriver, J. Am. Chem. Soc., 84, 4610 (1962).

N.M.R. COUPLING CONSTANTS AND HYBRIDIZATION IN ORGANO-TIN COMPOUNDS

J. C. Schapiro

March 5, 1963

A relationship between ^{13}C -H nuclear magnetic resonance coupling constants and bonding hybrid orbitals has been shown to exist in a series of hydrocarbons (1, 2). The development of this relationship and its application to the bonding in some organo-tin compounds will be considered in the following sections.

THEORY

The Hamiltonian for a molecular system in a magnetic field may be expressed as the sum of four principal parts,

$$H = H_1 + H_2 + H_3 + H_4$$

The term H_4 is for the direct magnetic interaction of the nuclei with each other. This term may be neglected since it averages to zero in nuclear magnetic resonance experiments in which frequent collisions average the molecule over all orientations(3).

The term H_1 includes terms for the kinetic energy of the electrons, their interaction as moving charged particles in the magnetic field of the nuclei, electrostatic potential energy, and the electron orbital-orbital, spin-orbital, and spin-spin interactions(4). H_2 is the term for the dipole-dipole interactions between the nuclear magnetic moments and the electron magnetic moments. H_3 is the contact term and is dependent upon the properties of electrons at the nucleus. The electron-coupled nuclear spin-spin interactions are those terms dependent on the spins of two interacting nuclei which arise when H_1 , H_2 , and H_3 are discussed to the accuracy of second-order perturbation theory.

Ramsey(3) and Karplus(5) have shown that in cases in which the interacting nuclei are separated by one or two bonds, the largest and dominant contribution to the nuclear spin-spin interaction comes from the mechanism which corresponds to second order perturbations involving H_3 . These terms tend to be largest because they correspond to the electrons being close to the nuclei. The second-order perturbation energy for the operator H_3 is

$$E_{(1)NN'}^{(2)} = - \sum_{n(\neq 0)} \frac{\langle 0 | H_3 | n \rangle \langle n | H_3 | 0 \rangle}{E_n - E_0} ; \quad H_3 = \frac{16\pi\beta h}{3} \sum_k \sum_N \gamma_N \delta(\vec{r}_{kN}) \vec{S}_k \cdot \vec{I}_N$$

where: $E_n - E_0$ = the excitation energy between the singlet ground electronic state and the nth triplet excited state.

β = Bohr magneton = $e\hbar/2mc$

γ_N = magnetogyric ratio of nucleus N

$\vec{r}_{kN} = \vec{r}_k - \vec{r}_N$

$\delta_{\vec{r}_{kN}}$ = Dirac function which picks out the value at $\vec{r}_{kN} = 0$ in any integration over the coordinates of electron k.

\vec{S}_k = spin of electron k

\vec{I}_N = spin of nucleus N

The coupling constant $J_{NN'}$, (in units of cycles per second) is defined by

$$J_{NN'} = \frac{1}{h} J_{NN'}^{(3)} \frac{1}{\Delta E}$$

where $E_{NN'}$ is the energy of interaction between two nuclei with spins I_N and $I_{N'}$ (units of \hbar) (6). If one uses the energy of interaction contributed by the $1/r^3$ term only, then the above equation defines $J_{NN'}^{(3)}$, the coupling constant which arises from the contact term alone. It can be shown that

$$J_{NN'}^{(3)} = - \frac{2}{3h} \left(\frac{16\pi\beta\hbar^2}{3} \right)^2 \chi_N \chi_{N'} \frac{1}{\Delta E} \times \left[\langle 0 | \sum_{k,j} \delta(\vec{r}_{kN}) \delta(\vec{r}_{jN'}) \vec{S}_k \cdot \vec{S}_j | 0 \rangle \right]$$

In deriving the above equation the singlet-triplet excitation energies are approximated by a mean value ΔE . The limits of this approximation have been discussed by Ripka (7). Evaluation of $J_{NN'}^{(3)}$ then depends upon the wave function, ψ_0 , for the ground electronic state.

Using a generalized valence-bond function

$$\psi_0(\mathbf{r}) = \eta(\lambda_N, \lambda_{N'}) \left\{ \phi_N(\mathbf{r}) \phi_{N'}(\mathbf{r}) + \phi_N(\mathbf{r}) \phi_{N'}(\mathbf{r}) + \lambda_N \phi_N(\mathbf{r}) \phi_{N'}(\mathbf{r}) + \lambda_{N'} \phi_{N'}(\mathbf{r}) \phi_N(\mathbf{r}) \right\}$$

where ϕ_N and $\phi_{N'}$ are hybrid orbitals on nuclei N and N' respectively, $\lambda_N, \lambda_{N'}$ are parameters determining the charge distribution and $\eta(\lambda_N, \lambda_{N'})$ is the appropriate normalizing factor with inclusion of overlap, $J_{NN'}^{(3)}$ becomes a function of the degree of hybridization in the atomic orbitals $\phi_N, \phi_{N'}$ and the bond polarizations $\lambda_N, \lambda_{N'}$ (1). The term $J_{NN'}^{(3)}$ is dependent upon the amount of s character in the hybrid atomic orbital, since the s orbital does not have a nodal plane at the nucleus.

Two additional points need to be considered in choosing the wave function for $J_{NN'}^{(3)}$. First if the two interacting nuclei are directly bonded NN' to each other, only the perfect pairing structure contribution to the ground-state wave function need be taken into account; if the two nuclei are not joined through a single sigma bond, the non-perfect pairing structures contribute to make $J_{NN'}^{(3)}$ non-zero. Secondly, Muller and Pritchard(2) and Juan and Gutowsky (8) have shown that J_{CH} in the C^{13} -H group is relatively insensitive to the value of λ_{C-H} (ionic character of the bonds) within certain limits. However, J_{Si-H} for the ^{29}Si -H group has been shown to be dependent upon λ_{Si-H} due to the greater polarizability of Si as compared to C(8). It might be expected then that ionic contribution to the wave function would be important in the case of the other group IVB elements.

APPLICATION OF THE THEORY

For any group IVB element, M , the Fermi contact contribution to the coupling constant J_{M-H} , using a valence bond wave function for the perfect pairing structure, may be expressed as

$$J_{M-H} = \frac{\chi_M \chi_H}{h \Delta E} \left(\frac{16\pi\beta\hbar^2}{3} \right)^2 \eta^2 \alpha^2 |1s_M(0)|^2 |1s_H(0)|^2$$

where α^2 is the s character of the M hybrid for the M-H bond. (8) $n_s(0)$ is the n_s wave function evaluated at the M nucleus. If non-perfect pairing structures are to be considered, the expression becomes more complicated but it is still linearly dependent on the amount of s character in the hybrid orbital on M. Therefore, one may to a first approximation determine changes in hybridization in a given series of compounds by observing trends in the coupling constants for those compounds.

METHYLTIN HALIDES

Natural abundance tin contains three nuclides with spin of $1/2$, ^{115}Sn , ^{117}Sn , and ^{119}Sn . The abundances are ^{115}Sn , 1%, ^{117}Sn , 7.67%, and ^{119}Sn , 8.67%. Spin-spin couplings due to ^{115}Sn are not observed experimentally. Holmes and Kaesz (9) and Van der Kelen (10) have studied the tin-hydrogen splitting on the proton resonance in a series of methyl tin halides. Holmes and Kaesz indicate that the contact mechanism makes the dominant contribution to Sn-C-H coupling and that except for rehybridization of the tin atom, other components of the bonding system Sn-C-H do not change appreciably. They point out that the assumption receives some support from the fact that the ^{13}C -H spin-spin interactions vary by a very small amount in this series of compounds. As a Cl atom is substituted for a CH_3 group the coupling constant $J_{\text{Sn-C-H}}$ increases indicating a larger amount of s character in Sn-C-H the tin bonding orbital towards carbon; this following Bent's argument for "second order hybridization" effects (11, 12).

In H_2O and D_2O solutions the formation of ions such as Me_3Sn^+ , $\text{Me}_2\text{Sn}^{2+}$ and MeSn^{3+} indicates entirely different hybridization states for the tin atom. Abrupt changes in $J_{\text{Sn-C-H}}$ are observed similar to those observed for $J_{\text{C-H}}$ in carbon compounds where hybridization changes from sp^3 - sp^2 to sp (10). Burke and Lauterbur (13), observing proton coupling on the ^{119}Sn resonance, found $J_{\text{Sn-C-H}}$ for $(\text{CH}_3)_2\text{SnCl}_2$ to be 80 c.p.s. in acetone and 98 c.p.s. $J_{\text{Sn-C-H}}$ in H_2O , which they attribute to the existence of alkyl-tin-chloride-solvent complexes. Although the $(\text{CH}_3)_3\text{SnBr}\cdot\text{py}$ complex is very stable, the observed change in $J_{\text{Sn-C-H}}$ is rather small; Van der Kelen attributes the small $J_{\text{Sn-C-H}}$ change in J to a small change in the hybridizational state of tin (10). In light of the recent work by Matwiyoff (14), who has found that the Sn-H coupling constants in $(\text{CH}_3)_3\text{SnCl}$ increase about 10 c.p.s. when measured in TMSO, it appears that the small change in $J_{\text{Sn-CH}_3}$ found for the pyridine complex in CDCl_3 solution may $J_{\text{Sn-CH}_3}$ be attributed to the diluteness of the solution and perhaps dissociation of the $(\text{CH}_3)_3\text{SnBr}\cdot\text{py}$ complex.

OTHER ORGANO-TIN SYSTEMS

Recently, Flitcroft and Kaesz (15) have observed in the SnH_4 , methylstannane series, trends similar to those in the methyltin halide series that are similarly explained by a combination of the theory of spin-spin coupling and the effects of the electronegativity of a substituent on the hybridization

TABLE 1 - COUPLING CONSTANTS IN METHYLTIN HALIDES (in c.p.s.)

<u>COMPOUND</u>	<u>SOLVENT</u>	$J_{C^{13}-H}$	$J_{Sn-CH_3}^{117}$	$J_{Sn-CH_3}^{119}$	<u>REF.</u>
$(CH_3)_4Sn$	neat	-	51.5	54.0	9
	CCl_4	-	51.4	53.8	9
	$CDCl_3$	128.4	52	54.3	10
$(CH_3)_3SnCl$	neat	-	57.4	59.7	9
	CCl_4	-	56.0	58.9	9
	$CDCl_3$	133	56.4	58.6	10
	H_2O	-	65.2	68.4	9
	D_2O	134	67.7	70.7	10
$(CH_3)_2SnCl_2$	neat:	-	68.0	71.0	9
	CCl_4	-	66.6	69.7	9
	$CDCl_3$	137.8	67	69.5	10
	H_2O	-	97.4	101.9	9
	D_2O	134	102.5	107.5	10
CH_3SnCl_3	neat:	-	95.7	100	9
	CCl_4	-	95.3	99.5	9
	$CDCl_3$	-	93.4	98.5	10
	H_2O	-	125.4	131.1	9
$(CH_3)_3SnBr$	$CDCl_3$	134	56	58.6	10
$(CH_3)_3SnBr \cdot py$	* $CDCl_3$	134	59.8	62.1	10
$(CH_3)_3SnCl$	* $TMSO$		68.0	70.0	14

* Probably a dilute solution

** $TMSO$ = tetramethylene sulfoxide

Table 1. Summary of the data for the different groups of patients.

Group	Number of patients	Mean age (years)	Mean duration (years)	Mean duration (months)	Mean duration (days)
1	10	55.2	12.5	150	4500
2	15	58.1	14.2	170	5130
3	20	60.3	16.8	200	6060
4	25	62.5	19.1	230	6975
5	30	64.7	21.5	260	7800
6	35	66.9	24.0	290	8700
7	40	69.1	26.5	320	9600
8	45	71.3	29.0	350	10500
9	50	73.5	31.5	380	11400
10	55	75.7	34.0	410	12300
11	60	77.9	36.5	440	13200
12	65	80.1	39.0	470	14100
13	70	82.3	41.5	500	15000
14	75	84.5	44.0	530	15900
15	80	86.7	46.5	560	16800
16	85	88.9	49.0	590	17700
17	90	91.1	51.5	620	18600
18	95	93.3	54.0	650	19500
19	100	95.5	56.5	680	20400
20	105	97.7	59.0	710	21300
21	110	99.9	61.5	740	22200
22	115	102.1	64.0	770	23100
23	120	104.3	66.5	800	24000
24	125	106.5	69.0	830	24900
25	130	108.7	71.5	860	25800
26	135	110.9	74.0	890	26700
27	140	113.1	76.5	920	27600
28	145	115.3	79.0	950	28500
29	150	117.5	81.5	980	29400
30	155	119.7	84.0	1010	30300
31	160	121.9	86.5	1040	31200
32	165	124.1	89.0	1070	32100
33	170	126.3	91.5	1100	33000
34	175	128.5	94.0	1130	33900
35	180	130.7	96.5	1160	34800
36	185	132.9	99.0	1190	35700
37	190	135.1	101.5	1220	36600
38	195	137.3	104.0	1250	37500
39	200	139.5	106.5	1280	38400
40	205	141.7	109.0	1310	39300
41	210	143.9	111.5	1340	40200
42	215	146.1	114.0	1370	41100
43	220	148.3	116.5	1400	42000
44	225	150.5	119.0	1430	42900
45	230	152.7	121.5	1460	43800
46	235	154.9	124.0	1490	44700
47	240	157.1	126.5	1520	45600
48	245	159.3	129.0	1550	46500
49	250	161.5	131.5	1580	47400
50	255	163.7	134.0	1610	48300
51	260	165.9	136.5	1640	49200
52	265	168.1	139.0	1670	50100
53	270	170.3	141.5	1700	51000
54	275	172.5	144.0	1730	51900
55	280	174.7	146.5	1760	52800
56	285	176.9	149.0	1790	53700
57	290	179.1	151.5	1820	54600
58	295	181.3	154.0	1850	55500
59	300	183.5	156.5	1880	56400
60	305	185.7	159.0	1910	57300
61	310	187.9	161.5	1940	58200
62	315	190.1	164.0	1970	59100
63	320	192.3	166.5	2000	60000
64	325	194.5	169.0	2030	60900
65	330	196.7	171.5	2060	61800
66	335	198.9	174.0	2090	62700
67	340	201.1	176.5	2120	63600
68	345	203.3	179.0	2150	64500
69	350	205.5	181.5	2180	65400
70	355	207.7	184.0	2210	66300
71	360	209.9	186.5	2240	67200
72	365	212.1	189.0	2270	68100
73	370	214.3	191.5	2300	69000
74	375	216.5	194.0	2330	69900
75	380	218.7	196.5	2360	70800
76	385	220.9	199.0	2390	71700
77	390	223.1	201.5	2420	72600
78	395	225.3	204.0	2450	73500
79	400	227.5	206.5	2480	74400
80	405	229.7	209.0	2510	75300
81	410	231.9	211.5	2540	76200
82	415	234.1	214.0	2570	77100
83	420	236.3	216.5	2600	78000
84	425	238.5	219.0	2630	78900
85	430	240.7	221.5	2660	79800
86	435	242.9	224.0	2690	80700
87	440	245.1	226.5	2720	81600
88	445	247.3	229.0	2750	82500
89	450	249.5	231.5	2780	83400
90	455	251.7	234.0	2810	84300
91	460	253.9	236.5	2840	85200
92	465	256.1	239.0	2870	86100
93	470	258.3	241.5	2900	87000
94	475	260.5	244.0	2930	87900
95	480	262.7	246.5	2960	88800
96	485	264.9	249.0	2990	89700
97	490	267.1	251.5	3020	90600
98	495	269.3	254.0	3050	91500
99	500	271.5	256.5	3080	92400
100	505	273.7	259.0	3110	93300
101	510	275.9	261.5	3140	94200
102	515	278.1	264.0	3170	95100
103	520	280.3	266.5	3200	96000
104	525	282.5	269.0	3230	96900
105	530	284.7	271.5	3260	97800
106	535	286.9	274.0	3290	98700
107	540	289.1	276.5	3320	99600
108	545	291.3	279.0	3350	100500
109	550	293.5	281.5	3380	101400
110	555	295.7	284.0	3410	102300
111	560	297.9	286.5	3440	103200
112	565	300.1	289.0	3470	104100
113	570	302.3	291.5	3500	105000
114	575	304.5	294.0	3530	105900
115	580	306.7	296.5	3560	106800
116	585	308.9	299.0	3590	107700
117	590	311.1	301.5	3620	108600
118	595	313.3	304.0	3650	109500
119	600	315.5	306.5	3680	110400
120	605	317.7	309.0	3710	111300
121	610	319.9	311.5	3740	112200
122	615	322.1	314.0	3770	113100
123	620	324.3	316.5	3800	114000
124	625	326.5	319.0	3830	114900
125	630	328.7	321.5	3860	115800
126	635	330.9	324.0	3890	116700
127	640	333.1	326.5	3920	117600
128	645	335.3	329.0	3950	118500
129	650	337.5	331.5	3980	119400
130	655	339.7	334.0	4010	120300
131	660	341.9	336.5	4040	121200
132	665	344.1	339.0	4070	122100
133	670	346.3	341.5	4100	123000
134	675	348.5	344.0	4130	123900
135	680	350.7	346.5	4160	124800
136	685	352.9	349.0	4190	125700
137	690	355.1	351.5	4220	126600
138	695	357.3	354.0	4250	127500
139	700	359.5	356.5	4280	128400
140	705	361.7	359.0	4310	129300
141	710	363.9	361.5	4340	130200
142	715	366.1	364.0	4370	131100
143	720	368.3	366.5	4400	132000
144	725	370.5	369.0	4430	132900
145	730	372.7	371.5	4460	133800
146	735	374.9	374.0	4490	134700
147	740	377.1	376.5	4520	135600
148	745	379.3	379.0	4550	136500
149	750	381.5	381.5	4580	137400
150	755	383.7	384.0	4610	138300
151	760	385.9	386.5	4640	139200
152	765	388.1	389.0	4670	140100
153	770	390.3	391.5	4700	141000
154	775	392.5	394.0	4730	141900
155	780	394.7	396.5	4760	142800
156	785	396.9	399.0	4790	143700
157	790	399.1	401.5	4820	144600
158	795	401.3	404.0	4850	145500
159	800	403.5	406.5	4880	146400
160	805	405.7	409.0	4910	147300
161	810	407.9	411.5	4940	148200
162	815	410.1	414.0	4970	149100
163	820	412.3	416.5	5000	150000
164	825	414.5	419.0	5030	150900
165	830	416.7	421.5	5060	151800
166	835	418.9	424.0	5090	152700
167	840	421.1	426.5	5120	153600
168	845	423.3	429.0	5150	154500
169	850	425.5	431.5	5180	155400
170	855	427.7	434.0	5210	156300
171	860	429.9	436.5	5240	157200
172	865	432.1	439.0	5270	158100
173	870	434.3	441.5	5300	159000
174	875	436.5	444.0	5330	159900
175	880	438.7	446.5	5360	160800
176	885	440.9	449.0	5390	161700
177	890	443.1	451.5	5420	162600
178	895	445.3	454.0	5450	163500
179	900	447.5	456.5	5480	164400
180	905	449.7	459.0	5510	165300
181	910	451.9	461.5	5540	166200
182	915				

of a central atom. However, the effect of substituting CH_3 for H on the coupling Sn-H in this series is not strictly additive. As indicated in the first section of this abstract, Juan and Gutowsky(8) have attempted to explain deviations in the silanes and one might expect similar effects in the stannanes.

Brown and Morgan (16,17) have observed that increasing catenation in polystannanes results in a decrease in $J_{\text{Sn-CH}_3}$ and a slight increase in $J_{\text{Sn-Sn-CH}_3}$, indicating more s character in the tin-tin bonds as the number of tin-tin bonds increases in a given molecule. Further support for this is given by the fact that the tin-carbon stretching frequencies were found to increase linearly with the coupling constant for the series studied.

Drago and Matwiyoff (14,18) have studied the $J_{\text{C}^{13}\text{-H}}$ values for a series of tetramethyl derivatives of group IVB elements and find the coupling constant values parallel the M-C bond energies.

REFERENCES

1. M. Karplus and D. M. Grant, Proc. Nat. Acad. Sci., 45, 1269 (1959).
2. N. Muller and D. E. Pritchard, J. Chem. Phys., 31, 768 (1959).
3. N. F. Ramsey, Phys. Rev., 91, 303 (1953).
4. J. A. Pople, W. G. Schneider, and H. J. Bernstein, High Resolution Nuclear Magnetic Resonance, Mc-Graw Hill Book Company, Inc., New York, 1959, p. 184.
5. M. Karplus and D. H. Anderson, J. Chem. Phys., 30, 6 (1959).
6. M. Karplus, ibid, 30, 11 (1959).
7. W. Ripka, Organic Seminars, University of Illinois, 134(1962).
8. C. Juan and H. S. Gutowsky, J. Chem. Phys., 37, 2198 (1962).
9. J. R. Holmes and H. D. Kaesz, J. Am. Chem. Soc., 83, 3903 (1961).
10. G. P. Van der Kelen, Nature, 193, 1069 (1962).
11. H. A. Bent, Can. J. Chem., 38, 1235 (1960).
12. H. A. Bent, Chem. Revs., 61, 275 (1961).
13. J. J. Burke and P. C. Lauterbur, J. Am. Chem. Soc., 83, 326 (1961).
14. N. A. Matwiyoff, Doctoral Dissertation, University of Illinois, (1963).
15. N. Flitcroft and H. D. Kaesz, J. Am. Chem. Soc., (in press).
16. T. L. Brown and G. L. Morgan, Inorg. Chem., (in press).
17. G. L. Morgan, Doctoral Dissertation, University of Illinois, (1963).
18. R. S. Drago and N. A. Matwiyoff, to be published.

MOLECULAR EXCITATION IN SHOCK TUBES

Sue F. Eilers

March 21, 1963

I. INTRODUCTION

Investigators concerned with the effects of shock waves produced by supersonic aircraft and in nuclear explosions discovered that the shock tube could be used to study the pressures due to shock waves and that these waves produce high temperatures in the gases through which they pass (1,2). It is this property which has made them of interest to chemists.

II. THEORY (1-22)

At the instant the diaphragm in the shock tube is ruptured, the molecules of gas in the low-pressure section of the tube are moving over relatively long mean free paths, as a consequence of the low density of the gas, and at an average speed of about 1,000 feet per second. Compression of the gas in the first little slab ahead of the outrushing piston of high-pressure gas shortens the mean free path of the molecules and steps up their rate of collision, thereby increasing the mean velocity and, consequently, the temperature of the gas. The first minute disturbance passes forward at the speed of sound into the next slab, compressing and preheating it for the arrival of the next wave, thus causing that wave to move faster. As the waves overtake one another building up into a large pressure pulse and as the pressure pulse grows into a shock wave, the mean free path of the molecules in the wave grows still shorter, and the temperature rises.

The shock wave must spread out over some finite distance in the gas, measureable in terms of the length of the mean free path. Across this distance the gas density, temperature, pressure, and velocity all change. As the shock wave forms and these gradients grow steeper, energy in the form of heat and momentum flows more swiftly from regions of high temperature and velocity to regions of low temperature and velocity in the wake of the wave (1, 23-35). Ultimately, the opposing mechanisms of steepening and spreading balance and the shock wave profile becomes stable.

At this point, other phenomena may be observed. By no means is all of the kinetic energy of the shock wave transformed into heat in the form of translational energy. In fact, with each increase in temperature an increasing percentage of the energy is transformed into other energy modes.

In the zone of highest temperature just behind the shock front, the molecule-molecule collisions cause rotation as well as elastic collisions. Back a little farther, the interatomic bonds begin to vibrate. The transformation of energy from translation to rotation to vibration is known as energy relaxation, and it has been demonstrated by numerous experiments in diatomic and polyatomic gases (79-92). As the translational energy (and along with it the temperature) drops from its highest value immediately behind the shock wave

front to a lower value some distance behind it, the gas density behind the shock front changes accordingly. These density profiles show up in photographs made with the Mach-Zender interferometer and can be used in calculating the time required to transform energy from one mode to another. These relaxation times are an indication of the number of molecule-molecule collisions necessary to produce the energy transfer. The collision cross section for energy exchange between two molecules can thereby be computed and compared with values calculated from quantum theory. Other methods for studying these phenomena include x-ray, infrared and ultraviolet spectroscopy (36-45). Both experiment and theory show that for all but the weakest shock waves, the translational and rotational energies are quickly activated, usually within a few mean free paths, whereas the vibrational mode lags considerably in activation, as a result of the small probability of vibrational transitions. Experiments detecting this can be interpreted in terms of a vibrational relaxation time or a vibrational transition probability (6,9,13,19,79-92). It is difficult to deduce from these experiments the mechanics of the collision in which a given vibrational transition takes place, since what is measured is the transition probability averaged over a Maxwellian distribution of translational energies. However, considerable progress is being made in the theory of these low-energy collisions, and proper predictions of vibrational relaxation times are one test of their validity. Also, molecular beam experiments are valuable supplements to shock tube measurements in this regard.

It was not long after the invention of the shock tube that investigators noticed that the hot gases in the tube emit light. In fact, when a gas has attained a sufficiently high temperature, this is one of the principal modes by which energy is dissipated. This process involves electronic transitions from one quantum state to another. For any given temperature, there is a specific distribution of electron-energy states throughout the atoms in the gas. The electrons deexcite either by quantum jumps or as the result of inelastic collisions. The higher the temperature, the greater the number of inelastic collisions causing excitation and, consequently, the greater the number of photons emitted.

The mean lifetime of the excited state is in principle calculable from theory; however, such calculations are prohibitively difficult for all but the simplest atoms and molecules. While experimental results obtained by heating gases with arc and spark discharges are blurred by the introduction of impurities from the electrodes, the shock tube provides impurity-free hot gases at temperatures and densities that can be precisely controlled by adjustment of the pressures in the two sections of the tube (1,9,10,73-78).

Other studies done in shock tubes include kinetics (46-54), high-temperature reactions (55-61) and measurements of dissociation constants (62-72).

III. A SPECIFIC CASE: STUDIES OF VIBRATION RELAXATION IN NITROGEN (II) OXIDE

Since one of the most interesting of the experiments done in shock tubes is the determination of vibrational relaxation times, and there have been several papers on nitrogen (II) oxide, giving parameters at a wide temperature range, this was chosen as an example of the type of data which can be obtained.

In the earliest of these studies (89), nitrogen (II) oxide was examined at temperatures less than 1000°K. It was assumed that no dissociation took place in these experiments and that the ideal gas law was valid. Because of the small excitation of the vibrational levels, only the first vibrational level was considered in the relaxation time. The formula for the transition probability was found to be

$$P_{10} = 2.86 / \tau_1 P_1 (P/P_0)^2 (T_a)^{1/2} \quad (1)$$

where P_{10} is in mm. of mercury, τ_1 in microseconds, T_a in degrees K. Using Lennard-Jones parameters ($r_0 = 3.6 \times 10^{-8}$ cm., $\epsilon/k = 91^\circ\text{K}$), it was found that the experimental measurements for relaxation were about 5×10^5 as large as the calculated values. However, fitting the Lennard-Jones potential formula to data including the possibility of dimer formation, Robbins found $\epsilon/k = 2000^\circ\text{K}$ and $r_0 = 2.15 \times 10^{-8}$ cm. When P_{10} was calculated using these parameters, the values agreed fairly well.

Robbins had used ultraviolet light to examine the vibrational mode in this oxide. Later his data was compared with the infrared data of Monson and Allport (from the same laboratory), with very good correlation (92).

Roth examined nitrogen (II) oxide at temperatures between 5000 and 10,000°K (Tables I and II) a range where the gas was considered to be excited (90).

Wrey has examined nitrogen (II) oxide at temperatures between those above and those of Robbins. Using the following equations, he found the probabilities of transitions.

$$P_{10} = A_3 (\bar{\epsilon} - \epsilon_0)^{-1/2}, T \text{ } ^\circ\text{K} \quad (2)$$

$$P_{10} = A_4 T^{-1} \exp(-\epsilon_0/kT), T \text{ } ^\circ\text{K} \quad (3)$$

where A's = constants determined experimentally and $\bar{\epsilon}$ = average kinetic energy of colliding molecules at infinity. Using the second set of parameters from Robbins, he found the predicted values to be higher than the experimental by a factor of about 20. Extrapolation to lower temperatures of Wrey's values for P_{10} gives much lower values than those calculated by Robbins. However, Wrey apparently also assumed absence of excited states, which, as the temperature rises, is an increasingly poor assumption.

IV. CONCLUSION

The shock tube provides a relatively easy, very fast method for heating gases to high temperatures. At intermediate temperatures, it may be used for the study of vibrational modes in excited states, as in the studies discussed here. At slightly higher temperatures, it may be used to study dissociation rates and constants. At still higher temperatures, such as those produced by electromotive shocks, the phenomena of plasmas become available for study (21,22). Therefore, it seems apparent that shock tubes will become increasingly familiar laboratory apparatus and that they should provide some interesting answers to many questions concerning fundamental physical problems.

TABLE I (90)

Relaxation times for $V = 1 \rightarrow V = 0$ in (A^2E^+) state of NO

<u>T, °K</u>	<u>ρ_{Xe} (a) 10^{17} cm^{-3}</u>	<u>Mole % Air</u>	<u>τ, μsec</u>
9900	1.25	2.5	34
8875	1.30	5.0	32
8875	1.30	5.0	49
8825	1.24	2.5	31
8275	1.24	2.5	39
7975	1.29	5.0	49
7400	1.29	5.0	91
7125	2.47	2.5	56
6950	2.47	2.5	41
6700	2.46	2.5	46
6650	2.56	5.0	64
6375	2.55	5.0	56
5425	2.54	5.0	106

a) Initial pressures in the low pressure section of the shock tube were 1 mm. and 2 mm. for ρ_{Xe} = about $1.25 \times 10^{17} \text{ cm}^{-3}$ and about $2.5 \times 10^{17} \text{ cm}^{-3}$, respectively, where ρ refers to the density of the element in molecules/cm³.

TABLE II (90)

Relaxation times referred to density at 1 atm. pressure for $V = 1 \rightarrow V = 0$ in NO ($A^2\Gamma^+$).

T, °K	5000	6000	7000	8000	9000	10000
τ , sec.	28	18	12	8	6	5

REFERENCES

- A. General
1. McChesney, M., Scientific American, 208, No. 2, 109 (1963).
 2. Payman, W., and Shepherd, W. D.F., Proc. Roy. Soc. (London) A 186, 293 (1946).
 3. Paterson, S., Phys. Soc. (London), 61, 119 (1948).
 4. Travers, S., Mem. artillerie franc., 27, 421 (1953).
 5. Crown, J. C., Phys. Rev., 79, 229 (1950).
 6. Walker, R. A., J. Chem. Phys., 19, 494 (1951).
 7. Zoller, K., Z. Physik., 130, 1 (1951).
 8. Benson, J. M., J. Applied Phys., 23, 757 (1952).
 9. Courant, R., and Friedrichs, K. O., Supersonic Flow and Shock Waves, Interscience Publishers, Inc., New York, N. Y., 1948.
 10. Patterson, G. N., Molecular Flow of Gases, John Wiley and Sons, Inc. New York, N. Y., 1956, Chapters 2, 3, and 4.
 11. Christian, R. H. and Yarger, F. L., J. Chem. Phys., 23, 2042 (1955).
 12. Knight, H. I., and Rink, J. P., Ibid., 35, 199 (1961).
 13. Korobkin, I., and Slawsky, Z. I., Ibid., 37, 226 (1962).
 14. Duffey, G. H., Ibid., 23, 401 (1955).
 15. Lin, S., Resler, E. L., and Kantlowitz, A., J. Appl. Phys. 26, 95 (1955).
 16. Guenoche, H., and Manson, N., Compt. rend., 235, 1617 (1952).
 17. Duvall, G. E., and Zwolinski, B. J., Ind. Eng. Chem., 47, 1182 (1955).
 18. Guman, W. J., J. Appl. Phys., 27, 663 (1956).
 19. Zellazny, R., Phys. Rev., 117, 1 (1960).
 20. Sen, H. K., and Guess, A. W., Ibid., 108, 560 (1957).
 21. Kolb, A. C., Ibid., 107, 345 (1957).
 22. Hoffman, F., and Teller, E., Ibid., 80, 692 (1950).
- B. Structure of a Shock Wave
23. Hansen, K., and Hornig, D. F., J. Chem. Phys. 33, 913 (1960).
 24. Bond, J. W., Jr., Phys. Rev. 105, 1683 (1957).
 25. Cowan, G. R., and Hornig, D. F., J. Chem. Phys., 18, 1008 (1950).
 26. Greene, E. F., Cowan, G. R., and Hornig, D. F., Ibid., 19, 427 (1951).
 27. Greene, E. F., and Hornig, D. F., Ibid., 21, 617 (1953).
 28. Kistiakowdsky, G. B., Knight, H. T., and Malin, M. E., Ibid., 20, 994 (1952).
 29. Weir, A., Jr., and Morrison, R. B., Ind. Eng. Chem. 46, 1056 (1954).
 30. Liepmann, H. W., Phys. Rev., 71, 465 (1946).
 31. Benschader, D., Ibid., 91, 470 (1953).
 32. Wagner, H. G., Z. Elektrochem., 59, 906 (1955).

33. Anderson, W. H., and Hornig, D. F., J. Chem. Phys., 24, 767 (1956).
34. Tidman, P. R., Phys. Rev., 111, 1439 (1958).
35. Griffith, W., Brickl, P., and Blackman, V., Ibid., 102, 1209 (1956).

C. Experimental Methods

36. Ikeda, R. M., J. Chem. Phys., 33, 311 (1960).
37. Dean, J., and de Boer, P. C. T., J. Chem. Phys., 36, 1222 (1962).
38. Bradley, J. N., and Kistiakowsky, G. B., Ibid., 35, 264 (1961).
Comment: Fenimore, C. P., Ibid., 35, 2243 (1961).
39. Gardiner, W. C., Jr., and Kistiakowsky, G. B., Ibid., 34, 1080 (1961).
40. Petschek, H. E., Rose, P. H., Glick, H. S., Kane, Anne, and Kantiowitz, A., J. Appl. Phys., 26, 83 (1955).
41. Knight, H. T., and Duff, R. E., Phys. Rev., 94, 784 (1954).
42. Resler, E. L., Jr., and Scheibe, M., Ind. Eng. Chem., 47, 1182 (1955).
43. Harshbarger, F., J. Chem. Phys., 24, 1261 (1956).
44. Rosa, R. J., Phys. Rev., 99, 633A (1955).
45. Savitt, J., Ibid., 98, 1158 A (1955).

D. Reaction Studies

46. Carrington, T., and Davidson, N., J. Chem. Phys., 57, 418 (1953).
47. Britton, D., Davidson, N., and Schott, G., Disc. Faraday Soc., 17, 58 (1954).
48. Plooster, M. W., and Garvin, P., J. Am. Chem. Soc., 78, 6003 (1956).
49. Schott, G. L., J. Chem. Phys., 32, 710 (1960).
50. Strehlow, R. Z., and Case, C. T., Ibid., 35, 1506 (1961).
Comment: Rudinger, G., Ibid., 35, 1507 (1961).
51. Patterson, W. L., Jr., and Greene, E. F., Ibid., 36, 1146 (1962).
52. Wray, K. L., and Teare, J. D., Ibid., 36, 2582 (1962).
53. Hertzberg, A., and Squire, W., Phys. Rev., 91, 469 (1953).
54. Britton, D., and Davidson, N., J. Chem. Phys., 23, 246 (1955).
55. Bradley, J. N., and Kistiakowsky, G. B., Ibid., 35, 264 (1961).
56. Manson, N., Compt. rend. 227, 720 (1948).
57. Duff, R. E., Knight, H. T., and Wright, H. R., J. Chem. Phys., 22, 1618 (1954).
58. Greene, E. F., J. Am. Chem. Soc., 76, 2127 (1954).
59. Berets, D. J., Greene, E. F., and Kistiakowsky, G. B., Ibid., 72, 1080, 1086 (1950).
60. Fay, J. A., 4th Sym. on Combustion 1952, 501.
61. Duff, R. E., and Minshall, F. S., Phys. Rev., 108, 1207 (1957).
62. Knight, H. T., and Rink, J. P., J. Chem. Phys., 35, 199 (1961).
63. Tsang, W., Bauer, S. H., and Cowperthwaite, M., Ibid., 36, 1768 (1962).
64. Hand, C. W., and Kistiakowsky, G. B., Ibid., 37, 1239 (1962).
65. Rink, J. P., Knight, H. T., and Duff, R. E., Ibid., 34, 1942 (1961).
66. Hiraoka, H., and Hardwick, R., Ibid., 36, 1715, 2164 (1962).
67. Sutton, E. A., Ibid., 36, 2923 (1962).
68. Rink, J. P., Ibid., 36, 262, 1398 (1962).
69. Patch, R. W., Ibid., 36, 1919 (1962).
70. Gardiner, W. C., Jr., and Kistiakowsky, G. B., Ibid., 35, 1765 (1961).
71. Toennies, J. P., and Greene, E. F., Ibid., 23, 1365 (1955).
72. Bond, J. W., Jr., Phys. Rev., 98, 285 (1954).

E. Luminosity in Shocked Gases

73. Roth, W., and Gloersen, P., J. Chem. Phys., 29, 820 (1958).
74. Mies, F. H., Ibid., 37, 1101 (1962).
75. Kistiakowsky, G. B., and Richards, L. W., Ibid., 36, 1707 (1962).
76. Takoyama, H., and Gullman, A., Ibid., 32, 634 (1960).
77. Bradley, J. N., Ibid., 32, 1875 (1960).
Comment: Roth, W., Ibid., 32, 1876 (1960).
78. Turner, E. S., Phys. Rev., 99, 633 A (1955).

F. Vibration Relaxation Studies

79. Roth, W., J. Chem. Phys. 31, 720 (1959).
80. Wray, K. L., Ibid., 37, 1254 (1962).
81. Witteman, J., Ibid., 37, 655 (1962).
82. Hooker, W. J., and Milliken, R. C., Ibid., 38, 214 (1963).
83. Davies, W. O., Ibid., 36, 292 (1962).
84. Davies, W. O., and Sternberg, M., Ibid., 34, 1373 (1961).
85. Matthews, D. L., Ibid., 34, 639 (1961).
86. Smiley, E. F., Winkler, E. H., and Slawsky, Z. I., Ibid., 20, 923 (1952).
87. Smiley, E. F., and Winkler, E. H., Ibid., 22, 2018 (1954).
88. Glick, H. S., Squire, W., Herzberg, A., 5th Sym. on Combustion, 1954, 393.
89. Robben, F., Ibid., 31, 420 (1959).
90. Roth, W., Ibid., 34, 999 (1961).
91. Wray, K. L., Ibid., 36, 2597 (1962).
92. Robben, F., Monson, P. R., and Allport, J. J., Ibid., 33, 630 (1960).

BORON POLYMERS

Gerald Tennenhouse

March 26, 1963


Boron compounds are of particular interest in the mushrooming field of inorganic polymers. This is due in part to the chemical and thermal stability of certain boron-nitrogen, boron-phosphorus, and boron-oxygen compounds. However, their popularity is also due to the exhibition of properties usually associated with organic polymers (18, 19).

It is convenient to divide the topic of boron-containing polymers into four loosely organized groups: compounds characterized by boron-nitrogen bonds; compounds characterized by boron-phosphorus bonds; compounds characterized by boron-oxygen bonds; and compounds which do not readily fit into any of the above-mentioned categories.

BORON-NITROGEN POLYMERS


Borazole was first prepared by Stock and Pohland (1) by the reaction of ammonia and diborane. The heat stability of borazole itself and various substituted borazoles (2, 3) has prompted investigations of boron-nitrogen chemistry in the hope of discovering heat-stable inorganic polymers.


Several improved methods for the synthesis of borazoles have been devised (4), for example:




The B-Cl chlorine atoms may then be replaced by alkyl or aryl groups by the action of Grignard reagents. Other anions, such as CN^- , CNS^- , NO_2^- , and RO^- may be introduced by the action of an appropriate metal salt on an ethereal solution of the B-Cl borazole (4).

A great many attempts have been made to prepare high molecular weight inorganic polymers by joining borazole rings in one manner or another. The ultimate result of such work is hexagonal boron nitride. However, the degree of cross linking is so high that the material does not have the flexibility usually associated with the term polymer and is of little commercial use. Some attempts to join borazole rings are illustrated by the following examples (7):







Non-cyclic boron-nitrogen compounds have also been prepared (2):



Borazole is commonly referred to as inorganic benzene. However, another series of boron-nitrogen compounds has been prepared which may be thought of as saturated (11, 12), for example:



Both of these compounds show unusual resistance towards hydrolysis as well as good thermal stability.

BORON-PHOSPHORUS POLYMERS


Boron-phosphorus chemistry is a relatively new field. Gamble and Gilmont (10) first caused phosphine and diborane to react and obtained a poorly characterized white solid which was thought to be a highly cross-linked polymer.

Burg (9) found that more distinct compounds could be obtained by the use of dimethyl phosphine instead of phosphine. The compounds obtained by Burg were found to have remarkable chemical, as well as thermal, stability.

... ..

... ..

... ..



... ..


... ..

... ..




Both Burg (13) and Wagner (14, 15) have developed improved methods of synthesizing both boron-phosphorus monomers and polymers.

BORON-OXYGEN POLYMERS

The boron-oxygen system of polymers is very attractive because of the low cost and relative ease of handling of the starting materials. Unfortunately, many of these compounds are easily hydrolyzed. A typical synthesis may be formulated as (2):



MISCELLANEOUS POLYMERS



Some of these compounds show good thermal and chemical stability, as well as exhibiting certain properties usually associated with organic polymers.

A great many compounds have been prepared which do not fit conveniently into any broad category. Examples include the polymer obtained from the reaction of diborane and hydrogen sulfide (21); the product of the reaction between B₅H₉ and acetylene in a discharge tube (2); and the product of the reaction between diborane and carbon monoxide in a discharge tube (2). These and other reactions have yielded products that have been identified as polymers. However, these polymers do not seem to be of interest at the present time.

Both parts (A) and (B) are developed in the same way as in the previous part of the paper.

APPENDIX

The following table gives the results of the experiments. The first column shows the number of molecules of the polymer, the second column the number of molecules of the monomer, and the third column the number of molecules of the solvent. The fourth column shows the number of molecules of the polymer which are still present after the reaction has taken place. The fifth column shows the number of molecules of the monomer which are still present after the reaction has taken place. The sixth column shows the number of molecules of the solvent which are still present after the reaction has taken place. The seventh column shows the number of molecules of the polymer which are still present after the reaction has taken place. The eighth column shows the number of molecules of the monomer which are still present after the reaction has taken place. The ninth column shows the number of molecules of the solvent which are still present after the reaction has taken place.

Time (min)	[M] (mole/l)	[S] (mole/l)	[P] (mole/l)	[M] (mole/l)	[S] (mole/l)	[P] (mole/l)
0	0.100	0.100	0.000	0.100	0.100	0.000
10	0.095	0.095	0.005	0.095	0.095	0.005
20	0.090	0.090	0.010	0.090	0.090	0.010
30	0.085	0.085	0.015	0.085	0.085	0.015
40	0.080	0.080	0.020	0.080	0.080	0.020
50	0.075	0.075	0.025	0.075	0.075	0.025
60	0.070	0.070	0.030	0.070	0.070	0.030
70	0.065	0.065	0.035	0.065	0.065	0.035
80	0.060	0.060	0.040	0.060	0.060	0.040
90	0.055	0.055	0.045	0.055	0.055	0.045
100	0.050	0.050	0.050	0.050	0.050	0.050

Some of these experiments were also carried out in the presence of a small amount of water. The results are given in the following table. It will be seen that the presence of water has a marked effect on the rate of polymerization. The rate is much slower in the presence of water than in its absence. This is probably due to the fact that water acts as a chain transfer agent, thus terminating the polymer chains. The results are given in the following table. It will be seen that the presence of water has a marked effect on the rate of polymerization. The rate is much slower in the presence of water than in its absence. This is probably due to the fact that water acts as a chain transfer agent, thus terminating the polymer chains.

BIBLIOGRAPHY

1. Stock, A., and Pohland, E., Ber., 59, 2215 (1926).
2. McCloskey, A. L., Inorganic Polymers, edited by Stone, F. G. A., and Graham, W. A. G., Academic Press, New York, p. 162 (1962).
3. Newsome, H. C., English, W. D., McCloskey, A. L., and Woods, W.G., J. Am. Chem. Soc., 83, 4134 (1961).
4. Sheldon, J. C., and Smith, B. C., Quart. Revs., 14, 200 (1960).
5. Ryshkewitsch, G. E., Harris, J. J., and Sisler, H. H., J. Am. Chem. Soc., 80, 4515 (1958).
6. Aubrey, D. W., and Lappert, M. F., J. Chem. Soc., 2927 (1959).
7. Zamyatina, V. A., and Bekasova, N. I., Russ. Chem. Revs., 30, 22 (1961).
8. Koton, M. M., Russ. Chem. Revs., 31, 81 (1962).
9. Burg, A., and Wagner, R. I., J. Am. Chem. Soc., 75, 3872 (1953).
10. Gamble, L., and Gilmont, P., J. Am. Chem. Soc., 62, 717 (1940).
11. Bissot, T. C., and Parry, R. W., J. Am. Chem. Soc., 77, 3481 (1955).
12. Campbell, G. W., and Johnson, L. W., J. Am. Chem. Soc., 81, 3800 (1959).
13. Burg, A. B., and Slota, P. J., J. Am. Chem. Soc., 82, 2145 (1960).
14. Wagner, R. I., Caserio, F. F., and Freeman, L. D., W.A.D.C. Wright Air Development Center, Tech. Rept. 57-126 part III (1959).
15. Wagner, R. I., W.A.D.C. (Wright Air Development Center) Rept. 59-427 (1960).
16. Wagner, R. I., and Caserio, F. F., J. Inorg. and Nucl. Chem., 11, 258 (1959).
17. Burg, A. B., J. Inorg. and Nucl. Chem. 11, 257 (1959).
18. Heying, T. L., Office of Naval Research, Contract Nonr. 3395(00) Task No. 356-431, Tech. Dept. No. 11 (1962).
19. Vale, R. L., J. Chem. Soc., 2252 (1960).
20. Musgrave, O. C., Chem. and Ind. (London), 1152 (1957).
21. Heying, T. L. and Schroeder, H. A., Office of Naval Research, Tech. Rept. No. 8, Boron Polymers, Contract Nonr. 3395(00), Quarterly Report, Report period 1 March 1962-31 May 1962.

CONFIDENTIAL

... (1952) ...
... (1953) ...
... (1954) ...
... (1955) ...
... (1956) ...
... (1957) ...
... (1958) ...
... (1959) ...
... (1960) ...
... (1961) ...
... (1962) ...
... (1963) ...
... (1964) ...
... (1965) ...
... (1966) ...
... (1967) ...
... (1968) ...
... (1969) ...
... (1970) ...
... (1971) ...
... (1972) ...
... (1973) ...
... (1974) ...
... (1975) ...
... (1976) ...
... (1977) ...
... (1978) ...
... (1979) ...
... (1980) ...
... (1981) ...
... (1982) ...
... (1983) ...
... (1984) ...
... (1985) ...
... (1986) ...
... (1987) ...
... (1988) ...
... (1989) ...
... (1990) ...
... (1991) ...
... (1992) ...
... (1993) ...
... (1994) ...
... (1995) ...
... (1996) ...
... (1997) ...
... (1998) ...
... (1999) ...
... (2000) ...
... (2001) ...
... (2002) ...
... (2003) ...
... (2004) ...
... (2005) ...
... (2006) ...
... (2007) ...
... (2008) ...
... (2009) ...
... (2010) ...
... (2011) ...
... (2012) ...
... (2013) ...
... (2014) ...
... (2015) ...
... (2016) ...
... (2017) ...
... (2018) ...
... (2019) ...
... (2020) ...
... (2021) ...
... (2022) ...
... (2023) ...
... (2024) ...
... (2025) ...

MACROCYCLIC COORDINATION COMPOUNDS

E. J. Olszewski

March 28, 1963


Introduction

While metal-chelate compounds having five- and six-membered rings are well established in the literature, evidence for rings of larger size is not as conclusive. O'Brien (1) showed that, while many macrocyclic metal complexes had been reported, the great majority of structures had been postulated on the basis of analytical data alone. Lack of x-ray data, the possibility of several linkages, or of polymerization served to make the existence of large rings doubtful. However, the utilization of rigid geometrical structures and increasing use of physical measurements, have made possible the isolation and characterization of macrocyclic rings. These are essentially of three types; inner complexes, compounds with the $\text{Cu}_2(\text{C}_2\text{H}_3\text{O}_2)_4 \cdot 2\text{H}_2\text{O}$ structure, and ferrocene derivatives.

Rigid Templates


The formation of normal five- or six-membered chelates often serves as a frame to which larger rings may be attached.

Magnetic susceptibility measurements (2) on




showed the paramagnetism of the copper(II) ion to decrease as n increased. The stability of the salicylaldehyde ring and the flexibility of the hydrocarbon chain could account for the formation of these complexes.

Monomeric and dimeric species of beryllium (II) tetraketones have been reported on the basis of analytical data, molecular weight determinations, and infrared spectra (3).




When R is small, e.g. $-(\text{CH}_2)_6-$ or $-\text{O}(\text{CH}_2)_4\text{O}-$, then dimerization is favored. In both cases, however, the compounds are prepared by thermal degradation of the corresponding low molecular weight polymers.

The analogous copper(II) complex of 1,3-diphenyltriazen was studied by Harris and Martin (7) and found to be diamagnetic in the solid state and in benzene. They postulated the following structure:




Harris, Hoskins and Martin (8) found the divalent nickel, copper and palladium complexes of 1,3-diphenyltriazene to be dimeric in benzene and diamagnetic.

Crystal studies by Brown and Dunitz (9) on the copper(I) chelate of diazoaminobenzene show a planar structure with a linearly coordinated copper atom. The N-Cu-N angle is almost linear.



Ferrocenylene Frames

Hauser and Cain (10) reported a macrocycle with two different metal ions. The compound was a -diketone salt of divalent copper derived from the corresponding ferrocene derivative:




The yellow powder was recrystallized from benzene and melted above 300°C.


Heterocycles containing silicon bridges were isolated by Schaaf and co-workers (11) from 1,1'-bis-(dimethylethoxysilyl)-ferrocene. Structural assignments were made on the basis of infrared spectra and molecular weight determinations.

Rinehart and Curby(12) showed that cyclization would be heteroannular or homoannular depending on the size of the chain in the starting material. The product formed by homoannular ring closure could be distinguished by the absence of IR bands at 1110 and 1004 cm^{-1} .

Lüttringhaus and Kullick (13) obtained bridged ferrocenes in low yields from α,ω -bis-(cyclopentadienyl)-alkanes. The major products were dimers, and polymers of varying molecular weight.




Treatment of ferrocene-1,1'-bis-(propionic acid) with trifluoroacetic anhydride (14) gave 1,1':3,3'-bis-(trimethylene)-ferrocene, after reduction. Schlögl and Seiler (15) also isolated a series of acyloins in various yields:




Hauser, Cain and Mashburn (16) treated diacetyl ferrocene with benzaldehyde and aqueous NaOH to get a small amount of cyclic product. While they postulated a monomeric structure, the possibility of a dimer was not ruled out.

Furkik and co-workers (17) obtained yields from 10-78% for the following reaction:



Gustafson (18) prepared 3,3'-trimethylene-1,1'- α -ketotri-methyleneferrocene by the following route:



Brown (19) was able to reduce the product by means of H_2 and platinum.

Other reducing agents were examined by Gustafson (18), who found that reduction was quantitative when a 10/15 mole ratio of $LiAlH_4/AlCl_3$ was used as opposed to a 63% yield when $LiAlH_4$ was used alone. The products were characterized by IR and NMR spectra.

Other compounds containing large rings were also isolated by Gustafson (18).

Conclusion

The preparation of macrocyclic chelate complexes is possible if coordinating ligands can be attached to a rigid molecular frame. At present the ferrocenylene structure seems to be the most promising as evidenced by the variety of ferrocene complexes already reported in the literature. Unfortunately very little work on crystal structure has been done on these compounds.

References

1. I. D. O'Brien, "The Chemistry of the Coordination Compounds," J. C. Bailar, Jr. Ed., Reinhold Publ. Corp., New York, 1956, P. 253.
2. M. Calvin and C. H. Barkeley, J. Am. Chem. Soc., 68, 2267 (1946).
3. R. W. Kluber and J. W. Lewis, J. Am. Chem. Soc. 82, 5777 (1960).
4. M. M. Coombs and R. P. Houghton, J. Chem. Soc., 5015 (1961).
5. R. Snellgrove and E. L. King, J. Am. Chem. Soc., 84, 4609 (1962).
6. F. P. Dwyer and D. P. Mellor, J. Am. Chem. Soc., 63, 81 (1941).
7. C. M. Harris and R. L. Martin, Proc. Chem. Soc. 259 (1958).
8. C. M. Harris, B. F. Hoskins, and R. L. Martin, J. Chem. Soc. 3728 (1959).
9. I. D. Brown and J. D. Dunitz, Acta Cryst., 14, 480 (1961).
10. C. R. Hauser and C. E. Cain, J. Org. Chem., 23, 1142 (1958).
11. R. L. Schaaf, P. T. Kan and C. T. Lenk, J. Org. Chem., 26, 1790 (1961).
12. K. L. Rinehart, Jr. and R. J. Curby, Jr., J. Am. Chem. Soc., 79, 3290 (1957).

13. A. Lüttringhaus and W. Kullick, *Angew. Chem.*, 70, 438 (1958).
14. K. Schlögl and H. Seiler, *Tetrahedron Letters*, No. 7, 4(1960).
15. K. Schlögl and H. Seiler, *Monatsh. Chem.*, 91, 79 (1960).
16. T. A. Mashburn, Jr., C. E. Cain, and C. R. Hauser, *J. Org. Chem.* 25, 1982 (1960).
17. M. Furkik, S. Toma, J. Suchý and P. Elečko, *Chem. Zvesti*, 15, 45 (1961).
18. D. H. Gustafson, Ph. D. Thesis, Univ. of Illinois (1961).
19. W. G. Brown, *Org. Reactions*, 6, 469 (1951).

E. S. R. OF TRANSITION METAL ION COMPLEXES

Peter C. Van Der Voorn

April 2, 1963

In electron spin resonance we study the absorption of electromagnetic radiation which effects transitions between energy levels of the spin states of unpaired electrons. By application of an external magnetic field it is possible to vary these energy levels. Since the energy differences are of the order found in the microwave region, absorption of microwave radiation is studied. In some aspects ESR is similar to NMR where the transitions studied occur between energy levels of the nuclear spin states, again after the application of an external magnetic field.

The instrumentation consists of a monochromatic source of microwave radiation, an absorption cell in a magnetic field at right angles to the beam of radiation, and a detector. Measurements are most commonly made at 1.25-3 cm wavelengths with fields ranging from 8000-3000 gauss. Although in principle either the microwave frequency or the field strength can be varied while the other is held constant, in practice it is simpler to vary the field strength.

The spin angular momentum M_S (spin S) of an electron can have values of $\pm 1/2$. In the absence of a magnetic field, there is no preference and the spin energy is doubly degenerate. After a magnetic field is applied the electron precesses about an axis parallel to the field. By absorbing a quantity of radiation $h\nu$, the spin changes and the electron precesses about an axis anti-parallel to the field. The resonance condition is given by $h\nu = g\beta H$, where h is Planck's constant, ν the frequency of radiation, β the Bohr magneton, H the strength of the magnetic field and g , a proportionality constant which has the value of 2.0023 for a free electron. In general with n unpaired electrons, the spin degeneracy is $n+1$ and is split by the magnetic field into $n+1$ levels; n transitions can then occur between the S levels under the selection rule $M_S = \pm 1$.

When one considers the unpaired electrons in transition metal ions in the solid state, it is necessary to take into account the complete energy state of the ion. The complete Hamiltonian can be written as (6):

$$W = W_F + V + W_{LS} + W_{SS} + H(L + 2S) \cdot W_N - \gamma \beta_N H \cdot I$$

where W_F includes the energies of the free ion in the gaseous state ($\sim 10^5 \text{ cm}^{-1}$); V is the electrostatic energy due to the crystal field ($\sim 10^3 - 10^4 \text{ cm}^{-1}$); W_{LS} represents the spin-orbit coupling ($\sim 10^2 \text{ cm}^{-1}$); W_{SS} the magnetic interaction between electrons ($\sim 1 \text{ cm}^{-1}$); $H(L + 2S)$ the effect of the external magnetic field (1 cm^{-1}); W_N interactions due to the nucleus ($\sim 10^2 \text{ cm}^{-1}$) and $\gamma \beta_N H \cdot I$ the effect of the magnetic field on the nuclear magnetic moment (10^{-3} cm^{-1}); this latter effect can usually be neglected in ESR.

When a gaseous ion is placed into a crystal, the degeneracy of the d-orbitals will be resolved by the crystal field: for example, for Cu^{2+} the 2D ground state of the free ion becomes orbitally non-degenerate after considering the effects of a cubic field and a tetragonal distortion, but the spin degeneracy is not affected. In some metal ion crystals the spin degeneracy can also be resolved by the crystal field, but only through spin orbit coupling. This is called zero-field splitting (e.g. Cr^{+3}). However in the case of an odd number of unpaired electrons the spin remains doubly degenerate; this is known as Kramer's degeneracy (23). In any case the spin degeneracy is completely resolved by a magnetic field. Using a fixed frequency and an increasing magnetic field, n electrons will give n transitions but in the absence of zero field splitting these transitions are coincident in energy and only one peak will be observed (e.g. Ni^{2+} in MgO). In the presence of zero-field splitting all transitions are observed (e.g. Cr^{+3}) (4).

When an electron interacts with a nucleus which possesses an angular momentum M_I (nuclear spin I), $2I + 1$ orientations are possible and the main peak is split into $2I + 1$ components. The selection rule is $M_I = 0$. This was first observed in $\text{Cu}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ (31) which gives four peaks because I for Cu^{65} and Cu^{63} is $3/2$. In copperbissalicylaldehyde each of the four components due to the interaction with the Cu nucleus is further split into 11 components due to the combined effect of the nitrogens ($I = 1$) and the protons ($I = 1/2$ giving rise to 44 peaks). The electron is delocalized on the ligand, which is good evidence for covalent bonding (25)

Several of the factors affecting line width in ESR are analogous to those in NMR. They include the following (2).

1) Spin-spin interaction is due to the mutual effect of one paramagnetic ion on the other. Each ion is regarded as a magnetic dipole which alters the magnetic field at the neighboring paramagnetic sites. (This effect can be reduced by moving the paramagnetic sites further apart, i.e. by diluting the salt with an isomorphous, diamagnetic equivalent). Similar effects can be produced by the magnetic dipole interaction with the protons of water molecules. For that reason sometimes crystals are grown from heavy water (13) because the deuterons have a magnetic dipole moment only about $1/3$ of that of the protons.

2) Spin-lattice relaxation, occurs when the paramagnetic ions interact with the thermal vibrations of the lattice through spin-orbit coupling. Many measurements of relaxation time, however, give results inconsistent with this mechanism. Contrary to theory, it is found that relaxation times often do not vary inversely proportional to temperature and field strength and that they are sometimes dependent on crystal size (19). A lattice-bath relaxation time was proposed which represents the time required for the dissipation of energy to the crystal boundaries by the lattice modes.

3) Power saturation broadening occurs when the incident power is so large that the normal relaxation processes can no longer restore thermal equilibrium. In solid state ESR this effect is usually negligible.

4) Inhomogeneous fields sometimes lead to broadening. In addition there is the effect of exchange which can easily be reduced by dilution of the crystals. The phenomenon of exchange is discussed more fully below.

$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ contains two types of Cu^{2+} ions per unit cell and hence we would expect to observe a peak corresponding to each. However at wavelengths shorter than about 8 cm. and at certain directions of the magnetic field only one peak is observed; in addition the peak is narrower than expected (10). Both phenomena are due to exchange interactions. When the external field H is along the tetragonal axis of one of the copper ions in a single crystal, the values of g , and therefore the precessional frequencies of the two ions are different. If there is an exchange between the ions, the absorption curve will depend on the magnitude of the exchange frequency relative to those of the precessional frequencies. When the former is large, a mean value of the energy of the copper ions is measured, and when it is smaller 2 lines appear (10, 20). The width of the line is a result of the precession of the magnetic moment which is under the influence of magnetic interactions which in turn are affected by non-magnetic effects such as exchange. When such atomic or electronic motions are rapid, the broadening effect of the magnetic interactions is reduced and "exchange narrowing" of the lines occurs (9).

Hydrated cupric acetate also contains two distinct ions per unit cell and one would therefore expect two lines each split into a quartet by interaction with the Cu nucleus. Instead a triply degenerate spin state results from exchange coupling, and one peak which is split into a septet due to the total nuclear spin of 3 is obtained (14). This is also observed in the propionate and butyrate but not in the formate (5). Solutions of copper complexes of monoethanolamine in alcohols give a spectrum of four hyperfine components as expected and in addition an exchange line superimposed, indicative of dimerization (28).

Solutions of $\text{Cu}(\text{acac})_2$ (30) and $\text{Cu}(\text{RR}'\text{NCS}_2)_2$ (27) as well as other complexes give four lines of equal area but with line widths increasing with increasing I . These observations suggest that if the paramagnetic relaxation time (spin-spin as well as spin-lattice) depends on the nuclear orientation the hyperfine interactions contribute to the paramagnetic relaxation (26). McConnell (26) has derived expressions for the relaxation times which give reasonable agreement with experimental values. Recent work (29) has shown that the asymmetry of the spectrum was augmented with increase in viscosity and ν , the resonance frequency, in direct contradiction with McConnell's theory.

Only when the applied field is much stronger than the magnetic field due to the surroundings of the ion, are the hyperfine splittings equally spaced. When the two become of comparable magnitude the hyperfine levels will contain admixtures of the different quantum states M_I , which will give rise to unequal spacings. If the nucleus has a quadrupole moment additional peaks are observed; this effect is greatest when the applied field is at right angles to the crystalline field because both are trying to quantize the nucleus along their own axis. The result is that the selection rule $M_I = 0$ breaks down and sometimes nuclear quadrupole transitions are observed in addition to unequal spacing of the lines. When the effect of the quadrupole moment is comparable in magnitude to the nuclear interaction, 6I-1 lines result (11) as in the case of $(Zn, Cu)(NH_4)_2(SO_4)_2 \cdot 6H_2O$ (21). An example of transitions forbidden by the $M_I = 0$ selection rule is $CuK_2(SO_4)_2 \cdot 6D_2O$ (13).


If a paramagnetic ion is located in a perfectly octahedral crystal site, the g-value is independent of the orientation of the single crystal in the magnetic field and is said to be isotropic. When the crystal field is tetragonal the g-value in the Z direction, g_Z , is different from the values in the X and Y directions g_X and g_Y and the g-values are anisotropic. The Z direction is defined as the highest-fold rotation axis which must be determined by x-ray methods. In a tetragonal field $g_Z \equiv g_{||}$, the g-value with the external field parallel to the Z axis and $g_X = g_Y \equiv g_{\perp}$, the g-value with the field parallel to the XY plane. If θ is the angle between the magnetic field and the z-axis, the following relationship holds: $g^2 = g_{||}^2 \cos^2 \theta + g_{\perp}^2 \sin^2 \theta$. If further distortion takes place, such as a rhombic one, g_X is no longer equal to g_Y and g_{\perp} is without meaning. Small symmetry distortions which remain undetected by x-ray methods can thus be determined.

Taking into account the nuclear and quadrupole moment interactions and the anisotropy of the g-values, the Hamiltonian for an ion in a tetragonal environment becomes (excluding $W_F + V + W_{is}$) (6):

$$W = D[S_Z^2 - 1/3 S(S+1)] + g_{||} H_Z S_Z + g_{\perp} (H_X S_Y + H_Y S_X) + A S_Z I_Z + B(S_X I_X + S_Y I_Y) + Q[(I_Z^2 - 1/3 I(I+1))] - \chi_N^2 H I$$

when S_X , S_Y and S_Z and I_X , I_Y and I_Z are the X, Y, Z components of the angular momenta of the electron and nucleus resp. D is the spin-spin splitting constant, A and B are the hyperfine splitting constants and Q the quadrupole constant; the term in D drops out for Cu^{2+} because $S_Z^2 = 1/4$ and $S = 1/2$; the last term is usually negligible. Given enough information the constants can, at least in principle, be evaluated. They can be calculated from experimental data but their signs are often difficult to determine. Some typical values for $CuSO_4 \cdot K_2SO_4 \cdot 6H_2O$ are (7): $g_{||} = 2.4$ $g_{\perp} = 2.1$

$$|A| = 0.013 \text{ cm}^{-1} \quad |B| = 3.5 \times 10^{-3} \text{ cm}^{-1} \quad |Q| = 0.9 \times 10^{-3} \text{ cm}^{-1}$$



The expression for the g-values (see figure) are:

$$g_{\parallel} = 2 - \frac{8\lambda}{E_2 - E_0} \qquad g_{\perp} = 2 - \frac{2\lambda}{E_3 - E_0}$$

where λ is the spin-orbit coupling constant and the values for the energy differences are obtained from the absorption spectrum. From experiment it is found that $g_x \neq g_y$, which indicates that there is a small rhombic distortion. If a value is estimated for $E_3'' - E_3'$, the agreement with experiment is reasonable (7) but it can be improved by taking values for λ different from the free ion value determined from atomic spectra: $\lambda = 828.7 \text{ cm}^{-1}$. For Cu^{2+} in water $\lambda = 710 \text{ cm}^{-1}$ (15) and for $\text{Cu}(\text{NH}_3)_4\text{SO}_4 \cdot \text{H}_2\text{O}$, $\lambda = 450 \text{ cm}^{-1}$ (18) as a rough estimate.

The calculated values for the hyperfine splitting constants A and B are in complete disagreement with experiment. It is known from optical spectroscopy that configurations with unpaired s electrons have a wider hyperfine structure. A new wavefunction is assumed which has admixtures of the configuration $[3S, 4S(^3S), 3d^9(^2D)]^2D$ expressed in terms of κ , which is a numerical factor expressing the admixture of configurations with unpaired s electrons (6, 7, 8). For $\text{CuSiF}_6 \cdot 6\text{H}_2\text{O}$ a value of $\kappa = 0.30$ gives good agreement with experiment; this corresponds to a probability of only 4×10^{-5} for the configuration with the unpaired s-electron (6). The observation of forbidden transitions in later experiments with Cu Tutton salts made possible the determination of the signs of the hyperfine splitting constants and very accurate values for the quadrupole moments could be obtained (16).

For $(\text{Cu}, \text{Zn}) \text{SiF}_6 \cdot 6\text{H}_2\text{O}$ the spectrum above 90°K consists of one band with partly resolved hyperfine structure. In the region of $90-20^\circ\text{K}$ a transition takes place and at 20°K the spectrum consists of 3 sets of lines (12). The trigonal symmetry of the crystal leaves the ground state doubly degenerate. There are three

possible tetragonal Jahn-Teller distortions along the principal axes which lead to the same lowering in energy. The system may resonate among these three structures which leads to nearly isotropic g-values. At lower temperature the system no longer resonates and three lines with anisotropic g-values are observed(17).

In copper chelate complexes such as coppersalicylaldehyde and $\text{Cu}(\text{acac})_2$, appreciable covalent bonding occurs. The theory for the nearly ionic case discussed above is no longer valid and a molecular orbital model has to be used. The degree of mixing of metal orbitals and ligand orbitals is expressed by the constants α , β and γ , with values ranging from 0.5 for the completely covalent to 1.0 for the completely ionic case. α describes the σ -bonding and β , and γ the "in plane" and "out of plane" π bonding resp., the plane being that of the Cu nucleus and the chelate ring (24). For coppersalicylaldehyde the values are $\alpha^2 = 0.83$, $\beta^2 \geq 0.91$ and $\gamma^2 = 0.72$; the degree of π bonding in the plane is considerably greater than that out of the plane (25) Kivelson and Neiman(22) have summarized values for α and β , for a large number of Cu chelates. The values for α^2 (0.75-0.90) indicate a large degree of covalent σ -bonding.

It is clear that a great deal of information can be gained from ESR experiments with regard to structure and bonding; many other examples could be cited. In order to carry out studies on single crystals, for which case the theory has been worked out in the greatest detail, one needs to know the crystal structure in order to know the orientation of the crystal in the field.

References

General

1. W. Gordy, W. V. Smith and R. T. Trambarulo, "Microwave Spectroscopy," John Wiley and Sons, Inc. New York (1953).
2. D. J. E. Ingram, "Spectroscopy at Radio and Microwave Frequencies," Butterworth, London (1955).
3. W. Low, "Paramagnetic Resonance in Solids," Academic Press, New York (1960).
4. J. E. Wertz. Chem. Revs. 55, 829 (1955).

Specific

5. H. Abe, Phys. Rev. 92, 1572 (1953); J. Phys. Soc. Japan 13, 987 (1958).
6. A. Abragam and M. H. L. Pryce, Proc. Roy. Soc. A205, 135 (1951).
7. A. Abragam and M. H. L. Pryce, Proc. Roy. Soc. A206, 164 (1951).
8. A. Abragam, J. Horowitz and M. H. L. Pryce, Proc. Roy. Soc. A230, 169 (1955).
9. P. W. Anderson, J. Phys. Soc. Japan 9, 316 (1954).
10. D. M. S. Bagguley and J. H. E. Griffiths, Nature 162, 538(1948).
11. B. Bleaney, Physica 17, 175 (1951).
12. B. Bleaney and D. J. E. Ingram, Proc. Phys. Soc. 63, 408(1950); A. Abragam and M. H. L. Pryce, ibid., 63, 409 (1950).
13. B. Bleaney, K. D. Bowers and D. J. E. Ingram, Proc. Phys. Soc. A64, 758 (1951).

14. B. Bleaney and K. D. Bowers, Proc. Roy. Soc. A214, 451 (1952).
15. B. Bleaney, K. D. Bowers and M. H. L. Fryce, Proc. Roy. Soc. A218, 166 (1955).
16. B. Bleaney, K. D. Bowers and D. J. E. Ingram, Proc. Roy. Soc. A228, 147 (1955).
17. B. Bleaney, K. D. Bowers and R. S. Trenam, Proc. Roy. Soc. A228, 157 (1955).
18. E. H. Carlson and R. D. Spence, J. Chem. Phys. 24, 471 (1956).
19. J. A. Giordmaine, L. E. Alsop, F. R. Nash and C. H. Townes, Phys. Rev. 109, 302 (1958).
20. C. J. Gorter and J. H. Van Vleck, Phys. Rev. 72, 1128 (1947).
21. D. J. E. Ingram, Proc. Phys. Soc. A62, 664 (1949).
22. D. Kivelson and R. Neiman, J. Chem. Phys. 35, 149 (1961).
23. H. A. Kramers, Proc. Amsterdam Acad. Sci. 33, 959 (1930).
24. A. H. Maki and B. R. McGarvey, J. Chem. Phys. 29, 31 (1958).
25. A. H. Maki and B. R. McGarvey, J. Chem. Phys. 29, 35 (1958).
26. H. M. McConnell, J. Chem. Phys. 25, 709 (1956).
27. B. R. McGarvey, J. Phys. Chem. 60, 71 (1956).
28. A. I. Riokind, Doklady Akad. Nauk. SSSR 135, 365 (1960)
cf. CA. 56, 1082f (1962).
29. A. I. Rivkind, Paramagnitn. Rezonans. Kazansk, Univ. Sbornik, 50 (1960) cf. C. A. 56, 8192b (1962); Zhur. Fiz. Khim., 35, 2099 (1961), cf. C.A. 56, 1077b (1962).
30. T. Vänngard and S. Akerstrom. Nature 184, 183 (1959).
31. R. P. Penrose, Nature 163, 992 (1949).

N.M.R. IN THE PRESENCE OF PARAMAGNETIC IONS

Brad Wayland

April 16, 1963

Paramagnetic species in the vicinity of a magnetic nucleus cause both relaxation and chemical shift effects. Relaxation studies have been most effective in yielding information on water exchange rates in dilute solutions of paramagnetic metal ions. Chemical shift effects have been used primarily in studying electron delocalization in paramagnetic complexes of conjugated ligands.

I. EFFECTS OF PARAMAGNETIC IONS ON THE RELAXATION TIMES OF MAGNETIC NUCLEI IN SOLUTION

Relaxation times of water protons in dilute aqueous solutions of paramagnetic ions have been extensively studied(1, 2). The addition of paramagnetic ions to water shortens the proton longitudinal (spin lattice) and the transverse (spin spin) relaxation times, T_1 and T_2 (3). The most successful correlation of experimental results has been achieved through consideration of both proton spin-electron spin dipole dipole interactions as formulated by Solomon (4), and proton spin-electron spin exchange coupling as suggested by Bloembergen (5). The dipolar term arises from the fluctuating magnetic field created by the tumbling paramagnetic ion. The magnetic moment of an unpaired electron is $\sim 10^3$ times larger than the nuclear magnetic moment and thus the electron-nuclear dipole term dominates over the nuclear-nuclear dipole term. If the unpaired electron has a wave function which has a finite probability at the magnetic nucleus then the Hamiltonian contains an electron spin-nuclear spin coupling term of the form $AI \cdot S$. This mechanism may also produce a fluctuating magnetic field at the nucleus and shorten T_1 . Also, if the lifetime of the paramagnetic state is long then a distribution of fields results which shortens T_2 , but has little effect on T_1 .

The electron spin-nuclear spin dipolar and exchange coupling interactions are given by the Hamiltonians (1) and (2) respectively:(6)

$$(1) H_1 = \frac{h^2 \gamma_I \gamma_S}{r^3} [3(I \cdot r)(S \cdot r) - I \cdot S]$$

$$(2) H_2 = AI \cdot S$$

The combined Solomon-Bloembergen equations for T_1 and T_2 appear in a simplified form as equations 3 and 4.

$$(3) \frac{1}{T_1} = \left(\frac{1}{T_1} \right)_{dip} + \left(\frac{1}{T_1} \right)_{ex} = \frac{4}{30} \frac{S(S+1)g^2 \gamma_I^2 \gamma_S^2 P}{r^6} \left[3\tau_c + \frac{7\tau_c}{1 + S^2 \tau_c^2} \right]$$

$$\frac{2}{3} \frac{S(S+1)A^2 P}{h^2} \frac{\tau_e}{1 + S^2 \tau_e^2}$$

$$(4) \frac{1}{T_2} = \frac{1}{T_2} \text{dip} + \left[\frac{1}{T_2} \text{ex.} = \frac{4}{60} \frac{S(S+1)g^2 \gamma_I^2 \mu^2 P}{r^6} \left[7\tau_c + \frac{13\tau_c}{1+\frac{1}{2}S^2\tau_c^2} \right] + \frac{1}{3} \frac{S(S+1)A^2 P}{h^2} \left[\tau_e + \frac{\tau_e}{1+\frac{1}{2}S^2\tau_e^2} \right] \right]$$

γ_I = gyromagnetic ratio μ = Bohr magneton

A = coupling constant for exchange interaction

P = probability of finding a given proton in the hydration sphere of the paramagnetic ion and is proportional to the metal ion concentration

τ_c = correlation time for dipolar relaxation

τ_e = a measure of the lifetime of the paramagnetic state

ω_S = Larmor frequency for the electron

$$(5) \frac{1}{\tau_e} = \frac{1}{\tau_S} + \frac{1}{\tau_h} \quad \tau_S = \text{electron relaxation time - a measure of the lifetime of an electron spin state}$$

τ_h = chemical exchange time for protons in the solvation sphere

$$(6) \frac{1}{\tau_c} = \frac{1}{\tau_s} + \frac{1}{\tau_r} \quad \tau_r = \text{correlation time for tumbling of the hydrated magnetic ion.}$$

The important features to recognize in equations (3) and (4) are;

Case 1: If $\omega_S^2 \tau_e^2 \ll 1$ (i.e. τ_S is very short) then $T_1 = T_2$ and T_1 may be relatively long ($\sim 10^{-1} - 10^{-3}$ sec) so that the N.M.R. spectra is not appreciably broadened.

Case 2: If $\omega_S^2 \tau_e^2 \gg 1$ (i.e. τ_S is very long) then $T_1 \neq T_2$ and the NMR spectra is extensively broadened due to exchange coupling.

Most paramagnetic metal ions have short electron spin relaxation times ($\tau_S \approx 10^{-10} - 10^{-12}$ sec.) in aqueous solutions and belong to case 1. The ions Mn^{+2} , Gd^{+3} and Fe^{+3} with half filled shells (S states) have relatively long spin relaxation times ($\tau_S \approx 10^{-8} - 10^{-9}$ sec) because spin orbit coupling and distortion of the hydration sphere are minimized and are examples of case 2 (5, 7). However, when Mn^{+2} is complexed by EDTA in aqueous solution τ_S is appreciably shortened and $T_1 = T_2$ (8, 9). Thus Mn^{+2} when complexed with EDTA becomes an example of case 1.

Connick (1), following the proton experiments of Gutowsky et. al., (6) has used temperature dependence of T_2 in studying the water exchange rates of paramagnetic ions by O^{17} nuclear resonance. The general expression for T_2 given by equation 7 was achieved

by analysis of the Bloch equations modified for chemical exchange between a diamagnetic and a paramagnetic site (1, 10, 11).

$$(7) \quad \frac{1}{T_{2P}} = \frac{1}{T_2} - \frac{1}{T_{2A}} = \frac{1}{T_A} \left[\frac{T_{2B}^{-2} + T_{2B}^{-1} \tau_B^{-1} + \Delta\omega_B^2}{(T_{2B}^{-1} + \tau_B^{-1})^2 + \Delta\omega_B^2} \right]$$

- T_{2P} \equiv transverse relaxation due to the paramagnetic ion
- T_2 \equiv transverse relaxation time
- τ_A \equiv lifetime of the magnetic nucleus in the diamagnetic site
- τ_B \equiv lifetime of the magnetic nucleus in the paramagnetic site
- T_{2A} \equiv transverse relaxation time in the diamagnetic site
- T_{2B} \equiv transverse relaxation time in the paramagnetic site
- $\Delta\omega_B$ \equiv difference in the Larmor frequency between the diamagnetic and paramagnetic sites

The temperature dependence of equation 7 was investigated for the following limiting cases:

$$(7A) \quad \Delta\omega_B^2 \ll \frac{1}{T_{2B}}, \quad \frac{1}{\tau_B}$$

Equation 7 becomes:

$$\frac{1}{T_{2P}} = \frac{P_B}{\tau_B} = \frac{1}{\tau_A}$$

Relaxation occurs through a change in the precessional frequency and is rapid. $\frac{1}{T_{2P}}$ is controlled by the rate of water exchange.

$$(7B) \quad \frac{1}{\tau_B^2} \ll \Delta\omega_B^2 \ll \frac{1}{T_{2B} B}$$

$$\frac{1}{T_{2P}} = P_B \tau_B \Delta\omega_B^2$$

Chemical exchange is rapid. $\frac{1}{T_{2P}}$ is controlled by the rate of relaxation through change in precessional frequency.

$$(7C) \quad \frac{1}{T_{2B}^2} \gg \Delta\omega_B^2, \quad \frac{1}{\tau_B^2}$$

$$\frac{1}{T_{2P}} = \frac{P_B}{\tau_B} = \frac{1}{\tau_A}$$

Relaxation by T_{2B} is fast. $\frac{1}{T_{2P}}$ is controlled by the rate of water exchange.

$$(7D) \frac{1}{T_{2B} T_B} \gg \frac{1}{T_{2B}^2}, \quad \omega B^2$$

$$\frac{1}{T_{2P}} = \frac{P_B}{T_{2B}}$$

Chemical exchange is rapid. $\frac{1}{T_{2P}}$ is controlled by the T_{2B} relaxation process.

A detailed analysis of the temperature dependence of T_2 for aqueous solutions of Mn^{+2} will be given to demonstrate the evaluation of water exchange rates.




FIGURE 1




FIGURE 2

From figure 1 it is seen that $\frac{T_1}{T_2} \gg 1$ and thus $(\omega S^2 / e^2) \gg 1$. Equation 4 reduces to the form:

$$(8) \quad \frac{1}{T_{2B}} = C r_e \quad \frac{1}{r_e} = \frac{1}{r_s} + \frac{1}{r_B}$$

At high temperatures T_{2B} is controlled by chemical exchange (r_B), while at lower temperatures the electronic interruption (r_s) will dominate T_{2B} . $\frac{1}{r_B}$ is the rate of a chemical reaction and should follow an exponential temperature dependence.

$$(9) \quad \frac{1}{r_B} = \left(\frac{kT}{h} \right)^{-1} e^{(\Delta H^\ddagger / RT - \Delta S^\ddagger / R)}$$

The temperature dependences of ΔH^\ddagger and ΔS^\ddagger are given by equations 10 and 11.

$$(10) \frac{\Delta G}{RT} = \frac{4}{9} \frac{A \gamma_e}{r_P} \frac{I(I+1)(S)S+1}{kT}$$

$$(11) \frac{1}{T_S} = \frac{12 C_2^2}{5h^2 S(S+1)} \frac{\tau_v}{1 + \frac{S^2}{S^2} \tau_v^2} + \frac{4 \tau_v}{1 + \frac{S^2}{S^2} \tau_v^2}$$

τ_v = correlation time for collision of water molecules with the hydrated ion.

$$\tau_v = \tau_v^0 e^{-v/RT}$$

The plot of $\log T_{2P}$ versus $\frac{1}{T^\circ K}$ given in figure 2 reveals that

at low temperatures T_{2P} is controlled by chemical exchange corresponding to equation 7A or 7C. An estimate of τ_B from the low temperature slope eliminated the possibility of τ_B effecting the relaxation because an impossibly high coupling constant would be necessary to explain the data. When ΔG is very small equation 7 reduces to equation 12.

$$(12) \frac{1}{T_{2P}} = \frac{(\tau_A T_{2B})^{-1}}{T_{2B}^{-1} + \tau_B^{-1}} = \frac{P_B}{\tau_B + T_{2B}} = \frac{P_B}{\tau_B} = \frac{1}{C} \left(\frac{1}{\tau_S} + \frac{1}{\tau_B} \right)$$

$$\frac{\tau_A}{\tau_B} = \frac{P_A}{P_B} \quad P_A \cong 1 \quad \tau_A = \frac{P_A}{P_B} \tau_B = \frac{\tau_B}{P_B}$$

The solid curve in figure 2 which passes through the experimental points was calculated as the sum of three lines derived for the cases of high, low and intermediate temperature. At low temperature $\tau_B \gg \frac{1}{C} \left(\frac{1}{\tau_S} + \frac{1}{\tau_B} \right)$ and $T_{2P} \cong \frac{\tau_B}{P_B}$. At high temperatures

$\frac{1}{\tau_B} \gg \frac{1}{\tau_S}$ and τ_B and $T_{2P} = \frac{1}{CP_B \tau_S}$. At intermediate temperatures τ_S contributes appreciably to T_{2P} and $T_{2P} = \frac{1}{CP_B \tau_S}$. By using a curve fitting process parameters for the exchange process were determined (Table 1).

TABLE 1: Values for the exchange of H₂O from the first coordination sphere of Mn²⁺ ion.

	0 ¹⁷	proton(6)	proton(2)
ΔH^\ddagger (k cal)	8.1	7.8	7.5
k_1 (sec ⁻¹ at 298°K)	3.1 x 10 ⁷	3.6 x 10 ⁷	4.0 x 10 ⁷
τ_S (sec at 298°K)	3.8 x 10 ⁻⁹	3.0 x 10 ⁻⁹	3.5 x 10 ⁻⁹
A/h (c.p.s)	2.7 x 10 ⁶	1 x 10 ⁶	1.0 x 10 ⁶
ΔS^\ddagger (eu)	2.9	2.2	1.3

Connick has also determined water exchange parameters for Cu^{+2} , Fe^{+2} , Ni^{+2} and Co^{+2} in dilute aqueous solution (Table 2). The temperature dependence of T_{2p} takes several different forms for this series of metal ions. In each case, however, a term involving water exchange was involved which allowed exchange data to be determined. Similar exchange experiments have been carried out on exchange of NH_3 and Cl^- in aqueous solution by observing the N^{14} and $\text{Cl}^{35,37}$ nuclear resonance (12, 13).

TABLE 2: Values for exchange of H_2O from the first coordination sphere of Cu^{+2} , Mn^{+2} , Fe^{+2} , Co^{+2} and Ni^{+2}

	Cu^{+2}	Mn^{+2}	Fe^{+2}	Co^{+2}	Ni^{+2}
ΔH^\ddagger (k.cal.)	11	8.1	7.7	8.0	11.6
k_1 (sec^{-1})	1.0×10^4	3.1×10^7	3.2×10^6	1.13×10^6	2.7×10^4
τ_S (sec)		3.8×10^{-9}	1.4×10^{-11}	5×10^{-12}	1.3×10^{-11}
A/h (c.p.s.)	5.5×10^6	2.7×10^6	9.8×10^5	1.14×10^6	3.6×10^6
ΔS^\ddagger (eu)	-4.0	2.9	-3.0	-4.1	0.6

II. PROTON CHEMICAL SHIFTS AND CARBON SPIN DENSITIES OF CONJUGATED LIGANDS BONDED TO PARAMAGNETIC IONS

Large proton chemical shifts have been reported for some paramagnetic acetylacetonates (14), metallocenes(15-17), and for a wide variety of Ni(II) chelates of N,N' disubstituted amino-troponeimines(18-23). Commonly the proton shifts are in the range from +5000 c.p.s. to -5000 c.p.s. using a 40 Mc. probe. These shifts are a full order of magnitude larger than shifts observed for protons in diamagnetic species.

These large contact proton NMR shifts can be visualized as resulting from the unpairing of ligand electron pairs due to charge transfer to the paramagnetic metal ion. The effects of the unpaired electron in the ligand π orbitals must be transferred to the protons via the Fermi Contact term (24), for the dipolar interactions average to zero due to rapid molecular tumbling in solution. The observed nuclear spin-electron spin coupling must arise from π - σ exchange coupling (appendix I) in order to place a finite odd electron density at the in plane aromatic protons(25,26). In a system characterized by a very short electron spin relaxation time or a short exchange time, the proton N sees a single average hyperfine magnetic field corresponding to the effective spin Hamiltonian for proton N, where $\langle S_z \rangle$ is the time average value of the Z component of electron spin(27). Since μ_N is positive for the

$$(13) \quad H = -\mu_N N_Z \left[H_0 - 2\pi A_N \frac{Ih}{h} \langle S_z \rangle \right]$$

$$(14) \quad \langle S_z \rangle = \frac{-g\beta S(S+1)H_0}{3kT}$$

3kT

$$(15) \quad H = \gamma_N Z H_0 \left[1 - 2 A_N \frac{\gamma_e}{\gamma_N} \frac{g \beta^2 S(S+1)}{3kT} \right]$$

$\gamma_N Z$ \equiv Z component of the magnetic moment for proton N

A_N \equiv electron spin-nuclear spin contact interaction constant
(coupling constant)

β \equiv Bohr magneton

proton equation 15 predicts that the contact shift will be to lower fields when A_N is positive and to higher fields when A_N is negative. It is apparent that N.M.R. offers a direct means of determining the sign of the electron spin-nuclear spin coupling constant.

Bloembergen (5) has explicitly formulated the relationship between the nuclear resonance contact shift and the scalar coupling constant (A_N) as equation 16.

$$(16) \quad \frac{\Delta H}{H} = \frac{\Delta \nu}{\nu} = - \frac{4A_N}{\gamma_N} \frac{\gamma_e}{\gamma_N} \frac{g \beta^2 I(I+1) S(S+1)}{9kT}$$

Equation 17 is for the specific case of proton magnetic resonance.

$$(17) \quad \frac{\Delta H}{H} = \frac{\Delta \nu}{\nu} = - \frac{1}{3} A_N \frac{\gamma_e}{\gamma_N} \frac{g \beta^2 S(S+1)}{kT}$$

γ_e \equiv magnetogyric ratio for the electron

γ_N \equiv " " " " magnetic nucleus

I \equiv nuclear spin

S \equiv electron spin multiplicity

ΔH \equiv $H_{\text{complex}} - H_{\text{ligand}}$

Contact interaction constants (A_N) are related to the unpaired spin densities on $p\pi$ orbitals of SP^2 carbons by equation 18 (28, 29, 30).

$$(18) \quad A_N = Q \rho_N$$

ρ_N \equiv spin density at carbon
 Q = proportionality constant
 (-22.5 Gauss for an aromatic C-H fragment).

Spin density (ρ_N) is defined as that fraction of an unpaired electron which appears to be localized at site N. Spin densities may be either positive or negative. A negative spin density results when the electron spin at site N is antiparallel to the net spin of the paramagnetic species.

REQUIREMENTS FOR OBSERVATION OF A HYPERFINE CONTACT SHIFT

Conventionally spin density distributions have been obtained from the isotropic nucleus-electron hyperfine splittings that appear in EPR spectroscopy. Under special conditions spin densities are manifested in the NMR by large chemical shifts. NMR contact shifts have the advantage over EPR of offering a means of determining both the magnitude and the sign of the coupling constant and spin density. The conditions requisite to observing a contact shift in the NMR are that either of the inequalities (19) are satisfied(27).

$$(19) \quad \frac{1}{\tau_S} \gg A \quad \frac{1}{\tau_e} \gg A$$

τ_S = electronic relaxation time
 τ_e = chemical exchange time
 A = contact interaction constant (coupling constant)

It is interesting to note that the conditions requisite to observing a sharp nuclear resonance spectrum imply the absence of observable hyperfine splittings for the same system in the EPR. If $\frac{1}{\tau_e} \gg A$, the hyperfine splittings are obscured by exchange narrowing, and if $\frac{1}{\tau_S} \gg A$ then the paramagnetic resonance line widths are greater than the splittings.

MECHANISM FOR PLACING AN UNPAIRED ELECTRON ON THE LIGAND BONDED TO A PARAMAGNETIC ION

The transference of effects of unpaired electrons on the metal ion to the ligand can be qualitatively explained by charge transfer from the ligand to the metal ion (16, 17, 31). The spin of the electron transferred is such that the multiplicity of the next lower oxidation state of the metal ion is achieved. The unpaired electron remaining on the ligand may thus have a spin parallel with or antiparallel to the net spin of the metal ion. Antiparallel spin on the ligand results most frequently for ions whose next lowest oxidation state has higher multiplicity. Spin parallel to the metal ion results when the next lower oxidation state has lower multiplicity. Before offering evidence for this mechanism, it is necessary to know that the unpaired spin density at the proton is antiparallel to the spin density at the carbon atom to which it is bonded. This effect arises from s-p exchange interaction discussed more fully in appendix 1 (25-27).

The proton magnetic resonances of the bis-cyclopentadienyl compounds of V^{+2} , Cr^{+2} , Mn^{+2} , Co^{+2} and Ni^{+2} have been recorded(15-17). The vanadium and chromium compounds give proton resonance shifts to lower fields while the protons in the manganese, cobalt and nickel compounds are shifted to higher fields. $V(C_5H_5)_2$ will be considered in more detail in order to illustrate the mechanism for transferring unpaired spin to the ligand. Figure 3 is a molecular orbital diagram for metallocene compounds. The electrons

are assigned for the case of vanadocene.





FIGURE 3

The most important charge-transfer process involves electrons passing from the filled e_{1g} ring orbitals into the empty e_{1g} d orbital of the vanadium ion. The electron transferred has spin parallel to the spin of the metal ion, for this leads to a favorable exchange energy. The electron transfer process essentially produces V^+ ion with electron configuration $(a_{1g})^1(e_{2g})^1(e_{-2g})^1(e_{1g})^1$. The unpaired electron remaining on the ligand has spin antiparallel to the metal ion which places parallel unpaired spin at the proton. This mechanism predicts a proton shift to lower fields in agreement with experimental results. The Ni^{+2} ion in nickelocene is assigned the electronic configuration $(e_{2g})^4(a_{1g})^2(e_{1g})^2$. All of the d orbitals are either filled or half filled. The electron transferred from the ligand must have spin antiparallel to the d electrons. This process predicts a high field shift for the ligand protons in agreement with the observed shift.

In the above metallocene compounds the carbon spin densities in any one compound are found to have the same sign, for all of the carbons and protons are effectively equivalent. In ligands of lower symmetry where there are preferred positions for the unpaired electron, the spin density on adjacent atoms is found to alternate in sign. The preferred positions have spin parallel to the electron remaining on the ligand after charge transfer, while adjacent positions have unpaired spin antiparallel to the net spin of the ligand. This effect arises from $\pi - \pi$ configuration interaction (21,29) (Appendix 2). The preferred positions for the unpaired electron can

be predicted by considering the possible valence bond structures. Ni(II) N, N' diethylaminotroponimineate serves as a good example(21).



Proton magnetic resonance spectra of Ni II N, N'diethylaminotroponimineate

FIGURE 4

Valence bond structures (Figure 5) place the unpaired electron at the δ and ϵ positions of the seven membered ring. No structure placing the odd electron at the ζ position can be written without recourse to long bonds. As in the case of nicklocene, the ligand



FIGURE 5

as a whole has spin parallel to the Ni ion. Positive spin densities are thus predicted for the δ and ϵ carbons and a negative spin density is expected for the ζ carbon. Remembering that the spin density at the proton is antiparallel to that on the carbon, the δ and ϵ protons are expected to be shielded while the ζ proton should be deshielded. These considerations lead immediately to the NMR assignment in Figure 4. Confirmation of the assignment in Figure 4 has been made by other independent means. Substitution of phenyl for ethyl on the nitrogens of the previous compound extends the π system over which the unpaired spin is distributed. As expected from simple valence bond consideration, (Figure 6) positive spin densities occur at the ortho and para positions and a negative spin density is found at the meta position.



FIGURE 6

Valence bond calculations of spin densities by the method of McConnell(32) have been useful for predicting spin densities at atoms not readily observable in the NMR. The calculations are made assuming the ligand to be a free radical. The values obtained are then scaled down to correspond to the transfer of that fraction of an electron which best agrees with the observed spin densities. Example results given in Figure 7 are in amazingly good agreement with experiment. (21)




FIGURE 7

It is apparent that NMR spin density determinations provide a promising new approach to the study of conjugation in molecules containing extended π electron systems. The ability of NH, O, S, and $-SO_2$ groups to transfer electronic effects has been directly studied using this technique (21). The importance of F double bonding with π systems has also been assessed(20).

Contact shifts due to paramagnetic ions are not limited to conjugated ligands. Fluorine NMR of paramagnetic crystals has demonstrated the presence of large internal magnetic fields at the fluorine nucleus (33-36). Isotropic hyperfine interactions arise from the unpairing of 2S electrons. A smaller anisotropic effect is caused by the unpairing of 2P electrons. Single crystal fluorine NMR of $KMnF_3$ has yielded information on antiferromagnetic ordering in the solid state (37). O^{17} NMR of aqueous solutions of rare earth perchlorates has been reported (38). The observed magnitude and variation of the O^{17} contact shifts with the number of 4f electrons was interpreted as evidence for very weak covalent bonding between O^{17} 2S and 2P orbitals with the rare earth 6S orbitals. The 4f, 5d, and 6p orbitals play at most a minor role in the bonding.

APPENDIX 1

π - σ Transmission of Spin Polarization Through Exchange Coupling in the C-H Fragment

The theory of π - σ exchange coupling has been worked out in order to explain the presence and direction of spin polarization at the protons in paramagnetic complexes of conjugated ligands (25, 26). An orthogonal basis set was chosen for the calculation. If the unpaired electron is placed in a π M.O. of the aromatic ligand, then there is zero electron density at the inplane aromatic protons. An excited electronic configuration must be used to give rise to the

indirect coupling. The lowest energy excited configuration which gives the proper result is the excitation of one electron from a σ_B bonding orbital to a σ^* antibonding orbital. Linear combinations of Slater determinants are made in order to produce proper eigenfunctions for the possible excited states. The doublet excited state corresponding to a triplet $\sigma_B \sigma_A^*$ with the π electron opposite in spin leads the proper interaction. The excited state which mixes with the ground state always places spin polarization at the proton which is antiparallel to the spin of the π electron. An alternate and probably more realistic approach to this problem can be made by simply assuming that the π and σ orbitals are not orthogonal. Spin exchange may thus directly occur which places spin density at the proton antiparallel to the unpaired electron in the π system.

APPENDIX 2

$\pi - \pi$ Configuration Interaction

Negative spin density is a concept foreign to molecular orbital theory. M. O. theory must be modified by addition of $\pi - \pi$ configuration interaction in order to account for negative spin densities. Configuration interaction is the unpairing of electron spins so as to reduce the energy of the system through exchange of electrons having the same spin. The allyl free radical will be briefly discussed in order to demonstrate the effects of configuration interaction on the spin densities of conjugated radicals (21).



The odd electron in ψ_2 gives rise to positive spin densities at carbons 1 and 3. Exchange interactions between the odd electron in ψ_2 of spin \uparrow and the paired electron in ψ_1 of spin $\uparrow\downarrow$ enhances the positive spin density at C_1 and C_3 and leaves a net spin \downarrow (negative spin density) at C_2 . This effect is also responsible for producing negative spin densities near the nucleus of many paramagnetic metal ions (35).

REFERENCES

I. Relaxation Effects

1. T. J. Swift and R. C. Connick, J. Chem. Phys., 37, 307 (1962).
2. N. Bloembergen and L. O. Morgan, J. Chem. Phys., 34, 842 (1961).
3. J. A. Pople, W. G. Schneider, and H. J. Bernstein, "High Resolution Nuclear Magnetic Resonance," McGraw-Hill, New York, 1959, Ch. 9.
4. I. Solomon, Phys. Rev. 99, 559 (1955).
5. N. Bloembergen, J. Chem. Phys. 27, 595 (1957).
6. R. A. Bernheim, T. H. Brown, H. S. Gutowsky, and D. E. Woessner, J. Chem. Phys., 30, 950 (1959).

7. M. Tinkham, R. Weinstein, and A. F. Kip, Phys. Rev. 84, 848 (1951).
8. J. King and N. Davidson, J. Chem. Phys., 29, 787 (1958).
9. J. Eisinger, R. G. Shulman, and B. M. Szymanski, J. Chem. Phys., 36, 1721 (1962).
10. H. M. McConnell and S. B. Berger, J. Chem. Phys., 27, 230 (1957).
11. R. G. Pearson, J. Palmer, M. M. Anderson, and A. F. Allred, Z. Elektrochem. 64, 110 (1960).
12. J. P. Hunt and H. W. Dodgen, J. Chem. Phys., 35, 2261 (1961).
13. J. E. Wertz, J. Chem. Phys., 24, 484 (1956).

Supplimentary References on Relaxation Experiments

- J. A. Jackson, J. F. Lemons and H. Taube, J. Chem. Phys. 32, 553, (1960).
- P. F. Cox, and L. O. Morgan, J. Am. Chem. Soc., 81, 6409 (1959).
- L. O. Morgan, J. Murphy, and P. F. Cox, J. Am. Chem. Soc. 81, 5043 (1959).
- V. A. Shcherbakov, Zhur. Strukt. Khim. 2, 484-517 (1961).
- R. Kh. Timerov and K. A. Valiev, Zhur. Eksptl. e Theoretic. Fig. 41, 1566 (1961).
- W. Lipscomb and A. Kaczmarczyk, Proc. Natl. Acad. Sci. U. S. 47, 1796 (1961).
- R. E. Connick and E. D. Stover, J. Phys. Chem. 65, 2075 (1961).
- N. C. Li, R. L. Scruggs, and E. D. Becker, J. Am. Chem. Soc. 84, 4650 (1962).
- J. Eisinger, R. G. Shulman, and W. E. Blumberg, Nature 192, 963(1961).
- F. Klanberg, J. P. Hunt, and H. W. Dodgen, Inorg. Chem. 2, 139(1963).

II. Contact Shift

14. A. Foreman, J. Murrell, and L. Orgel, J. Chem. Phys., 31, 1129 (1959).
15. H. M. McConnell and C. H. Holm, J. Chem. Phys., 27, 314 (1957).
16. H. M. McConnell and C. H. Holm, J. Chem. Phys., 28, 749 (1958).
17. D. A. Levy and L. E. Orgel, Mol. Phys., 3, 583 (1960).
18. R. E. Benson, D. R. Eaton, A. D. Josey and W. D. Phillips, J. Am. Chem. Soc., 83, 3714 (1961).
19. W. D. Phillips and R. E. Benson, J. Chem. Phys., 33, 607 (1960).
20. D. R. Eaton, A. D. Josey, W. D. Phillips, and R. E. Benson, Mol. Phys., 5, 407 (1962).
21. D. R. Eaton, A. D. Josey, W. D. Phillips, and R. E. Benson, J. Chem. Phys., 37, 347 (1962).

22. D. R. Eaton, A. D. Josey, R. E. Benson, W. D. Phillips, and T. L. Cairns, J. Am. Chem. Soc., 4100 (1962).
23. D. R. Eaton, W. D. Phillips, and D. J. Caldwell, J. Am. Chem. Soc., 397 (1963).
24. E. Fermi, Z. Physik. 60, 320 (1930).
25. H. M. McConnell, J. Chem. Phys., 24, 764 (1956).
26. S. I. Weissman, J. Chem. Phys., 25, 890 (1956).
27. H. M. McConnell and D. B. Chestnut, J. Chem. Phys., 28, 107 (1958).
28. H. M. McConnell, Proc. Natl. Acad. Sci. U. S. 43, 721 (1957).
29. H. M. McConnell, J. Chem. Phys., 28, 1188 (1958).
30. H. M. McConnell, and D. B. Chestnut, J. Chem. Phys., 27, 984 (1957).
31. L. E. Orgel, J. Chem. Phys. 30, 1617 (1959).
32. H. M. McConnell, and H. H. Dearman, J. Chem. Phys. 28, 51(1958).
33. A. Mukherji and T. P. Das, Phys. Rev. 111, 1479 (1958).
34. A. J. Freeman and R. E. Watson, Phys. Rev. Letters 6, 343(1961).
35. R. E. Watson, and A. J. Freeman, Phys. Rev. Letters 6, 277(1961).
36. F. Keffer, T. Oguchi, W. O'Sullivan, and J. Yamashita, Phys. Rev. 115, 1553 (1959).
37. R. G. Shulman and K. Knox, Phys. Rev. Letters, 4, 603 (1960).
38. W. B. Lewis, J. A. Jackson, J. F. Lemons, and H. Taube, J. Chem. Phys. 36, 694 (1962).

AQUEOUS INFRARED SPECTROSCOPY AND
ITS APPLICATIONS TO INORGANIC CHEMISTRY

Ronald H. Carlson

April 23, 1963


INTRODUCTION

Water has generally been recognized as being a very poor infrared solvent. However, with the advent of the modern double beam spectrophotometer and with the development of suitable optical materials for use as infrared cell windows, aqueous infrared spectroscopy has rapidly become an important tool for the investigation of many inorganic and biological systems.

Coblentz (1) was the first to show that H₂O can be used as an infrared solvent. More recently, Gore and coworkers (2) have shown that H₂O, when used in conjunction with D₂O, can be extremely useful in this respect. The field of aqueous infrared spectroscopy was not extensively pursued until about 1953 when Blout and Lenormant (3) published some infrared spectra of biologically important materials in H₂O and D₂O. Since then, interest in this field has grown considerably and the number of applications is increasing.

THEORY AND EXPERIMENTAL

Water is a nonlinear triatomic molecule and as such should show $3N-6=3$ normal modes of vibration. The normal modes of vibration of H₂O, along with their symmetry classifications and absorption frequencies are (4):



The corresponding normal modes of vibration of D₂O and HDO vapor have absorbance frequencies of $\nu_1 = 2666 \text{ cm.}^{-1}$, $\nu_2 = 1178 \text{ cm.}^{-1}$, $\nu_3 = 2789 \text{ cm.}^{-1}$ and $\nu_1 = 2719 \text{ cm.}^{-1}$, $\nu_2 = 1402 \text{ cm.}^{-1}$, $\nu_3 =$ (overlapped by ν_3 of H₂O), respectively.

In liquid H₂O, the aforementioned absorption bands are broadened out considerably, with the result that there is an almost continuous absorption over the entire rock salt region. There are, however, two "windows" in this region where water is reasonably transparent; these are from 3.5μ to 5.8μ (2860 cm.^{-1} to 1730 cm.^{-1}) and from 6.5μ to 10.5μ (1540 cm.^{-1} to 955 cm.^{-1}). It is generally agreed that hydrogen bonding is the principal cause

for the intense infrared absorption of water, since perturbations by neighboring molecules are regarded as a prime factor in producing line broadening in liquids (5). The compound D₂O is quite a bit more transparent than H₂O and exhibits windows in the rock salt region from 2.5 μ to 3.6 μ (4000 cm.⁻¹ to 2780 cm.⁻¹), 4.8 μ to 7.8 μ (2080 cm.⁻¹ to 1280 cm.⁻¹), and 8.7 μ to 13.3 μ (1150 cm.⁻¹ to 750 cm.⁻¹). Thus it can be seen that H₂O, when used in conjunction with D₂O, is a fairly transparent infrared solvent and as such enables one to obtain a spectrum over the entire rock salt region.

One of major experimental difficulties encountered in the use of aqueous infrared spectroscopy is the need to use very short path lengths and, consequently, fairly concentrated solutions in order to obtain appreciable solute absorption. The longest path lengths that can conveniently be used in this regard are of the order of 0.075 mm (6). Path lengths of the order of 0.025 mm are usually employed in most infrared cells for optimum results. This thickness of water, small though it may seem, transmits only a small percentage of the incident radiation at 1100 cm.⁻¹. Therefore, fairly large slit widths are required in order to insure that a sufficient amount of energy reaches the detector. Operating at large slit widths leads to a loss in resolution of the spectrum. However, infrared bands in aqueous solution are usually so broad that a high degree of resolution is not necessary. The limit imposed by aqueous solutions upon the cell path length requires that a material have a solubility of at least 1% in H₂O in order to produce a spectrum. The infrared spectrum of more difficultly soluble materials can nonetheless be obtained through use of an ordinate scale expansion attachment available with most modern spectrophotometers. This attachment allows the spectroscopist to expand an infrared band up to ten times its normal height, and thus record the spectra of difficultly soluble materials the spectra of which under normal conditions would be extremely weak and distorted.

Another major difficulty encountered in aqueous infrared spectroscopy lies in the selection of suitable materials for cell windows which are both resistant to H₂O and transparent to infrared radiation. Table I lists some optical materials which can be used as cell windows (7,8).

<u>Material</u>	<u>μ</u>	<u>Cm.⁻¹</u>
fused quartz	4	2500
sapphire	6	1670
LiF	7	1430
CaF ₂	10	1000
BaF ₂	13	770
As ₂ S ₃	12.5	800
KRS-5 (TlBr, I)	35	285
Kodak I.R. Trans.	14	715
Kodak 80-20 (As modified Se glass)	25	400
AgCl	28	355

One of the more useful window materials for the construction of fixed thickness cells is BaF_2 since it is transparent over practically the entire rock salt region and is quite resistant to water. Plates of BaF_2 are, however, readily attacked by strongly acidic solutions and are very fragile. Cells constructed from this material must, therefore, be used with extreme care. Fixed-thickness AgCl cells find considerable use in investigations of highly acidic solutions where use of BaF_2 cells is not practical. Kodak "I. R. Trans" plates are extremely valuable in obtaining qualitative identification spectra where the need for a fixed thickness cell does not arise. Usually only one drop of solution is required in using the "I. R. Trans" plates.

Adequate compensation for residual solvent absorption in any one of the window regions of H_2O or D_2O can be obtained by placing a second absorption cell containing pure solvent into the reference beam of the spectrophotometer. The sample and reference cells should be of the same path length in order to insure maximum compensation. Another method of compensating for solvent absorption is to insert a transmittance screen or optical attenuator into the reference beam (9).

APPLICATIONS

(1) Qualitative identification of molecules in solution


One of the widest applications of aqueous infrared spectroscopy to date has been in the identification and examination of many biologically important systems. Aqueous media provide a natural physiological environment for such systems, and hence a study of their infrared spectra in this natural environment can be most rewarding. A good review of some biological systems which have been investigated by aqueous infrared techniques is offered by Parker (10).

The infrared spectra of several inorganic acids have been recorded both in H_2O and in D_2O , and the existence of discrete H_3O^+ ions in the liquid state has been confirmed (11). Three absorption bands were observed in the spectrum of each acid corresponding to the six normal modes of vibration of H_3O^+ . The similarity between these spectra and that of isoelectronic NH_3 was taken as sufficient proof that a true chemical bond between the proton and one of the adjacent water molecules exists long enough for the resulting species to undergo normal vibrations. Evidence regarding the degree of ionization of these acids was obtained through spectroscopic observation of the relative concentration of H_3O^+ existing in solution.

(2) Molecular structure as a function of pH

Aqueous amino acids exhibit a molecular structure which depends upon the pH of the solution in which the amino acid is dissolved. In highly acidic media, amino acids exist in a

cationic form, whereas in neutral media they exist as dipolar ions or "zwitterions", and in basic media they exist in an anionic form. The ionization scheme for an amino acid can be represented by the following, using glycine as an example:



Infrared spectroscopy is remarkably well adapted for following the attendant structural changes in aqueous amino acids as a function of pH (2,12). The unionized carboxyl group of the cation gives a characteristic absorption in D₂O at approximately 1740 cm.⁻¹, whereas the ionized carboxylate group of the zwitterion gives a characteristic absorption in D₂O at approximately 1610 cm.⁻¹, and the ionized carboxylate group of the anion gives a characteristic absorption at approximately 1580 to 1590 cm.⁻¹ in D₂O. Since each of the three structural forms of an amino acid has a characteristic carboxyl(ate) frequency associated with it, it is very easy to ascertain which species is predominant in solution at a given pH by noting the position and relative intensity of the respective carboxyl(ate) bands.

Martell and coworkers (13, 14) used infrared spectroscopy in an investigation of the molecular structure of nitrogen containing polycarboxylic acid ligands as a function of pH. Among the ligands which were studied were iminodiacetic acid (IMDA), nitrilotriacetic acid (NTA), and ethylenediaminetetraacetic acid (EDTA). The criterion used for structural identification as a function of pH was the same as that used in the case of simple amino acids, namely the position and relative intensity of the asymmetric carboxyl(ate) stretching band. In these amino-acid type systems, three carboxyl(ate) asymmetric stretching bands can be distinguished: type (A), un-ionized carboxyl band; type (B), γ -ammonium carboxylate (R₂NHCH₂COO⁻) band; and type (C), γ -amino carboxylate (R₂NCH₂COO⁻) band. The type A bands absorb at 1700~1730 cm.⁻¹ and have an extinction coefficient roughly one-third that of the type B and C bands, which absorb at 1620~1630 cm.⁻¹ and 1575~1585 cm.⁻¹, respectively. The results obtained by Martell and coworkers are summarized in Table II. Chapman and coworkers (15) and Sawyer and Tackett (16) have investigated the aqueous infrared spectra of IMDA, NTA, and EDTA. Their results are also included in Table II.

Table II

Ligand	pH	Observed bands (cm. ⁻¹)			Predominant Species	Ref.
		COOH(A)	COO ⁻ (B)	COO ⁻ (C)		
Iminodiacetic Acid	0.4	1733(s)	-	-	$ \begin{array}{c} \text{H} \quad + \quad \text{CH}_2\text{COOH} \\ \quad \quad \quad \diagdown \quad / \\ \quad \quad \quad \text{N} \\ \quad \quad \quad / \quad \diagdown \\ \text{H} \quad \quad \quad \text{CH}_2\text{COOH} \end{array} $	13
"	2.2	1721(m)	1619(s)	-	$ \begin{array}{c} \text{H} \quad + \quad \text{CH}_2\text{COO}^- \\ \quad \quad \quad \diagdown \quad / \\ \quad \quad \quad \text{N} \\ \quad \quad \quad / \quad \diagdown \\ \text{H} \quad \quad \quad \text{CH}_2\text{COOH} \end{array} $	13
"	4.1	-	1619(vs)	-	$ \begin{array}{c} \text{H} \quad + \quad \text{CH}_2\text{COO}^- \\ \quad \quad \quad \diagdown \quad / \\ \quad \quad \quad \text{N} \\ \quad \quad \quad / \quad \diagdown \\ \text{H} \quad \quad \quad \text{CH}_2\text{COO}^- \end{array} $	13

Table II (Continued)


Ligand	pH	Observed bands (cm. ⁻¹)			Predominant Species	Ref.
		COOH(A)	COO ⁻ (B)	COO ⁻ (C)		
Iminodiacetic Acid	10.9	-	-	1575(vs)	$\text{HN} \begin{cases} \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \end{cases}$	13
Nitrilotriacetic Acid	<<1.5	1732	-	-	$\text{HN} \begin{cases} + \text{CH}_2\text{COOH} \\ \text{CH}_2\text{COOH} \\ \text{CH}_2\text{COOH} \end{cases}$	15
"	<1.5	1730	1635	-	$\text{HN} \begin{cases} + \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COOH} \\ \text{CH}_2\text{COOH} \end{cases}$	15
"	2.3	1720(vw)	1623(m)	-	$\text{HN} \begin{cases} + \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COOH} \end{cases}$	14
"	5.9	-	1625(vs)	-	$\text{HN} \begin{cases} + \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \end{cases}$	14
"	10.2	-	-	1584(vs)	$\text{N} \begin{cases} \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \\ \text{CH}_2\text{COO}^- \end{cases}$	14
Ethylenediamine-tetraacetic Acid	<2	~1730	~1635	-	$\begin{matrix} \text{H} & & \text{H} \\ & & \\ \text{HOOCCCH}_2 & \text{N}(\text{CH}_2)_2 & \text{N} \\ & & \\ \text{H} & & \text{H} \\ \text{CH}_2\text{COOH} & & \text{CH}_2\text{COOH} \end{matrix}$	15
"	2.7	1700(vw)	1620(s)	-	$\begin{matrix} \text{H} & & \text{H} \\ & & \\ \text{HOOCCCH}_2 & \text{N}(\text{CH}_2)_2 & \text{N} \\ & & \\ \text{H} & & \text{H} \\ \text{CH}_2\text{COOH} & & \text{CH}_2\text{COOH} \end{matrix}$	16
"	4.0-5.3	-	1625(vs)	-	$\begin{matrix} \text{H} & & \text{H} \\ & & \\ \text{HOOCCCH}_2 & \text{N}(\text{CH}_2)_2 & \text{N} \\ & & \\ \text{H} & & \text{H} \\ \text{CH}_2\text{COO}^- & & \text{CH}_2\text{COO}^- \end{matrix}$	14, 16
"	7.8	-	1624(m)	1585(m)	$\begin{matrix} \text{H} & & \text{H} \\ & & \\ \text{HOOCCCH}_2 & \text{N}(\text{CH}_2)_2 & \text{N} \\ & & \\ \text{H} & & \text{H} \\ \text{CH}_2\text{COO}^- & & \text{CH}_2\text{COO}^- \end{matrix}$	14, 16
"	12.2	-	-	1585(vs)	$\begin{matrix} \text{H} & & \text{H} \\ & & \\ \text{HOOCCCH}_2 & \text{N}(\text{CH}_2)_2 & \text{N} \\ & & \\ \text{H} & & \text{H} \\ \text{CH}_2\text{COO}^- & & \text{CH}_2\text{COO}^- \end{matrix}$	14

The structural data reported in Table II agree extremely well with results predicted from ionization schemes based upon pK_a data and the known relative acidities of the COOH and R₃NH groups.

Values of pK_a for EDTA were calculated from aqueous infrared data by Martell and coworkers (14). From a plot of the peak intensities of the 1623 cm.^{-1} or 1585 cm.^{-1} carboxylate band against the pH of the solution, the authors were able to obtain approximate pK_a values by taking the pH value at the midpoint of the extreme intensities for each dissociation step. Values of $pK_3=6.5$ and $pK_4=9.3$ were obtained from the plot presented in Figure I.

Figure I

Absorbancy of carboxylate bands of EDTA as a function of pH. A, 1623 cm.^{-1} band; B, 1585 cm.^{-1} band.



The intersection of the solid vertical line with curves A and B represents an extreme intensity for each of the two dissociation steps under investigation.

Either of the two curves in Figure I could have been used for the estimation of the pK_a values. The values of pK_3 and pK_4 obtained by this method are in fairly good agreement with those obtained from potentiometric studies ($pK_3=6.1$, $pK_4=10.1$).

Aqueous infrared spectroscopy adds another dimension to the study of neutralization reactions of complex molecules. Not only does this technique provide valuable information on solution equilibria (pK_a values), but it also gives information on the structure of the reacting species as well.

(3) Stability and chemical bonding of transition metal ion complexes

The aqueous infrared spectra of several transition-metal ion-amino acid chelates have been obtained by Martell and coworkers (17). From a qualitative study of the variation in the asymmetric and symmetric carboxylate stretching frequencies as a function of metal ion coordination, the authors were able to obtain information regarding the nature of the metal carboxylate link, as well as a relative order of metal-oxygen bond strengths. For a given amino acid in D_2O solution, it was found that the asymmetric carboxylate stretching frequency increased and that the symmetric carboxylate frequency decreased in the following order of metal ions: $Ni^{II}-Zn^{II}-Cu^{II}-Co^{III}$. This result can best be explained by assuming that the covalent character of the

metal-oxygen bond increases in the order of metal ions given above. Furthermore, since the asymmetric and symmetric carboxylate stretching frequencies shift in opposite directions with a change in metal ion, a chelate structure involving symmetrical coordination of the carboxylate group can be ruled out.

Jones and Penneman (18,19,20) have used infrared spectroscopy in a determination of the successive formation constants of aqueous cyanide complexes of Ag^{I} , Au^{I} , Cu^{I} , Hg^{II} , Cd^{II} , and Zn^{II} . A good discussion of this work was presented in an inorganic seminar given recently by Mr. Khalil (21).

Fronaeus and Larsson (22,23) have investigated a series of metal ion-thiocyanate complexes in aqueous solution by means of infrared spectroscopy. A metal thiocyanate complex in aqueous solution absorbs infrared radiation in the region 2000-2200 cm^{-1} , due to a vibration involving the $\text{C}\equiv\text{N}$ stretching mode. For most thiocyanate complexes, this absorption band has a frequency quite distinct from that of the free thiocyanate ion and hence gives unequivocal evidence for the formation of a new molecular species in solution. For some of the complexes studied, it was possible to calculate quantitatively the extinction coefficient and formation constant of the 1:1 species. In addition, a bonding scheme, based upon the variation in frequency and integrated intensity of the complex peak as a function of metal ion coordination, was proposed for all of the complexes studied. The work of Fronaeus and Larsson is summarized in Table III.

Table III

Bonding Class	System	ν (cm^{-1})	ϵ (l/mole cm^{-1})	$\Delta\nu_{1/2}$	$\epsilon \cdot \Delta\nu_{1/2} \times 10^{-4}$	B_1
	SCN^-	2066	537	37	1.99	
I	$\text{Mn}^{\text{II}} + \text{SCN}^-$	2093	655	31	2.03	4.4
I	$\text{Fe}^{\text{II}} + \text{SCN}^-$	2099	-	-	-	-
I	$\text{Co}^{\text{II}} + \text{SCN}^-$	2112	696	29	2.02	8.9
I	$\text{Ni}^{\text{II}} + \text{SCN}^-$	2119	640	32	2.05	15.0
I	$\text{Zn}^{\text{II}} + \text{SCN}^-$	2109	685	34	2.32	8.2
II	$[\text{Co}(\text{NCS})_4]^{2-}$	2075	1900	29	4.9	-
II	$\text{Fe}^{\text{III}} + \text{SCN}^-$	2045	1310	49	6.4	-
II	$[\text{Cr}(\text{NCS})_6]^{3-}$	2088	937	39	3.7	-
II	$\text{UO}_2^{\text{II}} + \text{SCN}^-$	2066	1100	40	4.4	-
III	$\text{Cd}^{\text{II}} + \text{SCN}^-$	2132	635	24	1.5	-
III	$[\text{Pt}(\text{SCN})_6]^{2-}$	2126	246	15	0.4	-
III	$[\text{Hg}(\text{SCN})_4]^{2-}$	2112	420	27	1.1	-

ν , frequency of absorption
 ϵ , extinction coefficient
 $\Delta\nu_{1/2}$, band half width
 $\epsilon \cdot \Delta\nu_{1/2}$, approximation to the integrated intensity
 B_1 , formation constant of 1:1 complex

The class I complexes are considered to be essentially ionic in character with only a very small degree of electron exchange between metal and ligand; they are characterized by an absorption frequency somewhat greater than that of the free thiocyanate ion and an integrated band intensity approximately equal to that of the free thiocyanate ion. The class II complexes are considered to be more covalent in character, with lone-pair donation occurring through the nitrogen end of the ligand; they are characterized by an absorption frequency approximately equal to or less than that of the free thiocyanate ion and an integrated band intensity greater than that of the free thiocyanate ion. The class III complexes are covalent in character, with lone-pair donation occurring through the sulfur end of the ligand; they are characterized by an absorption frequency somewhat greater than that of the free thiocyanate ion and an integrated band intensity somewhat less than that of the free thiocyanate ion.

CONCLUSION

Aqueous infrared spectroscopy is a comparatively new experimental technique. It is a technique which not only provides valuable information regarding the nature of metal-ligand bonding in solution but also is without equal in characterizing the structures of reacting species in aqueous neutralization reactions involving amino acid and amino-polycarboxylic acid molecules. There is no doubt that, in the future, infrared spectroscopy will play an important role in determining the structures of the metal chelates of these molecules in aqueous solution.

BIBLIOGRAPHY

1. W. W. Coblentz, "Investigations of Infrared Spectra," p. 56, Carnegie Institution of Washington, Washington, D.C., (1905).
2. R. C. Gore, R. B. Barnes, and E. Petersen, *Anal. Chem.*, 21, 382 (1949).
3. E. R. Blout and H. Lenormant, *J. Opt. Soc. Am.*, 43, 1093(1953).
4. G. Herzberg, "Infrared and Raman Spectra of Polyatomic Molecules," p. 171, D. Van Nostrand Co., Inc., Princeton, New Jersey (1945).
5. D. A. Ramsay, *J. Am. Chem. Soc.*, 74, 72 (1952).
6. J. D. S. Goulden, *Spectrochim. Acta*, 1959, 657.
7. H. Sternglanz, *Appl. Spectroscopy*, 10, 77 (1956).
8. F. C. Nachod and C. M. Martini, *Appl. Spectroscopy*, 13, 45 (1959).
9. W. J. Potts, Jr. and N. Wright, *Anal. Chem.*, 28, 1255 (1956).
10. F. S. Parker, "The Perkin-Elmer Instrument News for Science and Industry," 13, 1 (1962).
11. M. Falk and P. A. Giguere, *Can. J. Chem.*, 35, 1195 (1957).
12. H. Lenormant, *Compt. rend.*, 234, 1959 (1952).
13. K. Nakamoto, Y. Morimoto, and A. E. Martell, *J. Am. Chem. Soc.*, 84, 2081 (1962).
14. K. Nakamoto, Y. Morimoto, and A. E. Martell, *J. Am. Chem. Soc.*, 85, 309 (1963).
15. D. Chapman, D. R. Lloyd, and R. H. Prince, *Proc. Chem. Soc.*, 1962, 336.

16. D. T. Sawyer and J. E. Tackett, *J. Am. Chem. Soc.*, 85, 314 (1963).
17. K. Nakamoto, Y. Morimoto, and A. E. Martell, *J. Am. Chem. Soc.*, 83, 4528 (1961).
18. L. H. Jones and R. A. Penneman, *J. Chem. Phys.*, 22, 965 (1954).
19. R. A. Penneman and L. H. Jones, *ibid.*, 24, 293 (1956).
20. R. A. Penneman and L. H. Jones, *J. Inorg. Nucl. Chem.*, 20, 19 (1961).
21. F. L. Khalil, *Inorganic Seminars*, University of Illinois, 1961, 183.
22. S. Fronaeus and R. Larsson, *Acta Chem. Scand.*, 16, 1433(1962).
23. S. Fronaeus and R. Larsson, *ibid.*, 16, 1447 (1962).

FLASH PHOTOLYSIS AND KINETIC SPECTROSCOPY

Michael R. Rosenthal

May 7, 1963

INTRODUCTION AND EXPERIMENTAL (41)

Flash photolysis and kinetic spectroscopy is a powerful method for studying free radicals and transient intermediates in photochemical and thermal reactions. These intermediates, the lifetimes of which are only microseconds or milliseconds, may be photographed in absorption and their rates of growth and decay determined. The fact that they are often produced in vibrationally excited states has led to studies of their relaxation.

A bank of condensers is discharged through a gas, usually xenon, producing a high-intensity flash. Lying parallel and close to the flash tube is a reaction tube. By means of electronic delay circuits, a second flash automatically throws a beam of light through the length of the reaction vessel. The contents are then photographed in absorption by a Littrow spectrograph. A series of such photographs is taken at varying time intervals in order to follow the growth and decay of the species present with time. Plate photometry is used to make these observations quantitative. Representative data for an operation of this type are as follows:

	<u>Photolysis Flash</u>	<u>Spectroscopic Flash</u>
Gas Pressure	10 cm.	10 cm.
Tube Length	50 cm.	10 cm.
Capacity	40 μ F.	2 μ F.
Voltage	10 Kv.	10 Kv.
Duration	50 μ sec.	10 μ sec.
Energy	2000 J.	100 J.

The most common studies are kinetic and mechanism studies of explosions (adiabatic) in the gas phase, isothermal gas phase reactions (large excess inert gas), and isothermal flash photolysis of solutions. The spectra obtained are limited to 1900 Å by the transparency limit of quartz, unless special techniques are employed. The outstanding advantage of this method is that unlike conventional photolysis techniques, flash photolysis leads to nearly complete reaction and consequently, high radical or intermediate concentrations.


APPLICATIONS

A great deal of work has been done on iodine atom recombination. It was discovered early that the rate of recombination is second order with respect to iodine atoms and first order with respect to a third body (49, 28, 16).

The fact that the rate constant varied for different third bodies was attributed to varying Van der Waals forces in the third body. A reinvestigation, however, indicated that the simple third-order rate law was not obeyed and that the rate constant increased as the ratio of iodine to third body increased (17). This had been previously undetected as a consequence of an opposing thermal effect. New values of the rate constants for several third bodies were obtained by extrapolation. The studies were extended to bromine with argon and oxygen as third bodies (51). The detection of a transient spectrum in the flash photolysis of iodine in benzene (48) led to the proposal that a charge-transfer intermediate was involved. Similar results in other solvents strengthened this proposal (22). A search was undertaken to find such an intermediate in the gas phase (45). A transient spectrum was found when iodine was flash photolyzed in nitric oxide. This was attributed to [NOI] (46). Further work is in progress.

Because of space limitations, the presentation of the other work in this field shall be brief. For more complete information, the reader is referred to the references.

The flash photolysis of quinones has shown the existence of equilibria of the following type: (9, 10).



Rate constants and equilibrium constants have been determined.

The decomposition of carbon dioxide during flash photolysis has been studied and the effect of varying experimental conditions determined (8).

The flash photolyses of nitrosyl chloride, nitrosyl bromide (1,3), and nitric oxide (2,4) have led to the detection of vibrationally excited electronic ground states of nitric oxide.

The adiabatic flash photolysis of sulfur dioxide (37) produced a species identified as a high temperature isomer. Sulfur trioxide was also studied isothermally and mechanisms proposed for its decomposition.

The adiabatic flash photolysis of hydrogen sulfide (36) and the isothermal flash photolysis of carbon disulfide (11) were carried out and mechanisms for their decompositions were postulated.

Several combustion studies of the antiknock properties of tetraethyl lead in hydrocarbons have been made (20,21). The antiknock properties were attributed to the ability of the lead compound to remove oxygenated intermediate species through the formation of lead(II) oxide, which eventually was reduced to elemental lead. Several other hydrocarbon combustion studies have been carried out (34,35) as well as a study of the effectiveness of other additives as antiknock agents (12).

The production of ClO radicals was noted in the flash photolysis of both chlorine-oxygen mixtures (42) and of chlorine monoxide (18). Rate constants were calculated and mechanisms of dissociation and recombination were proposed.

The flash photolyses of aldehydes have been studied under both slow and explosive conditions and mechanisms proposed for their decomposition (24, 32).

The flash photolysis of anthracene-styrene mixtures was found to lead to polymerization of the styrene through the formation of the triplet state of anthracene (38). The triplet state of anthracene in hexane solution has been studied in great detail (43). The flash photolysis of mercury vapor has led to formation of the triplet state (15). The subject of radiationless transitions from triplet to singlet states has been discussed in detail (44).

Nickel tetracarbonyl was flash photolyzed under isothermal conditions in several gases and a mechanism for its decomposition was proposed (13,14).

The flash photolyses of ammonia (23), deuterated ammonia (23), phosphine (27,28), and deuterated phosphine (27,28) led to detection of NH_2 , ND_2 , PH_2 , and PD_2 radicals, respectively, and to postulation of mechanisms of formation and decay. The flash photolysis of phosphine in oxygen was also studied (40).

The color change of the substance



in water was studied by means of flash photolysis (53).

The isothermal flash photolysis of azomethane (50) and hydrogen azide (52) have been carried out. Intermediates and decomposition mechanisms have been postulated.

Flash photolysis has been used to study and characterize methyl and methylene radicals (25) and to determine acidity constants for several organic molecules in their lowest triplet states (26).

Flash photolysis has been used to study ozone decomposition (29,31) and the formation of vibrationally excited oxygen from the isothermal decomposition of chlorine dioxide and nitrogen dioxide(27).

Vibrationally excited CN radicals have been produced in the flash photolyses of cyanogen, cyanogen bromide, and cyanogen iodide (5).

The first observation of the spectrum of benzyne has been reported as a transient spectrum in the flash photolysis of -COO(6).

The spectral range has been extended to 1500 Å in a flash photolysis of water vapor (7).

A flash photolysis study of inorganic halides in solution has produced transient spectra associated with dihalide anions(19).

Flash photolysis has been used to study the combustion of methyl iodide in great detail (33).

The mechanism of relaxation of vibrationally excited states produced in flash photolyses and the energy distribution among them have been discussed (30).

CONCLUSION

The formation of short-lived intermediates and free radicals in concentrations higher than those produced by more conventional methods has prompted studies elucidating fast-reaction kinetics and mechanism. The growing interest in this area promises much more useful data and an interesting future for the study of "instantaneous" reactions.

REFERENCES

1. Basco, N., and Norrish, R. G. W., Nature, 189, 455 (1961).
2. Basco, N., Callear, A. B., and Norrish, R. G. W., Proc. Roy. Soc. (London), A260, 459 (1961).
3. Basco, N., and Norrish, R. G. W., Proc. Roy. Soc. (London), A268, 291 (1961).
4. Basco, N., Callear, A. B., and Norrish, R. G. W., Proc. Roy. Soc. (London), A269, 180 (1962).
5. Basco, N., Nicholas, E., Norrish, R. G. W., and Vickers, W.H.J., Proc. Roy. Soc. (London), A272, 147 (1963).
6. Berry, R. S., Spokes, G. N., and Stiles, R. M., J. Am. Chem. Soc., 84, 3570 (1962).
7. Black, G., and Porter, G., Proc. Roy. Soc. (London), A266, 185 (1962).

8. Bortner, M. H., Kovard, V. D., and Myerson, A. L., J. Opt. Soc. Am., 50, 172 (1960).
9. Bridge, N. K., and Porter, G., Proc. Roy. Soc. (London), A244, 259 (1958).
10. Bridge, N. K., and Porter, G., Proc. Roy. Soc. (London), A244, 276 (1958).
11. Callear, A. B., and Norrish, R. G. W., Nature, 188, 53 (1960).
12. Callear, A. B., and Norrish, R. G. W., Proc. Roy. Soc. (London), A259, 304 (1960).
13. Callear, A. B., Proc. Roy. Soc. (London), A265, 71 (1961).
14. Callear, A. B., Proc. Roy. Soc. (London), A265, 88 (1961).
15. Callear, A. B., and Norrish, R. G. W., Proc. Roy. Soc. (London), A266, 299 (1962).
16. Christie, M. I., Norrish, R. G. W., and Porter, G., Proc. Roy. Soc. (London), A216, 152 (1953).
17. Christie, M. I., Harrison, A. J., Norrish, R. G. W., and Porter, G., Proc. Roy. Soc. (London), A231, 446 (1955).
18. Edgecombe, F. H. C., Norrish, R. G. W., and Thrush, B. A., Proc. Roy. Soc. (London), A243, 24 (1957).
19. Edgecombe, F. H. C., and Norrish, R. G. W., Proc. Roy. Soc. (London), A253, 154 (1959).
20. Erhard, K. H. L., and Norrish, R. G. W., Proc. Roy. Soc. (London), A234, 178 (1956).
21. Erhard, K. H. L., and Norrish, R. G. W., Proc. Roy. Soc. (London), A259, 297 (1960).
22. Gover, T. A. and Porter, G., Proc. Roy. Soc. (London), A262, 476 (1961).
23. Herzberg, G., and Ramsay, D. A., Discussions Royal Soc., 14, 11 (1953).
24. Herzberg, G., and Ramsay, D. A., Proc. Roy. Soc. (London), A233, 34 (1955).
25. Herzberg, G., Proc. Royal Soc. (London), A262, 291 (1961).
26. Jackson, G., and Porter, G., Proc. Roy. Soc. (London), A260, 13 (1960).
27. Lipscomb, F. J., Norrish, R. G. W., and Thrush, B. A., Proc. Roy. Soc. (London), A233, 455 (1956).
28. Marshall, R., and Davidson, N., J. Chem. Phys., 21, 659 (1953).
29. McGrath, W. D., and Norrish, R. G. W., Proc. Roy. Soc. (London), A242, 265 (1957).
30. McGrath, W. D., and Norrish, R. G. W., Z. physik. Chem. (Frankfurt), 15, 245 (1958).
31. McGrath, W. D., and Norrish, R. G. W., Proc. Roy. Soc. (London), A254, 317 (1960).
32. McKellar, J. F., and Norrish, R. G. W., Proc. Roy. Soc. (London), A254, 147 (1960).
33. McKellar, J. F., and Norrish, R. G. W., Proc. Roy. Soc. (London), A263, 51 (1961).
34. Norrish, R. G. W., Discussions Royal Soc., 14, 16 (1953).
35. Norrish, R. G. W., Porter, G., and Thrush, B. A., Proc. Roy. Soc. (London), A216, 165 (1953).
36. Norrish, R. G. W., and Zeelenberg, A. P., Proc. Roy. Soc. (London), A240, 293 (1957).
37. Norrish, R. G. W., and Oldershaw, G. A., Proc. Roy. Soc. (London), A249, 498 (1959).

38. Norrish, R. G. W., and Simons, J. P., Proc. Roy. Soc. (London), A251, 4 (1959).
39. Norrish, R. G. W., and Oldershaw, G. A., Proc. Roy. Soc. (London), A262, 1 (1961).
40. Norrish, R. G. W., and Oldershaw, G. A., Proc. Roy. Soc. (London), A262, 10 (1961).
41. Norrish, R. G. W., Am. Sci., 50, 131 (1962).
42. Porter, G., and Wright, F. J., Discussions Royal Soc., 14, 23 (1953).
43. Porter, G., and Windsor, M. W., Discussions Faraday Soc., 17, 178 (1954).
44. Porter, G. and Wright, M. R., Discussions Faraday Soc., 27, 18 (1959).
45. Porter, G., and Smith, J. A., Proc. Roy. Soc. (London), A261, 28 (1961).
46. Porter, G., Szabo, Z. G., and Townsend, M. G., Proc. Roy. Soc. (London), A270, 493 (1962).
47. Ramsay, D. A., Nature, 178, 374 (1956).
48. Rand, S. J., and Strong, R. L., J. Am. Chem. Soc., 82, 5(1960).
49. Russell, K. E., and Simons, J., Proc. Roy. Soc. (London), A217, 271 (1953).
50. Sleppy, W. C., and Calvert, J. G., J. Am. Chem. Soc., 81, 769 (1959).
51. Strong, R. L., Chien, J. C. W., Graf, P. E., and Willard, J.E., J. Chem. Phys., 26, 1287 (1957).
52. Thrush, B. A., Proc. Roy. Soc. (London), A235, 143 (1956).
53. Wettermark, G., J. Am. Chem. Soc., 84, 3658 (1962).

THE MUSSBAUER EFFECT AND ITS
APPLICATIONS TO INORGANIC CHEMISTRY

K. F. Purcell

May 14, 1963

INTRODUCTION

Nuclear resonance spectroscopy is a spectroscopic method that possesses several unique features. As the name implies, nuclear resonance absorption is the resonant excitation of nuclei. Very briefly, the experimental procedure requires a source of the resonant γ -energy mounted on a variable velocity mechanism, an absorber (sample), and a detector. The intensity of γ -rays passing through the absorber is plotted as a function of the source velocity. The resulting spectrum will yield information about the electron density at the nucleus and molecular symmetry.

THEORY

Consideration of a particle which emits a photon shows that there is a recoil momentum associated with the emission(1). For low energy photons, the recoil is negligible, but in the present situation where $E_\gamma \approx 10$ -100 k ev., the recoil energy loss can be appreciable ($\approx 10^{-2}$ ev.). It is given by $R = \frac{E_\gamma^2}{2 mc^2}$. For emitting (absorbing) nuclei tightly bound in a crystal of "finite" size, the crystal absorbs the recoil and because of the large mass of the crystal the recoil velocity and energy transferred to the lattice are negligible. There is another mechanism for recoil energy loss to the lattice, however. Emission of a γ -ray may cause excitation of lattice vibrational modes and lead to a loss of energy (1,2,3,4). Any emission event which occurs leaving the lattice in its initial vibrational state is termed "recoilless". Recoilless resonance is also called Müssbauer resonance, after the man who discovered it (5).

We shall now consider the crystal properties and experimental conditions which lead to Müssbauer resonance (1,4). A real crystal is assumed to possess the properties of a Debye solid. For such a substance, Debye showed that the distribution of vibrational frequencies is proportional to frequency squared, as indicated in Figure 1.




Figure 1

There are no vibrations with $\omega > \omega_D$, the "cut-off" frequency. As far as we are concerned, this relationship means that there are more high-energy modes than low. Quantum theory tells us that the average energy transferred to the lattice for an emission event is

$$\bar{R} = \sum_f P(f \leftarrow i) (\underline{E}_f - \underline{E}_i) = \underline{E}_i^2 / 2\underline{MC}^2$$

where \underline{E}_i and \underline{E}_f are the initial and final lattice energies and $P(f \leftarrow i)$ is the probability for the lattice going from state i to state f during emission. If we assume that the Debye model applies, the Mössbauer effect will be greatest when the crystal possesses a high Debye temperature and when the temperature of the crystal is low.

The fact that $\bar{R} = \underline{E}_i^2 / 2\underline{MC}^2$ restricts the occurrence of the Mössbauer effect to nuclear transitions for which $\underline{E} < 150$ kev. (assuming a maximum realistic θ_D). The dependence of the Mössbauer effect on \underline{E} , θ_D , and \underline{T} is shown in Figure 2 (2).




Figure 2

Two aspects of chemical structure - electron distribution and molecular symmetry - play a significant role in determining the nature of the Mössbauer spectrum of a molecule. Molecular symmetry is reflected in the spectrum of a nucleus which possesses a quadrupole moment, through the presence (or lack) of a quadrupole split resonance line (6). The effect of a difference in electron densities at source and absorber nuclei is termed "isomer shift", and such a difference determines the position of the resonance line in the spectrum(1). This is because the nuclear energy levels are shifted slightly by a change in electrostatic interaction with the extranuclear electrons. Since "s" electrons have the greatest density at the nucleus, changes in "s" wave functions at the nucleus are of most importance in producing the isomer shift.

In the case of the iron compounds to be discussed later, increasing "s" density over a series of compounds produces a progressively smaller isomer shift with respect to a given

source (7,8). The nuclide Fe^{57} also possesses a quadrupole moment in the excited state but not in the ground state (6).

EXPERIMENTAL PROCEDURE

In order to "match" the energy of the emitted γ ray to that required for absorption, the frequency is shifted by the linear Doppler effect (1). The source (usually) is placed on a velocity drive and driven toward or away from the absorber with a known velocity. The intensity of γ rays passing through the absorber is measured by a NaI(Tl) scintillation counter and plotted as a function of the velocity. Mössbauer used a record turntable as a velocity mechanism (5). The apparatus most commonly used now is shown schematically in Figure 3 (1).




Figure 3

The source, mounted on the speaker, is driven with changing velocity as shown in the "velocity spectrum". The counting pulses from the detector are modulated with this velocity spectrum, and a multichannel analyzer sorts the pulses as a function of velocity.

Mössbauer resonance has been observed for 16 nuclei, all of interest to the inorganic chemist(1). These nuclei are Fe^{57} , Ni^{61} , Zn^{67} , Sn^{119} , Gd^{155} , Dy^{161} , Er^{166} , Tm^{169} , Yb^{170} , Hf^{177} , Ta^{181} , W^{182} , Re^{187} , Ir^{191} , Ir^{193} , and Au^{197} . To illustrate source and absorber preparation, the procedure for Fe^{57} experiments will be outlined. The nuclide Co^{57} undergoes electron-capture decay to excited state Fe^{57} , which falls to the ground state emitting the 14.4 kev. γ ray (1,3). The Co^{57} nuclide, then, is the source for Fe^{57} Mössbauer experiments. The standard "iron" source is prepared by depositing Co^{57} on a thin sheet of 310 stainless steel and heating to diffuse the cobalt into the steel (9). This source emits a monoenergetic γ ray. The sample, the spectrum of which is desired, may be a very thin crystal, a small amount of powder (10), or a thin layer of "frozen" liquid (11).

An important correction to the isomer shift must be made when one wishes to compare the results of experiments performed at different temperatures. The correction is for the "second-order Doppler effect" and arises as a result of the temperature dependence of the lattice particle kinetic energy. The "temperature shift" has been shown to be given by

$$\frac{\Delta \nu}{\nu} = -2.4 \times 10^{-15} \frac{v^2}{T} (\text{°K})^{-1}$$

where ν is source velocity at resonance (12,13).

CHEMICAL APPLICATIONS

One of the earliest papers in which an attempt was made to correlate the isomer shift (δ) and quadrupole splitting (ΔE) with molecular properties is that of Ingalls and coworkers(7). Their results (see Table) indicate that "ionic" Fe(II) has a $\delta \approx 1.0-1.5$ mm. sec.⁻¹ and "ionic" Fe(III) $0.40 < \delta < 1.0$ mm. sec.⁻¹. It is seen that ΔE for Fe(III) is zero except for Fe₂O₃, whereas ΔE for the Fe(II) salts ranges from 1.80 mm. sec.⁻¹ to 3.20 mm. sec.⁻¹. These authors explained the effect of the oxidation state of iron on the quadrupole splitting as due directly to asymmetry of charge distribution in the 3d shell.

Their argument breaks down, however, in the case of Fe Fe(CN)₆. In view of the data available it is more logical to consider ligand-field distortions as giving rise to the splittings. Such distortions may be transmitted to the nucleus through the 3p electrons and/or by asymmetric shielding of the "s" electrons by the "d" electrons (14). Although these "mechanisms" are useful in correlating the data there have been no calculations to support them. The large ΔE found for the Fe(II) salts is believed to be due to tetragonal and rhombic distortions of the octahedral field about the nucleus. X-ray investigations(16) have shown this to be the case for Fe(NH₄)₂(SO₄)₂·6H₂O.

Several workers (10,14) have reported Mössbauer resonance data for the simple and complex iron cyanides, and Epstein (10) has reported spectra for a number of "ionic" and "covalent" iron complexes. The isomer shifts for all of the iron cyanides indicate extensive Fe - CN π bonding and close approach of the ligands to the iron ion in the anions. The quadrupole splittings may be rationalized by considering the electronic configuration about the iron nuclei and the effects of crystal distortions. In the mixed complex cyanides, the spectra indicate that Fe(II) resides in the carbon hole(15). The ΔE of 0.4 mm. sec.⁻¹ for Fe(III) in KFe₂(CN)₆ is attributed to lattice water of crystallization. A similar proposal applies to Na₃Fe(CN)₆·H₂O. The spectrum of sodium nitroprusside, Na₂Fe(CN)₅NO·2H₂O, has been interpreted (10) as indicating iron in an oxidation state of +4.

Table I

<u>Compound</u>	ΔE (mm. sec. $^{-1}$)	δ (mm. sec. $^{-1}$)	<u>Reference</u>
Fe ₂ O ₃	0.24±0.03	0.47±0.03	7
Fe ₂ (SO ₄) ₃	0.0	0.55±0.05	7
FeCl ₃	0.0	0.45±0.05	7
FeCl ₂ ·4H ₂ O	3.00±0.05	1.35±0.05	7
FeSO ₄ ·7H ₂ O	3.20±0.05	1.40±0.05	7
Fe(NI ₄) ₂ (SO ₄) ₂ ·6H ₂ O	1.75±0.05	1.40±0.05	7
FeF ₂	2.68	1.40±0.05	7
K ₃ Fe(CN) ₆	0.26±0.05	-0.10±0.05	10, 14
K ₄ Fe(CN) ₆ ·3H ₂ O	0.0	0.0±0.05	14, 10
Na ₃ Fe(CN) ₆ ·H ₂ O	0.60	-0.05	10
Na ₂ Fe(CN) ₅ NO·2H ₂ O	1.76±0.01	-0.17±0.01	10
Fe ₄ [Fe(CN) ₆] ₃	-0.02 +0.55	0.0 (anion) 0.0 (cation)	10
Fe ₂ (CN) ₆	0	0.0±0.1 (Anion)	14
K ₂ FeFe(CN) ₆	0	0.53±0.1 (Cation)	14
	0	0.0±0.1 (Anion)	
	0.4±0.05	0.53±0.05 (Cation)	14
K ₂ FeFe(CN) ₆	0.6±0.2	0.3±0.2 (Anion)	14
	1.8±0.2	0.93±0.2 (Cation)	14

Epstein(10) and Wertheim and coworkers(11) have shown that Mössbauer resonance can be obtained from molecular crystals (e.g., Fe(II) Phthalocyanine, $\text{Fe}(\text{CO})_5$, $\text{Fe}_2(\text{CO})_9$, and $\text{Fe}_3(\text{CO})_{12}$). There has been a prodigious amount of infrared, x-ray, and kinetic data published on the structure of $\text{Fe}_3(\text{CO})_{12}$ (11). The Mössbauer spectrum of the compound shows the molecule to have the 3,3,3,3 structure in agreement with the kinetic studies and one interpretation of the infrared spectrum.

CONCLUSION

In view of the complex molecules to which Mössbauer resonance has been applied, it is not surprising that ambiguities exist in interpreting their spectra. It is necessary that a detailed systematic selection of compounds be made to examine the effects of σ and π bonding and ligand-field symmetry on the spectra.

REFERENCES

1. Frauenfelder, H., "The Mössbauer Effect," W. A. Benjamin Inc., New York, New York (1962).
2. Mössbauer, R. L., Science, 137, 731 (1962).
3. Mössbauer, R. L., Nuclear Science Abstracts, 12, 123 (1962).
4. Lipkin, H. J., Ann. Phys., 9, 332 (1960).
5. Mössbauer, R. L., Naturwissenschaften, 45, 538 (1958).
6. Kistner, O. C., and Sunyar, A. W., Phys. Rev. Letters, 4, 412 (1960).
7. Ingalls, R., Lang, G., and DeBenedetti, S., Phys. Rev. Letters, 6, 60 (1961).
8. Walker, L. R., Wertheim, G. K., and Jaccarino, V., Phys. Rev. Letters, 6, 98 (1961).
9. Wertheim, G. K., J. Appl. Phys., Suppl., 32, 1105 (1961).
10. Epstein, L. M., J. Chem. Phys., 36, 2731 (1962).
11. Herber, R. H., Kingston, W. R., and Wertheim, G. K., Inorg. Chem., 2, 153 (1963).
12. Pound, R. V. and Rebka, G. A., Phys. Rev. Letters, 4, 274 (1960).
13. Josephson, B. D., Phys. Rev. Letters, 4, 341 (1960).
14. Duncan, J. F. and Wigley, P. W. R., J. Chem. Soc., 1963, 1120.
15. Robin, M. B., Inorg. Chem., 1, 337 (1962).
16. Hückel, W., "Structural Chemistry of Inorganic Compounds," Vol. II, P. 920, Elsevier Publishing Company, New York (1951).


HYDROLYSIS OF METAL IONS

J. E. House, Jr.


May 23, 1963

INTRODUCTION


The aqueous solutions of most heavy metal ions are acidic. In the process of solution, some reaction takes place in which protons are liberated producing solutions which are acidic. Historically, the process has been written as



Since this requires the solid metal hydroxide to be present or a solution of the hydroxide, neither of which is observed in many systems, this is not a very accurate representation. Others are



and



Since many ions (e.g. Be^{2+} , Fe^{3+} , UO_2^{2+} , Bi^{3+} , etc.) produce acidic solutions, these reactions are fairly general. It is now known that fairly complicated hydrolysis products are produced in which two or perhaps even several metal ions form polynuclear species. Since the number of ions which hydrolyze to give acidic solutions is fairly large and several studies have been made on most of them, no attempt is made here to give a complete survey of the literature of hydrolytic processes. For a more complete survey of the literature, one should consult the Tables of Stability Constants (1). Neither are anionic hydrolysis reactions considered here.

THEORY AND TREATMENT OF DATA

To present an outline of the general methods used to describe hydrolytic equilibria, use will be made of the following:

List of Symbols

- A = the ligand species
- B = the central metal ion
- C_B = total concentration of species B
- C_A = total concentration of species A
- \bar{n} = the average number of ligands bound to B
- a = the concentration of free (unbound) A
- b = the concentration of free (unbound) B
- h = the concentration of H^+ set free by hydrolysis
- K_w = the ion product of water
- β_{qp} = the formation constant of complex $[B_qA_p]$, $[B_qA_p]/ab$
- χ_{qp} = the hydrolysis constant

INTRODUCTION

The purpose of this study is to investigate the effects of various factors on the rate of change of a certain variable. The results are presented in the following sections.

1. The first part of the study is devoted to a general discussion of the problem. It is shown that the rate of change of the variable is dependent on several factors, which are discussed in detail in the following sections.

2. In the second part, the effect of the first factor is investigated. It is shown that the rate of change increases as the value of this factor increases. This is illustrated by the following graph:

Figure 1 shows the relationship between the rate of change and the first factor. The curve shows that the rate of change increases rapidly at first and then levels off as the value of the first factor increases. This is due to the fact that the rate of change is proportional to the square of the first factor.

3. The third part of the study is devoted to an investigation of the effect of the second factor. It is shown that the rate of change is also dependent on this factor, and that the effect is similar to that of the first factor.

- References:
- 1. J. D. Smith, "The Rate of Change of a Variable," *Journal of Applied Mathematics*, vol. 10, pp. 1-10, 1950.
 - 2. A. B. Jones, "The Effect of Various Factors on the Rate of Change of a Variable," *Research Report*, 1950.
 - 3. C. E. Brown, "The Rate of Change of a Variable," *Journal of Applied Mathematics*, vol. 10, pp. 11-20, 1950.
 - 4. D. F. Green, "The Effect of Various Factors on the Rate of Change of a Variable," *Research Report*, 1950.
 - 5. E. G. White, "The Rate of Change of a Variable," *Journal of Applied Mathematics*, vol. 10, pp. 21-30, 1950.


For the hydrolysis of metal ions



where the ionic charges have been omitted. The hydrolysis constants χ_{qp} are given by

$$\chi_{qp} = \frac{[Bq(OH)_p]h^p}{b^q} = \beta_{qp}K_W^p \quad (5)$$

where the β_{qp} is the formation constant of the complex formed by the reaction



and is given by

$$\beta_{qp} = \frac{[Bq(OH)_p]}{[B]^q[OH]^p} \quad (7)$$

The problem of determining the extent of hydrolysis as occurring in equation (5) and the hydrolysis constants is intimately related to the stability of the hydroxo complexes formed. In order to discuss the extent of hydrolysis, the determination of stability constants must be considered. A brief survey of the methods used in the treatment of data will be given. The methods used for mononuclear complexes will be considered first.

MONONUCLEAR COMPLEXES

A. Bjerrum's Method (2).

The average number of ligands attached to the central metal ion B is given by

$$\bar{n} = \frac{C_A - a}{C_B} = \frac{\sum_{i=1}^p i[BA_i]}{b + \sum_{i=1}^p [BA_i]} \quad (8)$$

By making use of the definition of the formation constants

$$\bar{n} = \frac{\beta_1 a + 2\beta_2 a^2 + \dots + p\beta_p a^p}{1 + \beta_1 a + \dots + \beta_p a^p} \quad (9)$$

This can be written as

$$\bar{n} = (\bar{n} - 1)\beta_1 a + (\bar{n} - 2)\beta_2 a^2 + \dots + (\bar{n} - p)\beta_p a^p = 0 \quad (10)$$

The values of the β 's where p experiments have been performed giving p equations (8) are obtained by solution of the p equations using determinants.

The expression (6) cannot be used to determine \bar{n} when weak complexes are formed or where the central ion is present in very small concentrations because under these conditions $C_A \approx a$. In this case, the degree of formation of the complex, α_p , is used.

$$\alpha_p = \frac{[BA_p]}{C_B} = \frac{\beta_p a^p}{1 + \sum_{i=1}^p \beta_i a^i} \quad (11)$$

Using equation (9) and the definition of \bar{n} , it is possible to derive the expression for \bar{n} ,

$$\bar{n} = p - \frac{\partial \ln \alpha_p}{\partial \ln a} \quad (12)$$

If the concentration of the central ion or any one of the complexes is known as a function of the free ligand concentration, it is possible to obtain \bar{n} by graphical differentiation of the curve one gets from plotting $\log a$ against $\log \alpha_p$. The plot of \bar{n} against $\log a$ gives the formation curve of the system. Where the measured quantity is the free metal ion concentration, b ,

$$\bar{n} = \frac{\partial \log b}{\partial \log a} \quad (13)$$

Again \bar{n} can be determined for the system but here the approximation $C_A \approx a$ is not valid since the experimentally determined quantity is not "a" itself. The graphical method of successive approximation must be used to find \bar{n} and a . From equations (7) and (9), α_c and \bar{n} are functions of "a" only for mononuclear complexes.

This method applies to all systems whether the measured quantity be the concentration b , a , or one of the complexes BA_p . A plot of \bar{n} against a , the free metal ion concentration, immediately indicates the number of complexes which must be dealt with.

B. Leden's Method (3)

A function of the concentration of free A, $F(a)$, is defined by

$$F(a) = \frac{C_A - a}{ba} = \sum_{i=1}^p \beta_i a^{p-i} = \beta_1 + \beta_2 a + \dots + \beta_p a^{p-1} \quad (14)$$

From a plot of $F(a) = \frac{C_A - a}{ab}$ against a , the limit

$$\lim_{a \rightarrow 0} F(a) = \beta_1$$

is obtained graphically. The successive constants may be obtained

Let $f(x) = x^2 + 2x + 1$. Then $f(x) = (x+1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = -1$.

Let $f(x) = x^2 - 2x + 1$. Then $f(x) = (x-1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = 1$.

$$f(x) = x^2 + 2x + 1 = (x+1)^2$$

Let $f(x) = x^2 - 2x + 1$. Then $f(x) = (x-1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = 1$.

$$f(x) = x^2 - 2x + 1 = (x-1)^2$$

Let $f(x) = x^2 + 2x + 1$. Then $f(x) = (x+1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = -1$.

$$f(x) = x^2 + 2x + 1 = (x+1)^2$$

Let $f(x) = x^2 - 2x + 1$. Then $f(x) = (x-1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = 1$.

Let $f(x) = x^2 + 2x + 1$. Then $f(x) = (x+1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = -1$.

Let $f(x) = x^2 - 2x + 1$. Then $f(x) = (x-1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = 1$.

$$f(x) = x^2 + 2x + 1 = (x+1)^2$$

$$f(x) = x^2 - 2x + 1 = (x-1)^2$$

$$f(x) = x^2 + 2x + 1 = (x+1)^2$$

Let $f(x) = x^2 - 2x + 1$. Then $f(x) = (x-1)^2$. The function $f(x)$ is always non-negative, and it is zero only when $x = 1$.

from other functions $G(a)$ where

$$G(a) = \frac{F(a) - \beta_1}{a} = \beta_1 + \dots + \beta_p a^{p-2} \quad (15)$$

The graphical limit is determined again for $G(a)$ and yields β_2 since

$$\lim_{a \rightarrow 0} G(a) = \beta_2 \quad (16)$$

As an alternate to this method of extrapolation, the values of the β 's may be determined by solving a system of linear equations by means of determinants.

Fronaesus (4) has developed a similar method which uses the concept of \bar{n} along with a limiting process of evaluation of the formation constants. This method can be shown to be equivalent to the Leden method by a simple transformation of the $F(a)$ function of Leden to the function defined in Fronaesus' method. A complete review of these methods and typical calculations involving their use has been given by Sullivan and Hindman (5).

A large number of graphical and empirical methods have been developed which are less general in scope (6-14).

Methods involving the use of digital computers have also been devised for the solution of the linear equations for the values of the stability constants (15,16).

POLYNUCLEAR COMPLEXES

Most metal ions produce polynuclear complexes on hydrolysis. The theory of such processes has been worked out in great detail and the description of some of the more general methods specifically treating hydrolytic processes follows (17,19).

The total concentration of the central ion and ligand are given by

$$C_B = \sum_{q=1}^Q \sum_{p=0}^P q [B_q A_p] = \sum_{q=1}^Q \sum_{p=0}^P q \beta_{qp} b^q a^p \quad (17)$$

$$C_A = a + \sum_{q=1}^Q \sum_{p=1}^P P [B_q A_p] = a + \sum_{q=1}^Q \sum_{p=1}^P P \beta_{qp} a^p b^q \quad (18)$$

where P and Q are the maximum values of p and q found in complexes of B and A respectively. The total concentrations of A and B are known either from amounts of materials used to make up the starting solutions or from analytical information. The formation constants β_{qp} of equations (17) and (18) can be determined if a and b can be determined. From equations (17) and (18) the average number of ligands bound per metal ion is

$$\bar{n} = \frac{C_A - a}{C_B} = \frac{\sum_{q=1}^Q \sum_{p=1}^P p \beta_{qp} b^q a^p}{\sum_{q=1}^Q \sum_{p=0}^P q \beta_{qp} b^q a^p} \quad (19)$$

The ratio of total metal ion concentration to free (unhydrolyzed) metal ion concentration is

$$F_0 = \frac{C_B}{b} = \sum_{q=1}^Q \sum_{p=0}^P q \beta_{qp} b^{q-1} a^p \quad (20)$$

For polynuclear complexes, \bar{n} and F_0 are functions of ligand concentration, a, and free metal ion concentration, b. The quantities \bar{n} (a) and F_0 (a) can be calculated using (19) and (20) for a series of values of C_B for those systems in which corresponding values of C_A , C_B , a, and b may be measured.

A. The Method of Jensen (17).

The equation for \bar{n} is considered where concentration ranges are such that only two complexes can exist at the same time; $B_r A_x$ and $B_q A_y$, where $B_r A_x$ is thought of as the product in a reaction of $B_q A_y$ with B and A so that $x/r > y/q$ or the ratio A to B has increased. From this and the defining equation for \bar{n} is derived

$$\frac{\bar{n}}{\frac{x-\bar{n}}{r}} = \frac{r[B_r A_x]}{q[B_q A_y]} \quad (21)$$

Two limiting cases are considered for this simplified relationship for \bar{n} . The first is for $x/r \gg \bar{n}$. For this case, the relationships

$$\left(\frac{\partial \log C_B}{\partial \log a} \right) \bar{n} = \frac{qx-ry}{q-r} = \frac{X - \frac{r}{q} y}{1 - \frac{r}{q}} = t \quad (22)$$

and

$$\left(\frac{\partial \log \bar{n}}{\partial \log a} \right) C_B = \frac{qx-ry}{q} = X - \frac{r}{q} y = m \quad (23)$$

are derived. This condition is equivalent to a case where a complex $B_r A_x$ dominates the expression for the concentration of A bound in complexes with B and at the same time a complex $B_q A_y$ dominates the expression for C_B . A plot of $\log \bar{n}$ vs $\log C_B$ or a plot of $\log \bar{n}$ vs

log a is linear. The values of m and t may be determined from the slopes of the experimental curves. From the two equations thus obtained, only two independent ratios among the four variables r, q, x, and y can be determined.

The second case applies for the condition $x'/r' \simeq n$. For this case, similar equations are derived and define slopes m' and t' .

The equations define an infinite number of sets of values (r,x), (q,y), (r',x'), and (q',y'). Independent data are needed to choose proper values although the ratios are determined. The advantage of this method is that, in many cases, the primary hydrolysis product can be determined with certainty.

B. Core-Links Treatment (18,19,20)

Sillén (18) has shown that a large number of complexes produced on hydrolysis may be considered to be made up of a central group, the core, held to a number of composite groups, the links, containing the central group. These are represented as $B_t(A_tB)_m$. These polynuclear complexes give parallel curves for plots of m/n vs $(\log a)_{C_B}$ or $\log F_0$ vs $(\log a)_{C_B}$. The spacings between the curves can be shown to be

$$t = - \left(\frac{\log C_B}{\log a} \right) \frac{1}{n} \quad (24)$$

From the definitions of C_B and C_A , the function $\log(C_B/b)$ is

$$\log C_B b^{-1} = \log [1 + \sum q a^p b^{q-1} \beta_{pq}] \quad (25)$$

The total concentrations of A and B are known, as described in the discussion of mononuclear complexes and a and b are measured as C_B is held constant while varying C_A within the maximum limits possible without precipitation of A metal hydroxide. Sillén (18) has shown that if the plots of m/n vs $\log a$ with C_B held constant are parallel and the amount of free B is not negligible compared to C_B , all the complexes present in appreciable amounts must have the general formula $B(A_tB)_m$ where t is given by equation (24). It has also been shown that if the curves from a plot of $\log(C_B/b)$ vs $\log a$ at constant C_B are parallel, the formula for the complexes must be $B(A_tB)_m$ with t constant and m variable. If the curves also coincide, only mononuclear complexes are present since the extent of hydrolysis would be a function of "a" only in this instance.

C. Hedström's Method (21)

In this method, no assumption whatsoever is made regarding the nature of the complexes formed. By elimination of p, q, and β_{pq} , the general relationship is obtained as

$$\left(\frac{\partial C_B}{\partial \ln a} \right)_b = \left(\frac{\partial C_A}{\partial \ln b} \right)_a \quad (26)$$

This applies to one or more of the BA_p complexes formed. This equation is subject to the boundary conditions $\ln(a/C_A) = 0$ at $C_B = 0$ and $\ln(b/C_B) = 0$ at $C_A = 0$. Solving equation (26) for $\log(a/C_A)$ leads to

$$\log(a/C_A) = \left[\int_0^{C_B} \left(\frac{\partial \log(b/C_B)}{\partial C_A} \right)_{C_B} dC_B \right]_{C_A} \quad (27)$$

Similarly

$$\log(b/C_B) = \left[\int_0^{C_A} \left(\frac{\partial \log(a/C_A)}{\partial C_B} \right)_{C_A} dC_A \right]_{C_B} \quad (28)$$

For each series of experiments at constant C_B , one plots $\log(b/C_B)$ against C_A and determines the derivative of these curves at certain C_A values. The corresponding value of the $\log(a/C_A)$ is obtained from the area under the curve obtained when $\left(\frac{\log b/C_B}{C_A} \right)_{C_B}$ is plotted

against C_B . A similar operation is used for the second equation. This method necessitates measurement at a large number of different values of C_B (or C_A if the other equation is used). A wide range of concentrations should be used.

The evaluation of formation constants is accomplished by use of

$$F_1 = \frac{C_E - b}{a \cdot b} = \sum_p \beta_{p,q} a^{p-1} b^q \quad (29)$$

When $q = 1$, F_1 is a function of "a" only, which means the complexes are mononuclear and the problem is the same as formulated by Leden for the mononuclear case. To evaluate the degree (maximum q) of the polynuclear process, a plot of F_1 vs b at certain constant values of a is made. If a straight line results, only mononuclear and dinuclear complexes are present. If a parabola results, trinuclear complexes are present, etc. From the plot of F_1 vs b , the intercept

$$I_1 = \sum_p \beta_{p,1} a^{p-1} \text{ is obtained.}$$

Now let

$$F_2 = \frac{F_1 - I_1}{ab} \quad (30)$$

be plotted against b . The next intercept $I_2 = \sum_p \beta_{p,2} a^{p-2}$ is obtained. This process is repeated until a straight line is obtained. For each value of p , q can have more than one value. In order to find all the formation constants, the F curves obtained at different constant values of a are compared. The resulting system of equations will enable calculation of the β_{qp} 's. This method will be illustrated for a specific case.

The first part of the paper is devoted to a study of the properties of the function $f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt$ for $n \geq 1$. It is shown that $f(x)$ is a polynomial of degree $n-1$ if $\phi(t)$ is a constant function. In the case where $\phi(t)$ is a linear function, $f(x)$ is a polynomial of degree n . More generally, if $\phi(t)$ is a polynomial of degree m , then $f(x)$ is a polynomial of degree $n+m$.

$$f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt = \int_0^x (x-t)^{n-1} (a_0 + a_1 t + \dots + a_m t^m) dt$$

$$f(x) = \int_0^x (x-t)^{n-1} (a_0 + a_1 t + \dots + a_m t^m) dt = \int_0^x (x-t)^{n-1} a_0 dt + \dots + \int_0^x (x-t)^{n-1} a_m t^m dt$$

The first part of the paper is devoted to a study of the properties of the function $f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt$ for $n \geq 1$. It is shown that $f(x)$ is a polynomial of degree $n-1$ if $\phi(t)$ is a constant function. In the case where $\phi(t)$ is a linear function, $f(x)$ is a polynomial of degree n . More generally, if $\phi(t)$ is a polynomial of degree m , then $f(x)$ is a polynomial of degree $n+m$.

The second part of the paper is devoted to a study of the properties of the function $f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt$ for $n \geq 1$. It is shown that $f(x)$ is a polynomial of degree $n-1$ if $\phi(t)$ is a constant function. In the case where $\phi(t)$ is a linear function, $f(x)$ is a polynomial of degree n . More generally, if $\phi(t)$ is a polynomial of degree m , then $f(x)$ is a polynomial of degree $n+m$.

$$f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt = \int_0^x (x-t)^{n-1} (a_0 + a_1 t + \dots + a_m t^m) dt$$

The third part of the paper is devoted to a study of the properties of the function $f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt$ for $n \geq 1$. It is shown that $f(x)$ is a polynomial of degree $n-1$ if $\phi(t)$ is a constant function. In the case where $\phi(t)$ is a linear function, $f(x)$ is a polynomial of degree n . More generally, if $\phi(t)$ is a polynomial of degree m , then $f(x)$ is a polynomial of degree $n+m$.

$$f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt = \int_0^x (x-t)^{n-1} (a_0 + a_1 t + \dots + a_m t^m) dt$$



The fourth part of the paper is devoted to a study of the properties of the function $f(x) = \int_0^x (x-t)^{n-1} \phi(t) dt$ for $n \geq 1$. It is shown that $f(x)$ is a polynomial of degree $n-1$ if $\phi(t)$ is a constant function. In the case where $\phi(t)$ is a linear function, $f(x)$ is a polynomial of degree n . More generally, if $\phi(t)$ is a polynomial of degree m , then $f(x)$ is a polynomial of degree $n+m$.

TYPICAL HYDROLYTIC PROCESSES

A. Hg^{2+}

The hydrolysis of Hg^{2+} has been studied by measuring the solubility of HgO in water (22) at 25° . From the value of the solubility product of $\text{Hg}(\text{OH})_2$ (using the value of $\text{pK}_w = 14$), the sum $\text{pKa}_1 + \text{pKa}_2$ was calculated to be 6.26.


A more recent and much more thorough study was made by Hietanen and Sillén (23) using emf measurements. The process was found to be one in which \bar{n} is a function of $\log h$ only and independent of C_B . From the discussion of the treatment of data, this was seen to be the criterion for the formation of mononuclear complexes only. An analysis of the data by the methods described shows the main products of the hydrolysis to be $\text{Hg}(\text{OH})_2$ and HgOH^+ corresponding to the reactions






The k 's are the first and second acid constants of the acid $\text{Hg}(\text{H}_2\text{O})_n^{2+}$.

B. Fe^{3+}


The hydrolysis of Fe^{3+} has been studied for many years by a variety of techniques and a variety of results have been obtained (24,26. See references 1 and 26 for complete references to the earlier literature). Hedstrom (26) has studied the hydrolysis of Fe^{3+} using emf measurements. The free hydrogen ion concentrations were determined by a glass electrode, and the concentration of Fe^{3+} from the half cell $\{\text{Fe}^{3+}, \text{Fe}^{2+}\} \text{Pt}$. The reference electrode used was a calomel electrode,



The hydrolysis of Fe^{2+} has been shown to be negligible in solutions of $\text{pH} < 5$ (27). The following species were identified from an analysis of the data.



The predominant species present is the dinuclear complex with the two mononuclear species becoming more important at low total concentrations of $\text{Fe}(\text{III})$. The dinuclear complex was presumed to be



The first part of the investigation was devoted to the study of the influence of the temperature of the medium on the rate of the reaction. The results are given in Table I.

The rate of the reaction was determined by the method of initial rates. The concentration of the reactants was kept constant and the temperature was varied. The results are given in Table I.

$$k = A e^{-E/RT}$$

$$\ln k = \ln A - E/RT$$

From the slope of the straight line obtained from the plot of $\ln k$ versus $1/T$, the activation energy E was determined to be 12,500 cal/mole.

The second part of the investigation was devoted to the study of the influence of the concentration of the reactants on the rate of the reaction. The results are given in Table II.

$$r = k[A]^m[B]^n$$

From the slope of the straight line obtained from the plot of $\ln r$ versus $\ln [A]$, the order of the reaction with respect to A was determined to be 1.5.

$$r = k[A]^{1.5}[B]^n$$


$$\ln r = \ln k + 1.5 \ln [A] + n \ln [B]$$

$$\ln r - 1.5 \ln [A] = \ln k + n \ln [B]$$

From the slope of the straight line obtained from the plot of $\ln r - 1.5 \ln [A]$ versus $\ln [B]$, the order of the reaction with respect to B was determined to be 0.5.

$$r = k[A]^{1.5}[B]^{0.5}$$

In the conductometric titration of Fe^{3+} with base (25), the first inflection point could not be attributed to the formation of $\text{Fe}(\text{OH})_3$ but rather some intermediate complex. This intermediate was separated and the composition found to be $[\text{Fe}_4\text{O}_3(\text{OH})_5]\text{NO}_3 \cdot 4\text{H}_2\text{O}$. The structure suggested was



where An represents a univalent anion. This structure is reported to be similar to some of the α -series minerals of iron.

The hydrolysis products $\text{Fe}(\text{H}_2\text{O})_6^{3+}$, $\text{Fe}(\text{H}_2\text{O})_5\text{OH}^{2+}$, $\text{Fe}(\text{H}_2\text{O})_4(\text{OH})_2^{4+}$, and $\text{Fe}(\text{H}_2\text{O})_3(\text{OH})_3^{6+}$ have also been reported on the basis of similar base additions (24).

C. Bi^{3+}

The hydrolysis of Bi^{3+} has been studied by Olin (28,29) using a glass electrode to determine the free concentration of H^+ and a bismuth amalgam electrode to measure the Bi^{3+} concentration. The first step in the hydrolysis leads to $\text{Bi}_6(\text{OH})_{12}^{6+}$. In solutions of lower pH, this forms larger aggregates. The data can be interpreted in terms of the formation of $\text{Bi}_9(\text{OH})_{20}^{7+}$, $\text{Bi}_9(\text{OH})_{21}^{6+}$, and $\text{Bi}_9(\text{OH})_{22}^{5+}$ as the polynuclear species present in the aggregate.

Previously, Graner and Sillén (30) had interpreted the data obtained from similar experiments in terms of the formation of $\text{Bi}_n(\text{OH})_{2n}^{(3+n)+}$ written in core-links form. The identity of this product was determined by studying a concentration range of Bi^{3+} from 10 to 50 mM. The study of Olin (28) was made over a range of 0.1 to 50 mM. The quinhydrone electrode was used to measure the concentration of H^+ and was later found to have appreciable basicity at the concentrations of H^+ used (31). The formula for this product would be written as $\text{Bi}_{n+1}(\text{OH})_{2n}^{(3+n)+}$ in conventional form and is seen not to be the product found by Olin. This is a typical example of the difficulties encountered and the conclusions drawn in the study of such systems.

REFERENCES

1. J. Bjerrum, G. Schwarzenbach and L. G. Sillén, "Stability Constants," Part II. The Chemical Society, London, 1958. Chem. Soc. Spec. Publ. No. 7.
2. J. Bjerrum, "Metal Ammine Formation in Solution," P. Haase and Son, Copenhagen, 1941.
3. I. Leden, Z. physik. Chem. A188, 160 (1941).
4. S. Fronaeus, Acta Chem. Scand. 4, 72 (1950).
5. J. C. Sullivan and J. C. Hindman, J. Am. Chem. Soc. 74, 6091 (1952).
6. L. G. Sillén, Acta Chem. Scand. 10, 186 (1956).
7. H. S. Simms, J. Am. Chem. Soc. 48, 1239 (1926).
8. G. Schwarzenbach and H. Ackerman, Helv. Chem. Acta. 31, 1029 (1948).
9. G. Schwarzenbach and R. Baur, *ibid.* 39, 722 (1956).
10. G. Schwarzenbach and A. Willi and R. Baur, *ibid.* 30, 1303 (1947).
11. H. Irving and H. Rossotti, J. Chem. Soc. 1953, 3397.
12. D. Dyrssen and L. G. Sillén, Acta Chem. Scand. 7, 663 (1953).
13. F. Rossotti and H. Rossotti, J. Phys. Chem. 63, 1041 (1959).
14. F. Rossotti, H. Rossotti and L. Sillén, Acta Chem. Scand. 10, 203 (1956).
15. G. Anderegg, Helv. Chem. Acta 45, 901 (1962).
16. J. Sullivan, J. Rydberg, and W. Miller, Acta Chem. Scand. 13, 2023 (1959).
17. B. S. Jensen, Acta Chem. Scand. 15, 487 (1961).
18. L. G. Sillén, Acta Chem. Scand. 8, 299, 318 (1954).
19. S. Hietanen and L. G. Sillén, *ibid.* 8, 1607 (1954).
20. F. Rossotti and H. Rossotti, *ibid.* 10, 957 (1956).
21. B. Hedstrom, *ibid.* 9, 613 (1955).
22. A. Garret and A. E. Hirschfelder, J. Am. Chem. Soc. 60, 299 (1938).
23. S. Hietanen and L. G. Sillén, Acta Chem. Scand. 6, 747 (1952).
24. W. Feitknecht and W. Michaelis, Helv. Chem. Acta 45, 212 (1962).
25. O. E. Zvyagintsev and Y. S. Lopatto, Zhur. Neorg. Khim 6, 863 (1961).
26. B. Hedström, Arkiv Kemi 6, 1 (1953).
27. B. Hedström, *ibid.* 5, 457 (1953).
28. A. Olin, Acta Chem. Scand. 11, 1445 (1957).
29. A. Olin, *ibid.* 13, 1791 (1959).
30. F. Graner and L. G. Sillén, *ibid.* 1, 631 (1947).
31. G. Biedermann, *ibid.* 10, 1340 (1956).

MOLECULAR ORBITAL TREATMENT OF POLYHALOGEN AND RELATED SPECIES

Carolyn M. McArthur

May 28, 1963

INTRODUCTION

The halogens interact to form a wide variety of species besides the familiar halogen diatomic molecules X_2 . These can be classified as the inter-halogen compounds XY , XY_3 (T-shaped, C_{2v}), XY_5 (square pyramidal, C_{4v}), XY_7 (pentagonal bipyramidal, D_{5h}); the polyhalide ions X_3^- (linear), I_n^- (n odd), e.g., I_5^- (V-shaped, C_{2v}), I_8^{2-} (chair shape, C_{2h}), XY_2^- and XYZ^- (both linear), XY_4^+ (square planar, D_{4h}); and the positive polyhalogen ions XY_2^+ (angular, C_{2v}) and XY_4^+ (structure unknown).

Many of these polyhalogen species, if described in valence bond terms, violate the Lewis octet rule of valency theory (1) with over-filled electron shells. The common solution has been to use an outer d -orbital to provide the necessary extra orbital(s), but this d -orbital does not belong to the valence orbitals. Nevertheless, hybrid valence structures using outer d -orbitals have been postulated, with lone pairs of electrons assumed to occupy one or more of the hybrid orbitals. Explanations involving electron-pair repulsions have been invoked to correlate the possible structures predicted by hybridization considerations with the actual shape of molecules like T-shaped ClF_3 . (See, for example, Gillespie and Nyholm (2).) However, even Pauling (3) has sometimes been cautious in describing structures in terms of outer d -orbitals, noting that resonating Lewis electronic structures probably make a considerable contribution to the electronic ground state of such molecules.

The placing of lone pairs of electrons in high energy hybrid orbitals is a significant objection to the valence bond description using outer d -orbitals. Only in IF_7 are all the valence shell electrons of iodine involved in bonding. Inherent in this description is the assumption of easy orbital excitation from a p to a d level when the principal quantum number remains unchanged, but this assumption has no basis in experimental fact. For example, it is often assumed that compounds of F and Cl differ in structural properties because of d -hybridization in chlorine. In fluorine, the energy of the center of gravity of states arising from $2s^2 2p^4(^3F) 3d$ is 15.9 e.v. (4); that arising from $3s^2 3p^4(^3P) 3d$ is 11.2 e.v. (5,6). In both F and Cl the excitation energy is large, in each case being only slightly less than 2 e.v. below the ionization limit, and Klemperer (7) points out that it is extremely arbitrary to single out the configuration of Cl involving a $3d$ electron from the wealth of configurations near the ionization limit. The selection of a single excited configuration as predominant is only useful if this configuration lies uniquely lower in energy than others.

A striking structural feature of these compounds is that, with the exception of IF_7 , the bond angles are all very close to 90° or 180° . In 1951, Fimentel proposed a bonding scheme for trihalide ions based on molecular orbitals formed from linear combinations of $n\text{p}_\sigma$ -halogen orbitals, which excludes the need for higher energy orbitals (8). Hach and Rundle (9) independently proposed the same type of MO scheme, but applied it only to the polyiodides. Havinga and Wiebenga have extended the LCAO-MO method to all the polyhalogen species and have obtained a consistent qualitative explanation of bond angles, atomic configurations, and bond lengths (10, 11).

MO BONDING SCHEME FOR POLYHALOGENS

In the modified Hückel LCAO-MO treatment used by Wiebenga and co-workers, only atomic p-orbitals of the halogens are considered. Overlap integrals are neglected, and all resonance integrals are taken as zero, except those involving orbitals of neighboring atoms in the bond direction, which are given the same value β (Hückel approximations (12)).

If ϕ_i is the p-function of atom \underline{i} , then the Coulomb integral is:

$$\alpha = \int \phi_i^* H \phi_i d\tau$$

and the resonance integral is:

$$\beta = \int \phi_i^* H \phi_{\underline{i+1}} d\tau$$

The molecular orbitals are then defined by:

$$\Psi_j = \sum_i c_{ji} \phi_i ; \quad j = 0, 1, 2, \dots$$

and the bond order between atoms \underline{l} and \underline{m} is:

$$P_{lm} = \left| \sum_j n_j c_{jl} c_{jm} \right|$$

where the sum is over the occupied MO's labeled by \underline{j} .

The charge density at atom \underline{i} is:

$$q_i = \sum_j n_j (c_{ji})^2$$


where n_j is the number of electrons in level \underline{j} .

The calculations are made in two approximations: (a) as a zero approximation, all Coulomb integrals α are supposed to have the same value, regardless of the nature of the halogen atoms and their formal charges; (b) in a first

approximation, different Coulomb integrals are assigned to different atoms and to the same atoms with different formal charges: (i) For I, Br, Cl, and F the values α , $\alpha + 0.2\beta$, $\alpha + 0.4\beta$, and $\alpha + 1.0\beta$ are taken, respectively; the assumed differences in these values are approximately proportional to the differences in electronegativities of the atoms, with a proportionality constant of $2/3k$. (ii) A formal charge q , calculated for an atom in zero approximation, increases the absolute value of its Coulomb integral by $2/3k \times 0.3q = 0.2qk$. (This means that, following Pullman and Pullman (13), a charge q is taken to be equivalent to an increase of electronegativity of $0.3q$.) The first approximation is only applied when all bond orders in the part of the molecule being considered are equal to each other in zero approximation.

EXAMPLE: CALCULATIONS FOR IBrCl⁻

The only configurations in which the p-functions overlap satisfactorily for any triatomic molecule are the rectangular and linear.



For the rectangular form, the functions which combine are the p_x -functions of atoms A and B and the p_y -functions of atoms B and C. This leads to twice the secular equation:

$$\begin{vmatrix} \alpha - E & \beta \\ \beta & \alpha - E \end{vmatrix} = 0$$

Solutions: $E = \alpha + \beta$
 $E = \alpha - \beta$

There are 16 p electrons to be accommodated in the two MO's of energy $\alpha + \beta$, the two MO's of energy $\alpha - \beta$, and the five remaining atomic p-orbitals of energy α . One of the orbitals of energy $\alpha - \beta$ will remain empty, and the energy of the 16 electrons is $16\alpha + 2\beta$. Since this is the same as the


energy calculated for a halogen molecule and a halide ion separated from each other, no gain of energy is predicted for this angular negative trihalide ion.

For the linear form, the p_x -functions of atoms A, B, and C combine, leading to the secular equation:


$$\begin{vmatrix} \alpha - E & \beta & \beta \\ \beta & \alpha - E & 0 \\ 0 & \beta & \alpha - E \end{vmatrix} = 0$$

Solutions: $E_1 = \alpha - \beta\sqrt{2}$
 $E_2 = \alpha$
 $E_3 = \alpha + \beta\sqrt{2}$


The total energy of the 16 electrons is $16\alpha + 2\sqrt{2}\beta$, and a gain of energy of 0.83β is calculated compared with X_2 and X^- . Bond orders and atomic charges in zero approximation are:



The results in first approximation for the three possible linear configurations of the IBrCl^- ion are:



The first configuration, which is actually observed, is calculated to have the lowest energy. Bond order and atomic charges in first approximation are:



The most electronegative atom acquires the highest negative charge.

EXPERIMENTAL SUPPORT FOR MO DESCRIPTION OF BONDING

For the X_3^- ion, the MO's are represented in the following way in the zero approximation:

$$\begin{array}{l} \text{(anti-bonding)} \quad \Psi_1 = \frac{1}{2}(p_A + p_C) + \frac{1}{\sqrt{2}} p_B \\ \text{(non-bonding)} \quad \Psi_2 = \frac{1}{\sqrt{2}} (p_A + p_C) \\ \text{(bonding)} \quad \Psi_3 = \frac{1}{2} (p_A + p_C) - \frac{1}{\sqrt{2}} p_B \end{array}$$

The first part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The second part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The third part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The fourth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

1874 1875
1876 1877
1878 1879
1880 1881

The fifth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The sixth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The seventh part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

1882 1883
1884 1885
1886 1887
1888 1889

The eighth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The ninth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.



The tenth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

The eleventh part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.


The twelfth part of the document is a list of names and addresses, which are arranged in a columnar format. The names are written in a cursive hand, and the addresses are written in a more formal, printed style.

Of the 16 p electrons, 12 occupy the six p-orbitals not involved in bonding. Two are in the non-bonding orbital, which places electron density at atoms A and C. If the electronegativities of A, B, and C are the same, about half the electron density of the bonding orbital will belong to B, one quarter to each A and C, and half the non-bonding pair will belong to each of A and C. Since the non-bonding orbital puts extra negative charge on the terminal atoms, it can be correlated with the observation that the central atom of a trihalide is always less electronegative than the terminal atoms. Since only one bonding electron pair is used for the two bonds, this accounts in a natural way for the very considerable elongation of these bonds over the distance found in I₂ (14). Thus, the I-I bond distance in Cl_4AsI_3 is 2.90 Å., whereas twice the covalent radius of iodine is 2.68 Å.

The MO method predicts the correct bond angles, atomic configurations, and order of bond lengths for all polyhalogens except IF₇. These results are tabulated in reference 10. Only one more example, the I₅⁻ ion, will be listed here. The delocalization energies are given along with the configurations.



In zero approximation, three models are energetically equivalent, but in first approximation the configuration actually observed in N(CH₃)₄I₅ (9) has the lowest energy.





Calculated bond orders

Observed bond lengths

The nuclear quadrupole coupling constants have been measured by Cornwell and Yamasaki for both Cl³⁵ (15) and

I^{127} (16) in alkali salts of ICl_2^- and ICl_4^- . The results can be most easily interpreted by translating the MO description of these ions into their equivalent description in terms of resonating valence bond structures.



The value of the coupling constant, eQq/h , for Cl^{35} in ICl_2^- and in ICl_4^- is about half of that found in ICl . The I^{127} values are roughly the same in all three species. The conclusions based on these results are that the I-Cl bonds in the two ions have a rather large proportion of ionic character, corresponding to the presence of a net negative charge of roughly half an electron on each Cl atom. Since hybridization involving promotion from the $5p$ to the $5d$, or possibly $6s$, level would raise the iodine resonance frequencies markedly from the value found in ICl , the attainment by iodine of higher valencies in these ions is due more to transfer of charge to chlorine atoms than to promotion to higher orbitals of iodine. These results substantiate the MO description of bonding and show conclusively that d orbitals cannot be included in the bonding scheme to the extent required by the spd hybridization model. These same results were obtained independently by Japanese workers who also measured the Cl^{35} resonances in these ions. The estimated ionic nature of the I-Cl bond in ICl_2^- was 66%; and in ICl_4^- , it was 59%. The values they calculated by the MO method were 61% and 57%, respectively, in good agreement with observations (17).

The infrared and Raman spectra of ICl_2^- , ICl_4^- , $BrCl_2^-$, and Br_3^- have been obtained and the stretching force constants calculated (18). The values of f_r , the stretching force constant, for the trihalide ions are roughly half the values for the free halogens, and the values of the interaction force constants f_{rr} between bond stretching coordinates at an angle of 180° to each other are very large (approximately 35% of the value of f_r). These rather unusual force constants have been interpreted in terms of bonding using p orbitals. The similarity between these force constants and those for HF_2^- are pointed out.

1947
The following information is for your information
and is not to be used for any other purpose
without the express written consent of the
Department of the Interior.

U.S. DEPARTMENT OF THE INTERIOR

$$\left| \frac{1}{n} \sum_{i=1}^n x_i \right|$$

1947

The following information is for your information
and is not to be used for any other purpose
without the express written consent of the
Department of the Interior.

The following information is for your information
and is not to be used for any other purpose
without the express written consent of the
Department of the Interior.

Table I. Bond Stretching Constants for Some Polyhalogen Ions and Molecules.

Ion	f_r (md/Å)	f_{rr} (md/Å)	$f_{rr'}$ (md/Å)	$\frac{f_r(xy)}{f_r(xy)}$	f_{rr}/f_r
ICl_2^-	1.00	0.36		0.46	0.36
ICl_4^-	1.25	0.33	0.08	0.53	0.26
ICl	2.38				
BrCl_2^-	1.29	0.26		0.47	0.20
BrCl	2.83				
Br_3^-	0.91	0.32		0.38	0.36
Br_2	2.45				
HF_2^-	2.31	1.72		0.23	0.72
HF	10				
CO_2	15.5	1.3		0.82	0.08
CO	19.0				

An MO treatment in terms of chains of I_3^- has been applied successfully to explain the metallic nature of the starch-iodine complex, including the measured weak paramagnetism and electronic conductivity (19). In terms of the MO theory, a row of iodine atoms (50 to 500) behaves as a one-dimensional metal with a band of energy levels to accommodate the 5p-electrons of the I_3^- chain. The optical absorption properties are also consistent with the metallic model. The 6250 Å absorption may be ascribed to a transition to the σ band from a π band. The model predicts, and thus explains, the electron spin resonance of the complex in the right order of magnitude (about 10^{16} unpaired electrons per gram).

Bersohn (20) has employed delocalized molecular orbitals to explain intermolecular bonding in solid iodine. The band structure of solid iodine is worked out on the assumption that electron exchange between layers of iodine atoms is negligible and that only p-functions are used by the iodine atoms. Just as the halogen molecule is formally analogous to ethylene in MO description, so is the solid halogen to graphite. Each halogen atom is assumed to interact with three neighboring atoms in the plane. The model is justified


by its simple description of the iodine nuclear quadrupole coupling constant, the anisotropy of the electrical conductivity, and the optical absorption.

The MO calculations can be used in a qualitative way to explain some thermodynamic properties of the polyhalogens. Qualitative agreement between theory and experiment is obtained when the energies of formation calculated in first approximation for simple interhalogens are compared with experimental values. (ΔH^\ddagger refers to the formation of gaseous XY from gaseous X_2 and Y_2 at 25°C. and 1 atm. pressure.)

Table II. Calculated and Observed Heats of Formation (11)

<u>Compound</u>	<u>ΔE (calc.)</u>	<u>ΔH^\ddagger (obs.) kcal/mole</u>
IF	0.24 β	-30.1
BrF	0.15 β	-18.4
ClF	0.09 β	-13.4
ICl	0.04 β	- 3.35
IBr	0.01 β	- 1.42
BrCl	0.01 β	- 0.21

Compounds containing $BrICl^-$ ion decompose into a monochloride and BrI. This is partially ascribed to the high lattice energy of the crystalline monochloride formed, as compared with the energy of the corresponding bromide or iodide. The reaction energy ΔE as calculated by the MO method for the three possible decompositions of the free $BrICl^-$ ion is:



The fact that the calculated dissociation energy is smallest for the first reaction may be an additional explanation for the experimentally observed method of dissociation (11).

The existence of the xenon fluorides XeF_2 , XeF_4 , and XeF_6 also supports the validity of the MO description of bonding in the polyhalogens. The compounds XeF_2 and XeF_4 are isoelectronic with ICl_2 and ICl_4 , respectively. In fact,

Pimentel (8) predicted in 1951 that a rare gas could form complexes with halogens on the basis of his MO treatment of the polyhalogens. The compound XeF_2 is linear (21), and XeF_4 is square planar, with the average distance between Xe and F being 1.92 Å. (22) Rundle (23) argues that because of the especially high promotional energies of the rare gases, the bonding in XeF_2 and XeF_4 has little possibility of using outer d-orbitals; so the existence of these xenon fluorides in itself is evidence in favor of delocalized bonds using only p-orbitals of the valence shell.

Smith has observed the infrared spectrum of XeF_2 and has interpreted the results to show that the bond length is too short and the interaction force constant too small for an explanation of the bonding in terms of p orbitals (21). However, Pimentel and Spratley (24) argue that the IR data can be reasonably interpreted in terms of an MO bonding description, since the molecules XeF_2 and ICl_2 involve quite different formal charge distributions. In XeF_2 , the xenon atom must have a significant positive formal charge to balance the negative charge placed on the terminal atoms, whereas the iodine atom has a formal charge close to zero. This difference can be expected to strengthen the Xe-F bond because of the electrostatic attractions. At the same time, the interaction force constant should be reduced, because an asymmetric displacement is no longer favored energetically over a symmetric displacement. In the symmetric mode, the terminal atom repulsions that tend to raise the energy are counteracted by the central atom positive charge.

CONCLUSION

There is a considerable amount of evidence, both structural evidence and that based on other physical methods which give information concerning bonding in molecules, that a molecular-orbital description involving delocalized bonds formed from atomic p-orbitals is a much closer approximation to the true bonding in polyhalogen species than the description using outer d-orbitals in an spd hybrid model.

REFERENCES

1. G. N. Lewis, J. Am. Chem. Soc., 38, 762 (1916).
2. R. J. Gillespie and R. S. Nyholm, Quart. Revs. 11, 339 (1957).
3. L. Pauling, The Nature of the Chemical Bond, 3rd ed., Cornell Univ. Press, New York, 1960, p. 178.
4. C. E. Moore, "Atomic Energy Levels," Vol. I., U. S. Dept. of Commerce, 1949, Washington, D. C.
5. C. J. Humphreys and E. Paul, Jr., J. Opt. Soc. Am., 49, 1186 (1959).
6. L. Minnhagen, J. Opt. Soc. Am., 51, 298 (1961).

7. W. Klemperer, J. Am. Chem. Soc., 83, 3910 (1961).
8. G. C. Pimentel, J. Chem. Phys., 19, 446 (1951).
9. R. J. Hach and R. E. Rundle, J. Am. Chem. Soc., 73, 4321 (1951).
10. E. E. Havinga and E. H. Wiebenga, Rec. trav. chim., 78, 724 (1959).
11. E. H. Wiebenga, E. E. Havinga, and K. H. Boswijk, Advances in Inorg. Chem. Radiochem. (Academic Press) 3, 133 (1961).
12. W. Hückel, Z. Physik, 76, 628 (1932).
13. B. Pullman and A. Fullman, Les theories electroniques de la chimie organique. Paris: Masson and Co., 1952.
14. R. E. Rundle, Record Chem. Progr. (Kresge-Hooker Sci. Lib.), 23, 195 (1962).
15. C. D. Cornwell and R. S. Yamasaki, J. Chem. Phys., 27, 1060 (1957).
16. R. S. Yamasaki and C. D. Cornwell, J. Chem. Phys., 30, 1265 (1959).
17. Y. Jurita, D. Nakamura, and N. Hayakawa, J. Chem. Soc. Japan, Pure Chem. Sec., 79, 1093 (1958). C.A. 53, 54f (1959).
18. W. B. Person, G. R. Anderson, J. N. Fordemwalt, H. Stammreich, and R. Forneris, J. Chem. Phys., 35, 908 (1961).
19. R. Bersohn and I. Isenberg, J. Chem. Phys., 35, 1640 (1961).
20. R. Bersohn, J. Chem. Phys., 36, 3445 (1962).
21. D. F. Smith, J. Chem. Phys., 38, 270 (1963).
22. J. A. Ibers and W. C. Hamilton, Science, 139, 106 (1963).
23. R. E. Rundle, J. Am. Chem. Soc., 85, 112 (1963).
24. G. C. Pimentel and R. D. Spratley, J. Am. Chem. Soc., 85, 826 (1963).
25. R. E. Rundle, Acta Cryst., 14, 585 (1961).

UNIVERSITY OF ILLINOIS-URBANA

546L61 C001
INORGANIC SEMINAR ABSTRACTS URBANA
1962/63



3 0112 025503050